

Astrophysics and Space Science Library 434

Johan H. Knapen
Janice C. Lee
Armando Gil de Paz *Editors*



Outskirts of Galaxies

AS
SL

 Springer

Outskirts of Galaxies

Astrophysics and Space Science Library

EDITORIAL BOARD

Chairman

W. B. BURTON, *National Radio Astronomy Observatory, Charlottesville, Virginia, U.S.A. (bburton@nrao.edu); University of Leiden, The Netherlands (burton@strw.leidenuniv.nl)*

F. BERTOLA, *University of Padua, Italy*

C. J. CESARSKY, *Commission for Atomic Energy, Saclay, France*

P. EHRENFREUND, *Leiden University, The Netherlands*

O. ENGVOLD, *University of Oslo, Norway*

E. P. J. VAN DEN HEUVEL, *University of Amsterdam, The Netherlands*

V. M. KASPI, *McGill University, Montreal, Canada*

J. M. E. KUIJPERS, *University of Nijmegen, The Netherlands*

H. VAN DER LAAN, *University of Utrecht, The Netherlands*

P. G. MURDIN, *Institute of Astronomy, Cambridge, UK*

B. V. SOMOV, *Astronomical Institute, Moscow State University, Russia*

R. A. SUNYAEV, *Space Research Institute, Moscow, Russia*

More information about this series at <http://www.springer.com/series/5664>

Johan H. Knapen • Janice C. Lee •
Armando Gil de Paz
Editors

Outskirts of Galaxies

 Springer



Editors

Johan H. Knapen
Inst de Astrofísica de Canarias
San Cristobal de la Laguna, Spain

Janice C. Lee
Space Telescope Science Institute
Baltimore, USA

Armando Gil de Paz
Dept. Astrofísica
Universidad Complutense de Madrid
Madrid, Spain

ISSN 0067-0057 ISSN 2214-7985 (electronic)
Astrophysics and Space Science Library
ISBN 978-3-319-56569-9 ISBN 978-3-319-56570-5 (eBook)
DOI 10.1007/978-3-319-56570-5

Library of Congress Control Number: 2017942746

© Springer International Publishing AG 2017

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Cover illustration: This image of the spectacular galaxy NGC 4725 shows evidence of dust lanes in the area surrounding its active nucleus, a bright ring-like region of star formation, and outer spiral arm structure. Credit: DAGAL, Nik Szymanek, SDSS, and S⁴G, www.dagalnetwork.eu.

Printed on acid-free paper

This Springer imprint is published by Springer Nature
The registered company is Springer International Publishing AG
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

Preface

The outskirts of galaxies are mostly unexplored territory. Great advances have been made in particular in studying the star formation (through UV imaging) and gas (HI radio emission), but exploration of the stellar component, observed through optical and infrared imaging, can be considered to be still in its infancy. Yet the outskirts are key to understanding how galaxies form and evolve and how they have diversified into a class of objects with the wide range of morphologies of properties that we observe today. Their importance stems from two facts: the timescales in the outer regions are long, and stellar and gas densities are low. Both lead to slower evolution in the outskirts, implying that we are observing conditions at an earlier state there relative to the denser inner regions of galaxies which have been observed traditionally. In addition, accretion of pristine gas happens in the outskirts, stars are thought to migrate outwards and the material in the outer regions, when seen projected against the emission from background quasars, yields important information about the properties of the interstellar and intergalactic medium.

This volume brings together the views and insights of a number of world-renowned experts in this field, who have written a total of ten chapters summarizing our current knowledge of the outer regions of galaxies, as well as their views on how this field is likely to evolve in the near future. The topics described in detail range from studies of the structure and star formation history of our own Milky Way and the nearest external galaxies on the basis of individual star counts, via the deepest possible imaging of the integrated light of galaxies, to the study of the outskirts of galaxies at cosmological distances through the study of the light of background quasars passing through their outer regions. Other reviews discuss recent observations of molecular and atomic gas in the outskirts of galaxies and what we can learn from those about topics as varied as the current and past star formation and the shape of the dark matter haloes. Observed metallicities and their radial gradients are discussed in the context of chemical composition and star formation in the outskirts, touching on mechanisms such as metal-rich infall and metal mixing in disks. Stellar migration in galaxies is discussed in detail, paying particular attention to how observations and theoretical insights are improving our understanding of galaxy evolution, as is star formation in the outskirts of galaxies, which shines a

new light not just on the properties of the outer regions but also on the process of star formation itself.

Our knowledge of the outer regions of galaxies is rapidly improving because new data are now enabling detailed study at a variety of wavelengths and with a variety of techniques. As the authors of this volume discuss, the next generation of telescopes and instruments will accelerate the exploration of galaxy outskirts, which will without any doubt lead to breakthroughs in our understanding of how galaxies have formed and evolved. We hope that this collection of reviews will provide a resource for a full range of workers in the field—expert investigators in theory and observation, those intrigued by recent discoveries who wish to learn more and students and other researchers who are interested in entering this exciting field.

Acknowledgements

The current volume owes its existence to the research programme developed in the context of the Initial Training Network Detailed Anatomy of Galaxies (DAGAL), funded through the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme FP7/2007–2013/ under REA grant agreement number PITN-GA-2011-289313, and to the partnership with Springer developed within that network. Editors and authors of this book met and developed ideas during the International Astronomical Union Symposium number 321, held in March 2016 in Toledo, Spain, and thank the organizers of that meeting for bringing them together in such a beautiful and stimulating environment.

San Cristobal de La Laguna, Spain
Baltimore, MD, USA
Madrid, Spain
December 2016

Johan H. Knapen
Janice C. Lee
Armando Gil de Paz

Contents

1	Outer Regions of the Milky Way	1
	Francesca Figueras	
1.1	Introduction	1
1.2	The Outer Disk of the Milky Way: Stellar Content	2
1.2.1	Resolved Stellar Populations	3
1.2.2	The Outer Reaches	4
1.3	The Milky Way Outer Disk: Structure and Dynamics	6
1.3.1	Spiral Arm Impact on Disk Dynamics and Structure	6
1.3.2	The Galactic Warp and Flare	7
1.3.3	Gravitational Interaction with Satellites	9
1.3.4	Dynamics of the Vertical Blending and Breathing Modes	10
1.4	Towards a Chemodynamical Model of the Galactic Disk	11
1.4.1	Age-Metallicity-Kinematics Relations	11
1.4.2	The Galactic Thick Disk	12
1.4.3	The Radial Abundance Gradients	13
1.4.4	The “Outside-In” Versus “Inside-Out” Disk Formation Scenarios	17
1.5	Large Surveys in the Next Decade	17
1.5.1	The <i>Gaia</i> Mission	18
1.6	Conclusions	23
	References	23
2	Resolved Stellar Populations as Tracers of Outskirts	31
	Denija Crnojević	
2.1	The Importance of Haloes	31
2.1.1	Resolved Stellar Populations	33
2.1.2	The Low-Mass End of the Galaxy Luminosity Function	34
2.2	Local Group	35
2.2.1	Milky Way	35

2.2.2	M31 (Andromeda)	40
2.2.3	Low-Mass Galaxies In and Around the Local Group	47
2.3	Beyond the Local Group	48
2.3.1	Systematic Studies	50
2.3.2	Panoramic Views of Individual Galaxies	53
2.4	Summary and Future Prospects	61
	References	63
3	The Impact of Stellar Migration on Disk Outskirts	77
	Victor P. Debattista, Rok Roškar, and Sarah R. Loebman	
3.1	Introduction	77
3.1.1	Our Definition of Breaks	79
3.2	Demographics of Profile Type	79
3.2.1	The Role of Environment	80
3.2.2	Redshift Evolution	81
3.3	Stellar Migration	81
3.3.1	Migration via Transient Spirals	82
3.3.2	Multiple Patterns	84
3.3.3	Evidence for Migration in the Milky Way	89
3.4	Type II Profiles	89
3.4.1	Theoretical Models	90
3.4.2	Observational Tests	96
3.4.3	Synthesis and Outlook	99
3.5	Type I Profiles	100
3.5.1	Origin of Type I Profiles in Isolated Galaxies	101
3.5.2	Type I Profiles in Cluster Lenticulars	102
3.6	Type III Profiles	104
3.6.1	Formation of Type III Disk Profiles	104
3.7	Future Prospects	105
	References	107
4	Outskirts of Nearby Disk Galaxies: Star Formation and Stellar Populations	115
	Bruce G. Elmegreen and Deidre A. Hunter	
4.1	Introduction	115
4.2	Outer Disk Structure from Collapse Models of Galaxy Formation	116
4.3	Outer Disk Structure: Three Exponential Types	117
4.4	Outer Disk Stellar Populations: Colour and Age Gradients	118
4.5	Mono-Age Structure of Stellar Populations	120
4.6	Outer Disk Structure: Environmental Effects and the Role of Bulges and Bars	120
4.7	Outer Disk Structure: Star Formation Models	121
4.8	The Disks of Dwarf Irregular Galaxies	125
4.8.1	Radial Trends	125
4.8.2	Star Formation in Dwarfs	128

4.8.3	The $H\alpha$ /FUV Ratio	130
4.8.4	Breaks in Radial Profiles in dIrr Galaxies	132
4.9	Summary	134
	References	135
5	Metallicities in the Outer Regions of Spiral Galaxies	145
	Fabio Bresolin	
5.1	Introduction	145
5.2	Measuring Nebular Abundances	146
5.3	Chemical Abundances of H II Regions in Outer Disks	148
5.3.1	Early Work	148
5.3.2	M83: A Case Study	149
5.3.3	Other Systems	152
5.3.4	Results from Galaxy Surveys	155
5.3.5	Nitrogen Abundances	158
5.4	Additional Considerations	159
5.4.1	Relation Between Metallicity and Surface Brightness Breaks	159
5.4.2	An Analogy with Low Surface Brightness Galaxies?	159
5.5	The Evolutionary Status of Outer Disks	160
5.5.1	Flattening the Gradients	160
5.5.2	Bringing Metals to the Outer Disks	161
5.6	Conclusion	164
	References	165
6	Molecular Gas in the Outskirts	175
	Linda C. Watson and Jin Koda	
6.1	Introduction	175
6.2	Molecular Gas from the Inner to the Outer Regions of Galaxies	177
6.3	Molecular ISM Masses: Basic Equations	178
6.3.1	Brightness Temperature, Flux Density and Luminosity	178
6.3.2	Observations of the Molecular ISM Using CO Line Emission	180
6.3.3	Observations of the Molecular ISM Using Dust Continuum Emission	182
6.3.4	The ISM in Extreme Environments Such as the Outskirts	183
6.4	Molecular Gas Observations in the Outskirts of Disk Galaxies	187
6.4.1	The Milky Way	187
6.4.2	Extragalactic Disk Galaxies	190
6.5	Molecular Gas Observations in the Outskirts of Early-Type Galaxies	195
6.6	Molecular Gas Observations in Galaxy Groups and Clusters	197
6.7	Conclusions and Future Directions	198
	References	199

7	HI in the Outskirts of Nearby Galaxies	209
	Albert Bosma	
7.1	Introduction	209
7.2	HI in Galaxies and the Dark Matter Problem: Early Work	210
7.3	Warps	212
7.4	Further Data on HI in Galaxies and the Dark Matter Problem	214
7.5	The Disk-Halo Degeneracy in the Dark Matter Problem	215
7.6	Flaring of the Outer HI Layer: Probing the Shape of the Dark Matter Halo	219
7.6.1	Early Work on Case Studies	219
7.6.2	Recent Results for Small, Flat Galaxies	220
7.6.3	Large Galaxies with a High Star Formation Rate: Accretion	221
7.6.4	Velocity Dispersions in the Outer HI Layers of Spiral Galaxies	224
7.6.5	Star Formation in Warped HI Layers	227
7.7	The Core-Cusp Problem	227
7.8	Alternative Gravity Theories	230
7.9	Irregular Galaxies	232
7.9.1	Very Large HI Envelopes	232
7.9.2	Velocity Dispersions in Dwarf Irregular Galaxies	235
7.10	The Relation Between HI Extent and the Optical Radius	235
7.11	Concluding Remarks	238
	References	239
8	Ultra-Deep Imaging: Structure of Disks and Haloes	255
	Johan H. Knapen and Ignacio Trujillo	
8.1	Introduction	255
8.2	The Challenges of Ultra-Deep Imaging	257
8.2.1	Sky Brightness	257
8.2.2	Internal Reflections	257
8.2.3	Flat Fielding	258
8.2.4	Masking and Background Subtraction	259
8.2.5	Scattered Light	261
8.2.6	Galactic Cirrus	263
8.3	Approaches in Ultra-Deep Imaging	263
8.3.1	Survey Data	263
8.3.2	Small Telescopes	264
8.3.3	Large Telescopes	266
8.4	Disk and Stellar Halo Properties from Ultra-Deep Imaging	267
8.4.1	Thick Disks	267
8.4.2	Truncations	270
8.4.3	Tidal Streams	272
8.4.4	Stellar Haloes	273
8.4.5	Satellites	277

8.5	Conclusions and Future Developments	279
	References	281
9	Outskirts of Distant Galaxies in Absorption	291
	Hsiao-Wen Chen	
9.1	Introduction	291
9.2	Tracking the Neutral Gas Reservoir over Cosmic Time	295
9.3	Probing the Neutral Gas Phase in Galaxy Outskirts	302
9.4	The Star Formation Relation in the Early Universe	305
9.5	From Neutral ISM to the Ionized Circumgalactic Medium	309
9.6	Summary	317
	References	318
10	Future Prospects: Deep Imaging of Galaxy Outskirts Using Telescopes Large and Small	333
	Roberto Abraham, Pieter van Dokkum, Charlie Conroy, Allison Merritt, Jielai Zhang, Deborah Lokhorst, Shany Danieli, and Lamiya Mowla	
10.1	Motivation	334
10.2	Why Is Low Surface Brightness Imaging Hard?	335
10.3	Small Telescope Arrays as Better Imaging Mousetraps	337
10.4	The Dragonfly Telephoto Array	340
10.5	The Universe Below 30 mag/arcsec ²	343
	10.5.1 Galactic Outskirts	343
	10.5.2 Ultra-Diffuse Galaxies	349
	10.5.3 Imaging the Cosmic Web: The Next Frontier?	351
	References	353
	Index	359

Chapter 1

Outer Regions of the Milky Way

Francesca Figueras

Abstract With the start of the *Gaia* era, the time has come to address the major challenge of deriving the star formation history and evolution of the disk of our Milky Way. Here we review our present knowledge of the outer regions of the Milky Way disk population. Its stellar content, its structure and its dynamical and chemical evolution are summarized, focussing on our lack of understanding both from an observational and a theoretical viewpoint. We describe the unprecedented data that *Gaia* and the upcoming ground-based spectroscopic surveys will provide in the next decade. More in detail, we quantify the expected accuracy in position, velocity and astrophysical parameters of some of the key tracers of the stellar populations in the outer Galactic disk. Some insights on the future capability of these surveys to answer crucial and fundamental issues are discussed, such as the mechanisms driving the spiral arms and the warp formation. Our Galaxy, the Milky Way, is our cosmological laboratory for understanding the process of formation and evolution of disk galaxies. What we learn in the next decades will be naturally transferred to the extragalactic domain.

1.1 Introduction

The kinematical and chemical characterization of the stellar populations in the outer regions of the galactic disks is a crucial and key element to understand the process of disk formation and evolution. These outer regions are areas in the low-density regime and thus hard to observe in external galaxies. It is in this context that our Galaxy, the Milky Way, can truly be a laboratory and an exceptional environment to undertake such studies. The disk formation in our Milky Way was an extended process which started about 10 Gyr ago and continues to the present. Throughout this evolution, time-dependent dynamical agents such as radial migration or resonant scattering by transient or long-lived structures have been driving the orbital motion of the stars. Concerning star formation and evolution,

F. Figueras (✉)

Institute of Cosmos Science (IEEC-UB), University of Barcelona, Barcelona, Spain
e-mail: cesca@fqa.ub.edu

© Springer International Publishing AG 2017

J.H. Knapen et al. (eds.), *Outskirts of Galaxies*, Astrophysics and Space Science Library 434, DOI 10.1007/978-3-319-56570-5_1

key factors such as the initial mass function and the star formation history are fundamental ingredients to describe the growth of disks. Furthermore, it has to be kept in mind that the process of disk formation also involves agents which are not yet well understood, from the dynamical influence of the puzzling three-dimensional structure of the dark matter halo to the characterization of the infalling gas which gradually builds up the disk and forms stars quiescently. The scenario becomes even more complex when other decisive factors such as mergers or gravitational interactions with satellites come into play.

Two approaches have usually been considered. From the extragalactic point of view, disks at different redshifts can be studied. This approach is limited to global information integrated over the disk stellar populations but has the advantage of tracing the evolution of disk properties with time. In a second approach, normally referred to as galactic archaeology (the approach used for the Milky Way), the disk evolution is reconstructed by resolving the stellar populations into individual stars. Disk evolution is fossilized in the orbital distribution of stars, their chemical composition and their ages. A drawback of this approach is that this information may be diluted through dynamical evolution and radial mixing. Tracers can be stars, open clusters and/or gas. In this Chapter we will concentrate on the stellar component. Molecular gas is described by Watson and Koda (2017). Nonetheless, when studying the Milky Way, we should not forget the perspective that in several aspects, our Galaxy is not a typical late-type spiral galaxy. The unusually quiescent merger history of the Milky Way has been discussed in detail by van der Kruit and Freeman (2011), who remark

...the unique possibility to make very detailed chemical studies of stars in the Milky Way provides an independent opportunity to evaluate the merger history of our large disk galaxy...

In Sect. 1.2 we describe the current knowledge of the stellar content in the outer regions of the Galactic disk. Section 1.3 deals with the non-axisymmetric structures that most contribute to the dynamics of these regions. Later on, in Sect. 1.4, we introduce some of the essential pieces towards a future chemo-dynamical model of the Milky Way. Finally, in Sect. 1.5, we present a brief overview of the upcoming new astrometric and spectroscopic data, both from the *Gaia* space astrometry mission and from ongoing and future ground-based spectroscopic surveys.

1.2 The Outer Disk of the Milky Way: Stellar Content

In this Section we first describe some of the best tracers of the structure and kinematics of the outer regions of the Milky Way. Very useful tools such as the Besançon stellar population synthesis model (Robin et al. 2014; Czekaj et al. 2014) allow us to quantify the number of field stars belonging to each tracer population (thin and thick disk or halo spheroid in the case of the outer regions). The key ingredients used in the strategy to generate simulated samples are shown in Fig. 1.1




Fig. 1.1 General scheme describing the Besançon Galaxy model ingredients (Czekaj et al. 2014). Credit: M. Czekaj, et al. A&A, 564, 3, 2014, reproduced with permission © ESO

(from Czekaj et al. 2014). Although the model is continuously being updated, caution has to be taken when analysing the properties of the generated samples. The model includes several critical assumptions such as the radial scale length or the disk cut-off, which can induce significant discrepancies between model and observations. A separate paragraph is devoted to Cepheid variables, whose high intrinsic brightness makes them excellent tracers of the radial metallicity gradient (Lemasle et al. 2013). Properties of other very good tracers such as open clusters are not treated here. The reader can find an updated characterization of this population in recent papers such as the one by Netopil et al. (2016). A summarizing discussion on the present knowledge of the outer reaches is presented in Sect. 1.2.2.

1.2.1 Resolved Stellar Populations

Stellar Tracers As an example, in Fig. 1.2 we show the radial galactocentric distribution of three different stellar populations: Red Clump K-giant stars and main-sequence A- and OB-type stars. The estimated number of stars that *Gaia* will observe in the outer Milky Way region, i.e. at galactocentric radii between 9 and 16 kpc, is shown there. These estimates come from the work of Abedi et al. (2014) where test particles were generated fitting the stellar density at the position of the Sun, as provided by the Besançon galaxy model (Czekaj et al. 2014). In this case, the three-dimensional Galactic extinction model by Drimmel et al. (2003) has been used. From this model we estimate that the visual extinction, A_V , in the mean does not exceed ~ 2 magnitudes when looking towards the Galactic anticentre




Fig. 1.2 Histograms of the numbers of stars in galactocentric radius bins of 1 kpc to be observed by the *Gaia* satellite (see Abedi et al. 2014 for details). The samples with *Gaia* magnitudes $G \leq 20$, $G_{RVS} \leq 17$ and those with expected parallax accuracy less than 20% (end of mission) are shown, respectively, in red, purple and green. The histograms are plotted for Red Clump stars (left panel), A stars (middle) and OB stars (right)

at low galactic latitudes. These low values for the interstellar extinction and the large number of stars to be fully characterized by *Gaia* and the future spectroscopic surveys (see Sect. 1.5) encourage the studies of resolved stellar populations towards the Galactic anticentre and the outer regions of the Milky Way.

Classical and Type II Cepheids These pulsating intrinsically bright variable stars are excellent tracers of the extent of the thin/thick disk surface densities and also of the abundance distribution of numerous chemical elements. Whereas classical Cepheids are young stars (< 200 Myr; Bono et al. 2005) associated with young stellar clusters and OB associations, Type II Cepheids, with some characteristics similar to the classical ones (e.g. period and light curve), are fainter and much older, although their evolutionary status is still not firmly established. Both are interesting targets for the outer disk. Classical and Type II Cepheids can probably be associated with the thin and thick disk populations, respectively. Cepheids have been observed at radial galactocentric distances up to $R = 18$ kpc. Furthermore, as reported by Feast et al. (2014), a few classical Cepheid stars have been found at approximately 1–2 kpc above the plane in the direction of the Galactic bulge, at distances 13–22 kpc from the Galactic centre. Their presence, far from the plane, suggests that they are in the flattened outer disk and thus are excellent tracers of this at present very unknown structure.

1.2.2 The Outer Reaches

The Disk Cut-Off An apparent and sudden drop was reported by van der Kruit and Searle (1981) in the surface brightness of several edge-on galaxies at a radius of about four disk scale lengths. This has been a long-standing question both in external

galaxies and in our Milky Way. Nowadays, when looking at external galaxies, one plausible explanation is that the reported edges are, in fact, inflections in the stellar density, i.e. breaks in the exponential density profiles (Bland-Hawthorn and Gerhard 2016). Does our Milky Way have such a truncation? A first analysis using deep optical star counts at a low-extinction window in the Galactic anticentre direction showed a clear signature of a sharp cut-off in the star density at about 5.5–6 kpc from the Sun (Robin et al. 1992). Later on, Momany et al. (2006), using 2MASS data, inferred robust evidence that there is no radial disk truncation at $R = 14$ kpc. More recently Minniti et al. (2011), using the UKIDSS-GPS and VVV surveys, pointed out that there is an edge of the stellar disk at about $R_{GC} = 13.9 \pm 0.5$ kpc along various lines of sight across the galaxy. When analysing these data, it has to be kept in mind that changes in the star counts induced by the warp and flare may not be negligible. New data seem to disagree with a sharply truncated nature, and proof of this is the presence of stars and even star formation regions beyond the break radius. López-Corredoira and Molgó (2014), using also star count techniques and Sloan Digitized Sky Survey (SDSS) data, quantified the change of the vertical scale height with galactocentric radius of the Galactic thin and thick disks. The presence of the Galactic flare, quite prominent at large R , can explain the apparent depletion of in-plane stars that is often confused with a cut-off at $R \sim 14$ – 15 kpc. Furthermore, in a recent paper, Carraro et al. (2016) studied the spatial distribution of early-type field stars and open clusters in the third Galactic quadrant of the Milky Way. Their Fig. 12 summarizes the spatial distribution of the young population. According to these authors, the field star sample extends up to 20 kpc from the Galactic centre, with no indication of the disk cut-off truncation at 14 kpc from the Galactic centre previously postulated by Robin et al. (1992).

The Monoceros Ring and Beyond The Monoceros Ring, a coherent ring-like structure at low Galactic latitude spanning about 100° and discovered by Newberg et al. (2002), thanks to the SDSS, deserves special attention. This structure was first identified as an overdensity of stars at ~ 10 kpc, with a metallicity in the range $-1 < [\text{Fe}/\text{H}] < 0$ (with a substantial scatter). It is a poorly understood phenomenon with no clear association to other structures such as the Canis Major or the Triangulum-Andromeda overdensity. Recent observations from PAN-STARSS1 (Slater et al. 2014) indicate a larger extent of the stellar overdensity, up to $|b| \sim 25$ – 35° and about 130° in Galactic longitude. Its origin and gravitational interaction with the Milky Way are unclear (see Sect. 1.3.3). More recently, Xu et al. (2015), using SDSS data, have reported the existence of an oscillating asymmetry in the main-sequence star counts on either side of the Galactic plane in the anticentre region. These stellar overdensities, identified in a large Galactic longitude range, $[110^\circ, 229^\circ]$, are oscillating above and below the plane and have been observed up to distances of about 12–16 kpc from the Sun. As shown in Fig. 1.3, the three more distant asymmetries seem to be roughly concentric rings, open in the direction of the Milky Way spiral arms. The Monoceros Ring is identified as the northernmost of these structures, and the others are the so-called Triangulum-Andromeda overdensities (first detected by Rocha-Pinto et al. 2003), which could extend up to at least 25 kpc from the Galactic centre.




Fig. 1.3 Stellar overdensities above (*left*) and below (*right*) the plane reported by Xu et al. (2015), using SDSS data. The (X, Y) coordinates are centred at the position of the Sun

1.3 The Milky Way Outer Disk: Structure and Dynamics

Here we will focus on those non-asymmetric structures that most contribute to the dynamics of the outer regions of the Milky Way. For an exhaustive and updated description of other components such as the Galactic bar or inner spirals, the reader is referred to the recent review by Bland-Hawthorn and Gerhard (2016).

1.3.1 *Spiral Arm Impact on Disk Dynamics and Structure*

As is well known, spiral arms have a great impact on the evolution of galactic disks, from driving the formation of massive clouds to the perturbation of the stellar orbits of the old populations. Phenomena such as resonant trapping at corotation and Lindblad resonances, streaming motion and radial migration (Sellwood and Binney 2002) can be understood in this context. Furthermore, the impact of the evolution of these non-axisymmetric structures on the current radial chemical gradient of the Galactic disk or on the age-metallicity relation cannot be understood without considering spiral arms (see Sect. 1.4).

Accurate kinematic data are, without doubt, a first requirement to distinguish between spiral structure theories and thus the mechanisms of formation and evolution of spiral arms. Some of the current theories under investigation are the density wave theory (Lin and Shu 1964), with a rigid spiral density wave travelling through the disk; swing amplification (Toomre 1981; Masset and Tagger 1997), with perturbations on the disk that can be swing amplified; invariant manifolds (Romero-Gómez et al. 2006; Athanassoula 2012), with manifolds originated in the periodic

orbits around the equilibrium points; external interactions (Sellwood and Carlberg 1984), for a certain fraction of mass accretion; or chaotic orbits (Voglis et al. 2006; Patsis 2006), important for a large perturbation, especially near corotation. It is difficult to test these theories with current available data, and complicating the situation even more, several of them may coexist. As an example, the invariant manifold mechanism has been tested using N -body simulations of barred galaxies (Athanasoula 2012; Roca-Fàbrega et al. 2013). A detailed characterization of the motion of the particles through the arms or crossing them is required (Antoja et al. 2016). It is clear that an accurate knowledge of the kinematics is mandatory to distinguish between theories (see Sect. 1.5).

Recently, Monguió et al. (2015) published the first detection of the field star overdensity in the Perseus arm towards the Galactic anticentre. Using a young population of B- and A-type stars, the authors placed the arm at 1.6 ± 0.2 kpc from the Sun, estimating its stellar density amplitude to about 10%. Moreover, these authors show how its location matches a variation in the dust distribution congruent with a dust layer in front of the arm. This favours the assumption that the Perseus arm is placed inside the corotation radius of the Milky Way spiral pattern. The obtained heliocentric distance of the Perseus arm is slightly smaller than the 2.0 kpc recently proposed by Reid et al. (2014). These authors used VLBI trigonometric parallaxes and proper motions of masers, thus star-forming regions, to accurately locate many arm segments in the Galactic disk. The outer spiral arm seems to be located at 13.0 ± 0.3 kpc from the Galactic centre, with an amplitude 0.62 ± 0.18 kpc and a pitch angle of $13.8 \pm 3.3^\circ$. Another critical issue is the age dependence of the radial scale length. Several determinations can be found in the literature. Recently, Monguió et al. (2015) derived it using the Galactic disk young population. They obtained values of 2.9 ± 0.1 kpc for B4-A1 type stars and 3.5 ± 0.5 kpc for B4-A0 stars. Unfortunately, the uncertainty associated with these data still prevents a direct application of these results in discriminating between inside-out or out-inside formation scenarios.

1.3.2 *The Galactic Warp and Flare*

It is widely accepted that warps of disk galaxies are a common phenomenon (as common as spiral structure), yet warps are still not fully understood (see García-Ruiz et al. (2002) for a historical review). From the time when the first 21 cm observations of our Galaxy became available, the large-scale warp in the HI gas disk has been apparent (Burke 1957; Westerhout 1957). More than 50 years later, Levine et al. (2006) re-examined the outer HI distribution, proposing a more complex structure with the gaseous warp well described by two Fourier modes. The warp seems to start already within the Solar circle. Reylié et al. (2009), using 2MASS infrared data, found the stellar component to be well modelled by an S-shaped warp with a significantly smaller slope than the one seen in the HI warp. Since then, several authors have tried to set up the morphology of the Galactic warp, reaching




Fig. 1.4 The effect of the Galactic warp on the distribution of the mean heliocentric vertical velocity component (W) as a function of galactocentric azimuth for Red Clump stars in the galactocentric ring $13 < R < 14$ kpc. Two warp models (A and B) described in López-Corredoira et al. (2014) are plotted here (dashed purple and solid blue lines) and compared to the best fit to PPMXL proper motion data (in black). See López-Corredoira et al. (2014) for details

no clear conclusion. More importantly, the current uncertainties do not allow us to disentangle which mechanisms can explain it. Critical issues are the kinematics of the warped population and basic properties such as its stellar age dependence. Data available currently (PPMXL proper motions) allowed López-Corredoira et al. (2014) (see Fig. 1.4) to perform a vertical motion analysis of the warp towards the Galactic anticentre. They point out that whereas the main S-shaped structure of the warp is a long-lived feature, the perturbation that produces an irregularity in the southern part is most likely a transient phenomenon. Again, this is a complex kinematic feature from which a definitive scenario is difficult to constrain. We refer to the recent work of Abedi et al. (2014) to know the capabilities of *Gaia* in detecting and characterizing the kinematic properties of the warp. Accurate proper motions and radial velocities (see Sect. 1.5) will certainly add a new dimension to this study.

Lozinskaya and Kardashev (1963) were the first to describe the observational evidence of a flare in the HI gas in the Milky Way. Since then, the question of whether the stellar component takes part in this flaring has been a matter of debate. Momany et al. (2006) presented a first comparison of the thickness of the stellar




Fig. 1.5 Comparison by Kalberla et al. (2014) of the current observational data (*points*) and exponential scale height fits (*continuous lines*) to quantify the flare of the Galactic thin (*left*) and thick (*right*) components. See details in their paper

disk, neutral hydrogen gas layer and molecular clouds. The stellar flare was traced using Red Clump and red giant stars. These authors found that the variation of the disk thickness (flaring)—treating a mixture of thin and thick stellar populations—starts at $R = 15$ kpc and increases gradually until reaching a mean scale height of ~ 1.5 kpc at $R = 23$ kpc. More recently, Kalberla et al. (2014) confirmed these results, showing strong evidence for a common flaring of gas and stars in the Milky Way. Several sources such as HI gas, Cepheids, 2MASS, SDSS and pulsar data show an increase of the scale height, growing with galactocentric radius (see Fig. 1.5). Even more, it has been proposed that flaring at large galactocentric distances could be stronger for the thin than for the thick disk. Although this is still a matter of debate, with *Gaia* we will have the opportunity not only to analyse the structure but also the kinematic properties of these outer populations.

Several models have been proposed to account for this flare. Whereas Kalberla et al. (2007) explored several mass models reflecting different dark matter distributions, others, such as Minchev et al. (2012), by preassembling N -body disks, showed that purely secular evolution could lead to flared disks. Others, like Roškar et al. (2010), proposed misaligned gas infall to provoke the flaring of the Galactic disk. Recent observational analysis of SDSS data points towards a smooth stellar distribution (López-Corredoira and Molgó 2014), thus supporting a continuous structure for the flare and not a combination of a Galactic disk plus some component of extragalactic origin.

1.3.3 Gravitational Interaction with Satellites

Several approaches have been followed in the last years to characterize the effects on the Galactic disks induced by dynamical perturbations by satellites (e.g. Purcell et al. 2011; Gómez et al. 2013, among others). All of them reinforce the picture that the Galactic disk can exhibit complex structure in response to close satellite

passages, from tidal debris in a disrupted dwarf galaxy to a strong gravitational perturbation by the accretion of a satellite. Work is in progress from both the observational and the theoretical side. As an example, Peñarrubia et al. (2005) proposed a model for the formation of the Monoceros Ring by accreted satellite material. Their model has recently been compared to PAN-STARSS1 data (Slater et al. 2014) at different distances, showing broad agreement with the observed structure at mid-distances but significant differences in the far regions. These and newer simulations have since been compared to the data. Recently, Gómez et al. (2016) studied the vertical structure of a stellar disk obtained from a fully cosmological high-resolution hydrodynamical simulation of the formation of a Milky Way-like galaxy. The disk's mean vertical height can have amplitudes as large as 3 kpc in its outer regions as a result of a satellite-host halo-disk interaction. The simulations reproduce, qualitatively, many of the observable properties of the Monoceros Ring. Nonetheless, as pointed out by Slater et al. (2014), the crucial question for future simulations is whether such Monoceros Ring-like features can be created without causing such an unrealistically large distortion of the disk; maybe less massive satellites or particular infall trajectories could be more favourable.

1.3.4 Dynamics of the Vertical Blending and Breathing Modes

We currently have observational evidence of oscillations of the Milky Way's stellar disk in the direction perpendicular to the Galactic midplane. Some of these are possibly related to the Monoceros and Triangulum-Andromeda overdensities in the outer disk (Sect. 1.2.2), while others are more local, within ~ 2 kpc of the Sun. Widrow et al. (2012), using SDSS/SEGUE, RAVE and LAMOST data, found a Galactic north-south asymmetry in the number density and bulk velocity of Solar neighbourhood stars which showed a gradual trend across the Galactic midplane and thus the appearance of a wavelike perturbation. This perturbation has the characteristics of a breathing mode, with compression and rarefaction motions and with the displacements and peculiar velocities having opposite signs above and below the plane. Widrow et al. (2014) demonstrate that both breathing and bending modes can be generated by a passing satellite or dark matter subhalo, with the nature of the perturbation being controlled by the satellite's vertical velocity relative to the disk (a slow-moving satellite would induce a bending mode, whereas higher vertical velocities would induce breathing modes). More recently, Monari et al. (2016) discussed that breathing modes can be induced by several effects such as bar or spiral perturbation, spiral instabilities or a possible bombarding by satellites. Even the existence of a dark matter substructure could play a role. These authors have derived explicit expressions for the full perturbed density function of a thin disk stellar population in the presence of non-axisymmetric structures such as spiral arms and bar. Undoubtedly, upcoming data will yield a more accurate and complete map of bulk motions in the stellar disk, up to about 3–4 kpc from the Sun (Sect. 1.5).

1.4 Towards a Chemodynamical Model of the Galactic Disk

Prantzos (2008, 2011) are two excellent and pedagogical papers to review the basic principles and hot topics for the development of galactic chemical evolutionary models. The state of the art is well summarized there. These models shall assume, among other factors, the evolution of several chemical elements up to the iron peak and the different compositions for the infalling material. Accurate observations are required to test these models, the products of which are usually expressed in terms of radial abundance gradients of several elements (C, N, O, Ne, Mg, Al, Si, S, Ar, Fe) and their time evolution. Thanks to the extent of present and future large surveys (see Sect. 1.5), any formation model must be able to account not only for local but also for radial and vertical large-scale chemical element distributions.

Radial migration has been firmly accepted to be an inseparable part of disk evolution in numerical simulations (Sellwood and Binney 2002). In radial migration models (e.g. Schönrich and Binney 2009), metal-poor stars born in the outer disk move inwards to the Solar neighbourhood, while metal-rich stars born in the inner disk migrate outwards. These authors suggest two mechanisms for this motion¹: *blurring*, due to the scattering and subsequent increase of eccentricities over time, and *churning*, mostly triggered by resonant scattering at corotation. These effects would produce a large heterogeneity in the chemical abundance in the Solar neighbourhood and its environment.

1.4.1 Age-Metallicity-Kinematics Relations

It is well established that the velocity dispersion of the thin disk population increases with age, a well-known phenomenon often referred to as disk heating (Binney and Tremaine 1987). Aumer and Binney (2009) used a power law to fit the thin disk age-velocity relation (AVR) in the Solar neighbourhood, using the most accurate data at that time: the Hipparcos astrometry and photometric and spectroscopic data from the Geneva-Copenhagen survey (Holmberg et al. 2007). They favour continuous heating, but as proposed also by Seabroke and Gilmore (2007), a saturation at ages ≥ 4.5 Gyr could not be excluded. Concerning the age-metallicity distribution (AMD), it is important to mention the work done by Haywood (2006). Biases were evaluated in detail by this author, and a new AMD with a mean increase limited to about a factor of two in Z over the disk age was proposed. Again, it was emphasized that dynamical effects and complexity in the AMD clearly dominate. All the above relations have been established using data in the Solar neighbourhood. Nonetheless, and as pointed out by Casagrande et al. (2011), the Solar neighbourhood is not only

¹Following the Schönrich and Binney (2009) terminology, churning implies a change of guiding-centre radii, while blurring means a steady increase of the oscillation amplitude around the guiding centre.

assembled from local stars but also from stars born in the inner and outer Galactic disk that migrated to their present positions (Schönrich and Binney 2009). New surveys (see Sect. 1.5) will open a new window and different approaches to these two key and fundamental relations.

In this direction, a new stellar chemo-kinematic relation has recently been derived by Minchev et al. (2014) using RAVE and deeper surveys such as SEGUE. Stars with $[\text{Mg}/\text{Fe}] \geq 0.4$ dex show a peculiar kinematic behaviour. These authors used this index as a proxy of the stellar age to identify the oldest stars in the sample. These stars, born during the first years of the Galaxy's life, have velocity dispersions too large to be accounted for by internal disk heating. Minchev et al. (2014) showed that a chemo-dynamical model incorporating massive mergers in the early Universe and a subsequent radial migration of cool stars could explain the observed trends. More imprints such as the ones reported there should be expected in the outer regions of the Galactic disk, so this work is a good example of the chemo-kinematic relations that the combination of future *Gaia* spectroscopic surveys will provide. They will surely bring new constraints to the formation scenarios of galactic disks.

1.4.2 *The Galactic Thick Disk*

The characterization of the thick disk is an important milestone when trying to understand the assembly of disk galaxies. Despite the increasing amounts of observational data in our local environment, to date we have lacked the means to discriminate among different thick disk formation scenarios for the Milky Way. A first key and basic question which arises is: Do the thin and thick disks have a different origin? Two approaches are being used to answer this question, observational evidence and galaxy modelling. Observational evidence at present is, in some sense, highly inconsistent. Whereas Bovy et al. (2012), using SDSS data, clearly favoured a vertical structure composed of a smooth continuum of disk thicknesses with no discontinuity between the thin and thick disk, a bimodal distribution in the $([\text{Fe}/\text{H}], [\alpha/\text{Fe}])$ relation with two sequences of high and low $[\alpha/\text{Fe}]$ seems well established (Adibekyan et al. 2012; Nidever et al. 2014) with high $[\alpha/\text{Fe}]$ values more prominent in the inner disk and lower values dominating the outer parts. In this line, Recio-Blanco et al. (2014) point towards a clear kinematic distinction between thin and thick disk (two distinct populations). Concerning the structure, the thick disk occupies (in agreement with its hotter kinematics) a larger vertical volume around the midplane (e.g. Jurić et al. 2008), perhaps at a shorter scale length than the thin disk (Bensby et al. 2011; Robin et al. 2014) and a clear uncertainty in the radial metallicity gradient, as will become evident in Sect. 1.4.3.2. Nonetheless, as discussed by Bland-Hawthorn and Gerhard (2016), one of the strongest pieces of evidence of the existence of the thick disk will be a robust statistical confirmation of its unique chemistry (see the important work by Bensby et al. 2014 in this line).

Several models have been proposed to explain the existence of the thick disk: (1) a heating process of the Galactic disk due to satellite mergers (e.g. Abadi et al. 2003); (2) the formation of a puffed-up structure by mere radial migration (Sellwood and Binney 2002); (3) a so-called “upside-down” disk formation, where old stars were formed in a relatively thick component, while younger populations form in successively thinner disks (e.g. Bird et al. 2013); or (4) in situ formation by early accretion of gas. The scenario for the Milky Way thick disk is presently unclear. Several observational constraints have to be fixed. As an example, some of these models predict a vertical gradient in $[\alpha/\text{Fe}]$; others do not. The in situ formation or the direct accretion of small satellites would be ruled out if α -gradients are observed (Recio-Blanco et al. 2014). Progress in this field is further discussed in Sect. 1.4.3.2. Other future constraints could be, as pointed out by B. Lemasle (private communication), the ratio of classical versus Type II Cepheids. The galactocentric radial change of these ratios, available from *Gaia*, would help us to analyse the extent of the thin/thick disk structures.

1.4.3 The Radial Abundance Gradients

Examples of two reference papers on chemical evolutionary models that show different but complementary approaches are those of Chiappini et al. (1997) and Pilkington et al. (2012a). Chiappini et al. (1997) developed a new model for the Galaxy assuming two main infall episodes for the formation of the halo-thick disk and thin disk. The model also predicts the evolution of the gas mass, the star formation rate, the supernova rates and the abundances of 16 chemical elements as functions of time and galactocentric distance. In summary, a long list of detailed model results can be constrained when compared with observational data. The approach of Pilkington et al. (2012a) is different and complementary. These authors used cosmological hydrodynamical simulations of dwarf disk galaxies to analyse the distribution of metals. Both approaches require the comparison of models and data and hence the selection of good stellar tracers, as discussed below.

The differences between the models concern effects such as the efficiency of the enrichment processes in the inner and outer regions and the nature of the material (primordial or pre-enriched falling from the halo onto the disk). These questions can help answer fundamental questions such as whether our Galactic disk has a flattening or a steepening radial metallicity gradient with time. Open clusters and Cepheids are proposed as good tools to derive the time evolution of the metallicity gradients as the need to answer this question is pressing. As an example, in the future WEAVE spectroscopic survey (Sect. 1.5), the requirement on the accuracy of the open cluster metallicity is 0.1 dex over the full age and metallicity range.

Tracers of Chemical Abundance Gradients As mentioned by Boissier and Prantzos (1999), all the existing observational data only inform us on the present properties of the Galactic disk, but not on its past history, except for the tentative

Table 1.1 Some of the ongoing and future ground-based surveys for chemodynamical studies of field stars in the outer regions of the Milky Way

Survey	Dates	Magnitude	Resolution	Trace populations
<i>Gaia</i> -ESO	2011–2016	$V \sim 16\text{--}20$	$\sim 20,000$	Red Clump giants
<i>Gaia</i> -ESO	2011–2016	$V < 16$	$\sim 45,000$	Thick disk population
LAMOST	2012–	$r \sim 19\text{--}20$	1800	North hemisphere
APOGEE	2011–	$H < 12.2$	22,500	Red giants and subgiants
EMIR	2017–	$J, K \sim 18$	4000	Red giants
WEAVE	2018–2022	$V = 16\text{--}20$	4000	Disk OBA and Red Clump giants
WEAVE	2018–2022	$V < 16\text{--}17$	20,000	Anti-centre OBA and RC disk
4MOST	2022–	$V \leq 20$	> 2000	From Sun-like to RGBs

indications obtained from the study of abundances in planetary nebulae or in open clusters. Alibes et al. (2001) reviewed the main observables for the disk of the Milky Way, which can be compared with the products of a chemical evolutionary model. As emphasized by these authors, homogeneous samples are required to derive radial gradients. The young population (age < 1 Gyr) provides information on the present-day abundances in the Galactic disk, the most important tracers being the HII regions and the B-type stars. HII regions have the drawback of large uncertainties due to the derivation of electron temperature. The B-type stars are bright, thus observed up to large distances, with a photospheric composition which supposedly reflects the composition of the material from which they formed, except for high rotators where there could have been some mixing of atmospheric and core material. Tracers older than 1 Gyr provide insight into the past evolution of the abundance gradients and thus into the evolution in abundance profiles. Among these are the planetary nebulae, with large uncertainties in distances, ages and masses; open clusters, which can be as old as 8 Gyr; and the FGK stars, dwarfs and giants, a population selected as tracer in most of the ongoing large spectroscopic surveys (see Table 1.1).

1.4.3.1 Cepheids as Tracers of Galactic Abundance Gradients

Since the early work by Harris (1981), the determinations of the Galactic $[\text{Fe}/\text{H}]$ gradient from Galactic Cepheids have been remarkably consistent, all of them reporting values of ~ -0.06 dex/kpc (Lemasle et al. 2013). From open clusters, a linear gradient of approximately -0.06 dex/kpc has been proposed (e.g. Bragaglia et al. 2008) but with a flattening between 10 and 14 kpc, that is, in the outer regions of the Galactic disk. This possible flattening of the abundance gradients is a long-standing question. Whereas it is not observed from Galactic Cepheids (Lemasle et al. 2013), a clear flat (bimodal) gradient in the outer disk is derived from open clusters. Lemasle et al. (2013) measured the Galactic abundance gradient of 11 chemical elements using high-resolution spectra of more than 60 Galactic Cepheids and focussed their analysis on the outer Galactic disk. They confirm the existence of

a gradient for all the heavy elements and noted that current data are not supporting a flattening of the gradient in the outer disk. Later on, the same team (Genovali et al. 2014) compiled homogeneous data for a sample of 450 Cepheids deriving a linear metallicity gradient over a broad range of galactocentric distances ($R \sim 5\text{--}19$ kpc). These authors pointed out clear evidence for the increase in the intrinsic scatter when moving towards the outer disk. It cannot be ruled out that this scatter is due to contamination by Type II Cepheids misclassified as classical Cepheids (Sect. 1.2.2).

1.4.3.2 Recent Outcomes of the *Gaia*-ESO Survey

From current surveys (SEGUE, RAVE, APOGEE, GALAH, *Gaia*-ESO Survey, among others), the large-scale characteristics of the thin and thick disks have started to emerge. It is difficult to summarize here the huge amount of new results published in the last 5–10 years. In this section, as it is impossible to make an exhaustive review, we will take one of them as an example, in this case the effort being done by the European *Gaia* research community (more than 300 scientists) to build the *Gaia*-ESO Survey (hereafter GES). The survey, planned for 5 years (2011–2016) and still ongoing, uses the FLAMES/GIRAFFE and UVES spectrographs on the Very Large Telescope (ESO), and important preliminary results have already been published. Two approaches have been followed to derive the vertical and radial trends of the chemical gradient, from open clusters and from FGK main-sequence field stars.

Open clusters are, apart from Cepheids (Sect. 1.4.3.1), the best tracers to derive metallicity over large ranges in galactocentric distance. However, accurate and homogeneous observations of open clusters are required to properly trace the chemical abundance distribution in the Galactic disk. Recently, Jacobson et al. (2016) showed that the GES open clusters at galactocentric distances [5.5, 8] kpc exhibit an [Fe/H] radial metallicity gradient of -0.10 ± 0.02 dex/kpc, consistent with values obtained by Netopil et al. (2016) with a larger open cluster sample (172 open clusters) at distances $R = [6, 14]$ kpc compiled using different criteria. Both datasets are showing us no evidence for a steepening of the inner disk metallicity gradient inside the Solar circle. Furthermore, Netopil et al. (2016) compared open clusters and Cepheids with the Galactic chemical model by Minchev et al. (2013) (see Fig. 1.6). Open clusters show a much flatter gradient than Cepheids or these model predictions. Possible explanations proposed for this flattening are an underestimation of radial migration in the models or other not yet included mechanisms acting at large galactocentric radius. It is evident that more open cluster data are needed in the outer Galactic disk. Open clusters are introducing an important constraint on the chemical evolutionary models and their time evolution scenarios. Note that other independent data, in this case from APOGEE DR10 and with high-resolution metallicity measurements, reproduce interesting features such as the lack of gradient in the logarithmic α -elements-to-metal ratio, $[\alpha/M]$, across the range $7.9 < R < 14.5$ kpc (Frinchaboy et al. 2013).




Fig. 1.6 Radial metallicity gradient derived from Cepheids and open cluster data compiled, respectively, by Genovali et al. (2014) and Netopil et al. (2016). The *left panel* compares the youngest populations of these tracers with models. The *grey line* and the *light grey area* show the running average and the spread in iron of the Cepheid data. The *black dashed line* shows the gradient derived from a subset of young open clusters with ages less than 0.5 Gyr. For comparison, the *red line* indicates the gradient for the gas from the Galactic chemical model by Minchev et al. (2014). The *right panel* presents the metallicity gradient derived from an intermediate population. The *grey area* shows the running average obtained from open clusters with ages between 0.5 and 3.0 Gyr. The *horizontal black solid line* is the mean value of those clusters located in the outer disk, with galactocentric distances larger than 12 kpc. The *black dashed line* is the gradient derived for all clusters in the outer disk. For comparison, the *red dashed line* shows the Galactic chemical model by Minchev et al. (2013) that includes the dynamical effects of the Galactic disk after 2 Gyr. This Figure is taken from Netopil et al. (2016); see this paper for details

Concerning field stars, the GES FGK dwarf and giant survey allows, for the first time, to extend high-resolution spectroscopic surveys for field stars further away from the Solar vicinity. Mikolaitis et al. (2014) published a first radial metallicity abundance gradient of eight elements (Mg, Al, Si, Ca, Ti, Fe, Cr, Ni and Y) from this survey. The bimodality observed in the $[Mg/M]$ distribution was attributed to thick and thin disk populations. Both radial (4–12 kpc) and vertical (0–3.5 kpc) gradients in metallicity were derived for the thin and the thick disk separately. The interpretation of this huge amount of data is complex, with the thin disk showing a positive radial $[\alpha/M]$ gradient in contrast to the flat gradient derived for the thick disk population. Concerning vertical gradients, the thick disk hosts a shallower negative vertical metallicity gradient than the thin disk in the Solar cylinder. These results and upcoming new analyses of GES, APOGEE and SEGUE data, among others, are being examined in the context of new and complex models: from detailed analytic local disk models (e.g. Just and Rybizki 2016) to N -body simulations of Milky Way-like galaxies (e.g. Kawata et al. 2017; Roca-Fàbrega et al. 2016). In all cases, kinematic data will play a fundamental role here (see, e.g. Kordopatis et al. 2016).

1.4.4 The “*Outside-In*” Versus “*Inside-Out*” Disk Formation Scenarios

The comparison of observations and models involves global properties of the Milky Way such as rotation curves, scale lengths, masses (gas, stars and dark matter), gas flows and stellar abundances. Moreover, dynamical aspects also play a critical role. They can be analysed and taken into account through analytical or semi-analytical models or from cosmological N -body plus hydrodynamical models, the latter ones needed to assess the effect of the environment and satellite accretion on shaping the outer disks. Several efforts have been conducted up to now in this direction. As an example of the heterogeneity in the obtained results, we can mention the works of Cescutti et al. (2007), Pilkington et al. (2012b), Haywood et al. (2013) and Minchev et al. (2014). Whereas the first three proposed an inside-out disk formation scenario, Haywood et al. (2013) put forward an outside-in scenario to explain Solar neighbourhood data. These models are all only partially compatible with GES data (Sect. 1.4.3.2), and others have only been tested against external galaxy data (e.g. Pilkington et al. 2012a). The addition of upcoming new observational constraints to these models is complex. Although a general prediction of inside-out formation would be a fast decrease in $[\alpha/\text{Fe}]$ with increasing radius for stars in a narrow metallicity range, different slopes for the thin and thick disk population (GES results; see Mikolaitis et al. 2014) increase the complexity. This research field will be a key priority for the next decade. It will be possible to combine *Gaia* and spectroscopic data from the Milky Way with extragalactic work such as that done in the context of the CALIFA or MaNGA surveys.

1.5 Large Surveys in the Next Decade

The beginning of the twenty-first century is, without doubt, a new golden age for Galactic astronomy. Data coming from both space missions and ground-based surveys would be a dream in 1962, when Eggen, Lynden-Bell and Sandage derived the first model of galaxy formation and evolution (Eggen et al. 1962). Astrometric (*Gaia*, the Large Synoptic Survey Telescope, LSST) and spectroscopic ground-based surveys are fully complementary, providing both local and global measures of the six-dimensional phase-space distribution function, needed to encode valuable dynamical information. The status and products of the *Gaia* mission (2014–2022) are detailed in Sect. 1.5.1. In the coming years, more than 10^9 stars up to *Gaia* apparent magnitude $G \sim 20$ will have an astrometric accuracy never envisaged up to now. These data will be complemented with the data collected by the LSST. This telescope, expected to be fully operational in 2022, will measure proper motions with an accuracy of about 1 mas/year to a limit 4 mag fainter than the *Gaia* survey. Geometric parallaxes will also be provided by LSST, with an accuracy similar to that of the faintest stars in *Gaia* (~ 0.3 mas) at $V = 20$ and up to ~ 3 mas at $V = 24$.

The acquisition of good signal-to-noise spectra at intermediate and high resolution is mandatory to complement these accurate astrometric data with good radial velocities and chemical abundances for a significant number of stars. In Table 1.1 we present a non-exhaustive list of the most promising spectroscopic surveys ongoing or planned for the near future. A detailed description of the capabilities of each of these surveys can be found in their respective web pages. Here, we will focus on one of them, the WEAVE² spectroscopic survey. This survey, to be undertaken in the Northern hemisphere, will open a new window to the Galactic anticentre and thus to the outskirts of the Milky Way.

WEAVE is a multiobject spectroscopic instrument using about 1000 fibres that will be operated at the William Herschel Telescope (WHT, Canary Islands). Its science case and survey plan are almost fixed. During the first 5 years of operation (2018–2022), several Galactic plane radial velocity surveys will be executed using as kinematic tracer disk Red Clump and young OBA-type stars. WEAVE’s low-resolution mode ($R = 5000$) will provide radial velocities for millions of stars up to the *Gaia* limiting magnitude. We estimate to obtain radial velocities for about one million Red Clump stars with an accuracy better than ~ 5 km/s. Expected accuracies for stars brighter than $V = 18$ are even better, with errors as small as 1–3 km/s. Complementarily, several chemical tagging programmes for Galactic archaeology will be undertaken using the high-resolution mode of WEAVE ($R = 20,000$). Both kinds of data are critical ingredients to establish the chemical gradients of the Galactic thin and thick disk (Sect. 1.4.3) and the dynamics of the non-axisymmetric components of the outer regions of the MW (Sect. 1.3). A promising survey of well-selected open clusters (ages > 100 Myr) will significantly contribute to the derivation of the radial metallicity gradient from these excellent tracers in the outer regions of the Milky Way.

1.5.1 The *Gaia* Mission

The high-quality astrometric and photometric data already collected by *Gaia* during its first almost 2 years of successful scientific operation allow us to anticipate that, in a few years, *Gaia* will revolutionize our understanding of our own Galaxy, the Milky Way, and its surroundings. In this section we will briefly describe the *Gaia* products to be published in the first and end-of-mission data releases (end of 2022). More in detail, we will quantify the expected accuracy in position, transversal and radial velocity and stellar astrophysical parameters of some of the key stellar tracers of the outer regions of the Milky Way.

²See <http://www.ing.iac.es/confluence/display/WEAV/The+WEAVE+Project>.