

Giuseppe Milano
Andrea Grasso
Angel Calvo
Roman Brzóska
Editors



Management of Failed Shoulder Surgery



Management of Failed Shoulder Surgery

Giuseppe Milano • Andrea Grasso
Angel Calvo • Roman Brzóska
Editors

Management of Failed Shoulder Surgery

 Springer



Editors

Giuseppe Milano
Department of Orthopaedics
Catholic University
Rome
Italy

Andrea Grasso
Department of Orthopaedics
Casa di Cura Villa Valeria
Rome
Italy

Angel Calvo
Clínica del Pilar
Zaragoza
Spain

Roman Brzóska
Orthopaedic/Upper Limb
St. Luke's Hospital
Bielsko-Biała
Poland

ISBN 978-3-662-56503-2 ISBN 978-3-662-56504-9 (eBook)
<https://doi.org/10.1007/978-3-662-56504-9>

Library of Congress Control Number: 2018940098

© ESSKA 2018

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Printed on acid-free paper

This Springer imprint is published by the registered company Springer-Verlag GmbH, DE part of Springer Nature
The registered company address is: Heidelberger Platz 3, 14197 Berlin, Germany

*To my father and my mother, fragile and brave persons,
for their example of living (GM)*

*To my family, my Dad, my Mum and my wife. Thanks for
always helping and supporting me (AG)*

Dedicated to my four sons: Mar, Rocío, Angel and Alex (AC)

To my family, my co-workers and to the whole ESA team (RB)

Preface

In recent years, as an effect of technological innovation of surgical instruments and devices, and also the increasing number of surgical techniques being proposed and validated in the literature, we have seen a considerable expansion of the surgical options for the treatment of shoulder disorders. Furthermore, advancing globalisation, the growth of web-based scientific dissemination and education, and the constant and systematic training and information activities carried out by scientific societies and the research world have all contributed to an overall improvement in the level of theoretical and practical knowledge in the field of shoulder surgery, with the result that there is now very little difference, in terms of quality and surgical efficiency, between the health systems of different countries. All this has contributed to an exponential increase in the number of shoulder repair, reconstruction and replacement surgeries performed every year in the world. Inevitably, this has brought an increase in the number of failures and complications, which have also become more complex to manage, especially in patients with repeated failures. For this reason, the various scientific societies with an interest in shoulder disorders have recently become inclined to examine more closely the problem of shoulder surgery complications and failures, from different perspectives: prevention, diagnosis and management. In particular, European Shoulder Associates (ESA), the ESSKA section devoted to shoulder disorders and surgery, decided that its first biennial meeting should focus on this important and highly topical issue. This meeting, entitled “Management of Failed Shoulder Surgery”, was held in Rome on 2–3 October 2015, and this book springs from that event.

We are particularly pleased and proud to have the task of presenting this monograph, which has the same title as the Rome congress, as it offers readers a valuable opportunity to explore aspects of a subject that is both complex and controversial. This is the first time in over a decade that a book has been published that deals exclusively and exhaustively with the management of failed shoulder surgeries, aiming to help us recognise these events, understand why they occur and find successful solutions.

The book is structured in the same way as the Rome meeting. There are five parts, each focusing on a specific area of shoulder surgery: glenohumeral instability surgery, sports injury surgery, standard anatomical shoulder replacement, reverse shoulder replacement and rotator cuff surgery. Each part is made up of chapters that analyse problems and solutions related to complications and failures specific to each surgical procedure. The parts also contain

case studies illustrating the diagnostic and therapeutic approach used by the authors to manage particularly complex cases.

All the speakers at the Rome congress agreed to take part in this book project, and all have provided a contribution, reviewed and updated, on the subject of their particular presentation. Our sincere thanks go to all of them. The enthusiastic support of all the authors has been crucial, helping us to produce volume of great scientific quality. We are confident that readers will appreciate the format the authors have chosen for their chapters, based mainly on a decision-making and problem-solving approach.

Finally, we thank ESSKA's Board for approving and supporting this initiative, and all those at Springer for their great professionalism, and also for the book's excellent quality in both graphic and editorial terms.

Rome, Italy
Rome, Italy
Zaragoza, Spain
Bielsko-Biala, Poland

Giuseppe Milano
Andrea Grasso
Angel Calvo
Roman Brzóška

Contents

Part I Failed Shoulder Instability Repair

- 1 Shoulder Instability Repair: Why It Fails** 3
Emilio Calvo, Gia Rodriguez-Vaquero, and David Haeni
- 2 How to Manage Failed Anterior Arthroscopic Repair** 15
Vito Bongiorno
- 3 How to Manage Failed Anterior Open Repair** 19
Boris Poberaj
- 4 Management of Failed Posterior and Multidirectional Instability Repair** 23
Roman Brzóška, Hubert Laprus, Paweł Ranosz, Patryk Kłapocz, and Tomasz Rynkiewicz
- 5 Failed Arthroscopic Anterior Instability Repair: Case Example** 35
Emmanouil Antonogiannakis, Emmanouil Brilakis, and Anastasios Deligeorgis
- 6 Failure Posterior Instability Repair: Case Example** 41
Angel Calvo, Nestor Zurita, Alfredo Rodríguez, and Pablo Carnero
- 7 Failed Open Anterior Instability Repair: Case Example** 45
Ettore Taverna and Vincenzo Guarrella

Part II Failed Biceps and Acromioclavicular Joint Treatment

- 8 The Disabled Throwing Shoulder: When and How to Operate** 51
Nuno Gomes, Ricardo Aido, and Joana Gomes
- 9 How to Manage Failed Slap Repair** 67
Néstor Zurita and Angel Calvo Díaz
- 10 Biceps Disorders: When and How to Operate** 73
Sebastian Kwisda and Matthias Flury

11 Acromioclavicular Joint Instability: When and How to Operate	85
Theresa Diermeier, Felix Dyrna, Andreas B. Imhoff, and Knut Beitzel	
12 Failed Slap Repair: Case Example	91
Antonio Cartucho	
13 Failed AC Joint Treatment: Case Examples	97
Francesco Franceschi and Marco Spoliti	

Part III Failed Anatomical Shoulder Arthroplasty

14 Anatomical Shoulder Arthroplasty: How It Works	103
Thierry Joudet, Christophe Charousset, and The Shoulder Friends Group	
15 Anatomical Shoulder Arthroplasty: Why It Fails	111
Pascal Gleyze, Nikolaos Tzanakakis, and Constantina Moraiti	
16 Total Shoulder Arthroplasty: How to Prevent Failure	121
Philipp R. Heuberger and Leo Pauzenberger	
17 Anatomical Shoulder Arthroplasty: How to Manage Failure	127
Friso A. de Boer and Pol E. Huijsmans	
18 Failed Anatomical Shoulder Arthroplasty: Case Example 1	135
Radovan Mihelic and Zdravko Jotanovic	
19 Failed Anatomical Shoulder Arthroplasty: Case Example 2	139
Ladislav Kovacic	
20 Failed Anatomical Shoulder Arthroplasty: Case Example 3	147
Pietro Simone Randelli, Chiara Fossati, and Alessandra Menon	

Part IV Failed Reverse Shoulder Arthroplasty

21 Reverse Shoulder Arthroplasty: How It Works	155
Bruno Toussaint and Jérôme Bahurel	
22 Reverse Shoulder Arthroplasty: Why It Fails	163
Vladimir Senekovič	
23 Reverse Shoulder Arthroplasty: How to Prevent Failure	169
Eric Petroff and Johnathan Edwards	
24 Reverse Shoulder Arthroplasty: How to Manage Failure	177
Srinath Kamineni	

25 Failed Reverse Shoulder Arthroplasty:	
Case Example 1	189
Berte Bøe and Tom C. Ludvigsen	
26 Failed Reverse Shoulder Arthroplasty:	
Case Example 2	193
Maristella F. Saccomanno, Cristina Rossi, Carmine Latte, and Andrea Grasso	
Part V Failed Rotator Cuff Repair	
27 Rotator Cuff Repair: Why It Fails	201
Olaf Lorbach	
28 How to Manage Failed Rotator Cuff Repair:	
Arthroscopic Revision Surgery	207
Alexander Otto, Knut Beitzel, and Andreas B. Imhoff	
29 How to Manage Failed Rotator Cuff Repair:	
Biologic Augmentation	219
Paolo Avanzi, Luca Dei Giudici, Antonio Gigante, and Claudio Zorzi	
30 How to Manage Failed Rotator Cuff Repair:	
Latissimus Dorsi Transfer	229
Enrico Gervasi, Enrico Sebastiani, and Enrico Cautero	
31 Failed Rotator Cuff Repair: Case Example 1	235
Taner Gunes, Umut Akgun, and Recep Kurnaz	
32 Failed Rotator Cuff Repair: Case Example 2	239
Maristella F. Saccomanno and Giuseppe Milano	
Index	245

Part I

Failed Shoulder Instability Repair



Shoulder Instability Repair: Why It Fails

1

Emilio Calvo, Gia Rodriguez-Vaquero,
and David Haeni

1.1 Introduction

The glenohumeral (GH) joint is the least constrained joint in the body and allows a wide range of motion (ROM). On the other hand, it is more susceptible to high rates of instability. In the United States, the incidence of shoulder dislocations is 23 per 100,000 person-years, with the highest rates in adults in their 20s [1]. Anterior shoulder instability is the most frequent, and it is estimated that it affects 1.7% of the population. Current surgical techniques treating anterior shoulder instability are classified in soft tissue and bone augmentation procedures [2]. In the past, the open Bankart repair was considered the “gold standard,” obtaining satisfactory surgical results since its first description [3]. Concerns

regarding this technique were related to the extensive non-sparing subscapularis approach, immediate postoperative pain, loss of external rotation, and secondary osteoarthritis [4]. With the advent of new techniques and the development of new implants, the arthroscopic Bankart repair showed similar recurrence rates and functional outcomes than the open technique [5, 6]. Despite these results, reported recurrence rates after open or arthroscopic Bankart repair ranges between 5% and 15% [7, 8]. Bone augmentation procedures are usually preferred in young and active patients with recurrent shoulder dislocation in the presence of bone loss (Hill-Sachs lesions and/or bony Bankart) [9]. Recently, a prospective multicenter study found that the Latarjet procedure (open or arthroscopic) improves significantly shoulder function [10].

The main complication after surgical shoulder stabilization (whether open or arthroscopic) is recurrent instability. Revision instability surgery is usually a challenge, and patients with postoperative shoulder instability should be carefully evaluated not only to diagnose the failure but also to clearly identify the underlying causes that determined the outcome and to establish a successful therapeutic strategy [7, 8]. Careful preoperative evaluation is critical for the selection of the best treatment. The clinician must collect detailed information about the cause of the instability, the number and frequency of

E. Calvo (✉)
Shoulder and Elbow Reconstructive Surgery Unit,
Department of Orthopedic Surgery and
Traumatology, Fundacion Jimenez Diaz, Universidad
Autónoma, Madrid, Spain
e-mail: ecalvo@fjd.es

G. Rodriguez-Vaquero
Upper Extremity Surgery Unit, Department of
Orthopedic Surgery and Traumatology, Hospital
General de Villalba, Madrid, Spain
e-mail: gia.rodriguez@fjd.es

D. Haeni
Shoulder and Elbow Surgery Unit, LEONARDO,
Hirslanden Klinik Birshof, Basel, Switzerland

episodes, the degree of trauma necessary for recurrence, the arm position at the time of the initial injury, and the arm position that provokes symptoms [11].

Any patient with surgical treatment failure after shoulder stabilization can be classified in at least one of the following groups (Table 1.1). The first group is composed of patients in whom the problem was misdiagnosed, either because surgery was not indicated (i.e., voluntary instability), because the specific joint abnormalities to be corrected at surgery were not precisely identified, or because the direction of instability was not adequately understood (i.e., patients with multidirectional instability treated only for anterior instability). Patient-related risk factors may also increase the risk of postoperative recurrence and should be taken into account in the decision-making process in order to offer the best surgical treatment for every patient. Another group of subjects includes properly diagnosed patients in whom the treatment was inadequate, in terms of procedure selection or technical execution. Obviously, there could also be patients with combined misdiagnosis and inadequate treatment leading to surgical treatment failure. The last group includes those patients that were properly diagnosed, and in whom joint abnormalities were recognized and corrected with the optimal procedure, but who

suffered a new trauma causing postoperative dislocation or subluxation [12, 13].

1.1.1 Misdiagnosis

In order to properly address failed surgical treatment, it is essential first to clearly identify if surgery was indicated. Voluntary GH dislocation tends to occur in the young adult, and it is sometimes related to emotional and psychological problems. Huber et al. showed that voluntary subluxation in the childhood shows usually a favorable long-term outcome with conservative treatment and that is not associated with osteoarthritis [14]. Therefore, recurrent postoperative instability in this setting should be managed conservatively with physical therapy.

Once voluntary instability is ruled out, and considering that instability interferes with patient's activities, the most challenging issue is identifying which is the suitable surgical technique for each patient. For this purpose, it is crucial to recognize the direction of the instability, as well as the abnormalities responsible for recurrence to be addressed. Zabinski et al. [15] reported the comparative results of revision instability surgery in two groups of patients diagnosed of anterior and multidirectional instability, respectively. They found that persistent Bankart lesions were less common and the presence of hyperlaxity was almost constant in those diagnosed of multidirectional instability and concluded that while revision shoulder stabilization is a reliable procedure for patients who have recurrent anterior instability, it is unpredictable in patients who have multidirectional instability with surgical failure and reoperation occurring frequently.

Clinical history and meticulous physical examination allow identifying the direction of the instability, providing evidence about the possible causes of failure and potential associated lesions [16]. Physical examination should be performed always comparing the index shoulder to the contralateral side. The degree of instability (dislocation, subluxations, or apprehension) is also important information. The apprehension test is

Table 1.1 Causes of failure of anterior shoulder stabilization

Misdiagnosis
– Surgical treatment not indicated
– Anatomical abnormalities not identified
– Direction of instability
Patient-related risk factors
– Age, sex
– Number of dislocation
– Type of sport
– Concomitant/trigger disease: Epilepsy, Ehlers-Danlos disease
Surgery-related risk factors
– Technical errors
– Inadequate treatment
– Implant failure: Anchor or graft related
Trauma after surgery
Unknown causes

performed with the arm held at 0°–90°–140° abduction and is considered positive for anterior instability if the patient fears subluxation/dislocation or feels high discomfort during the maneuver. The sulcus sign is considered positive if during inferior traction of the shoulder held in neutral position a “sulcus” between acromion and humeral head is appreciated. A positive painful jerk test suggests postero-inferior labrum tear and a surgical repair should be discussed with the patient [17].

Examination under anesthesia before any revision surgery can be useful since it may overcome the clinical examination limitation due to patient’s apprehension. Mechanical symptoms, such as catching or locking, may suggest a displaced labral tear, a loose body, or a large osseous defect that is engaging. Instability that occurs in the midrange of motion or during the sleep may indicate an osseous defect. Decreased ROM may be secondary to postoperative stiffness, chondrolysis, GH osteoarthritis, or excessive tension of the capsulolabral ligamentous complex. Loss of strength could be related to rotator cuff tear or neurological injury. Accurate rotator cuff testing should be performed, especially with regard to subscapularis muscle function in patients with previous open surgery. Sachs et al. [18] found that 23% of the patients undergoing open Bankart repair had a deficient subscapularis function and only 57% of them obtained good or excellent results after revision surgery.

Conventional radiography (CR) represents the first level of investigation in postoperative shoulder instability and should include outlet view, “true” anteroposterior view, and the axillary view. With the axillary view, we can evaluate anterior or posterior humeral head subluxation and the state bone graft healing.

Magnetic resonance imaging (MRI) with intra-articular contrast medium (MR arthrography, MRA) can be used both in presurgical and postsurgical care for shoulder instability giving a good assessment of capsulolabral-ligamentous complex and to evaluate postoperative recurrence or complication. MRA identifies soft tissue injuries, rotator cuff tears, humeral avulsion of the glenohumeral ligament (HAGL) lesions, capsu-

lolar lesions, chondral lesions, and laxity or rupture of the joint capsule better than standard MRI [19]. MRA in abduction and external rotation (ABER) position is useful to identify patients with atraumatic multidirectional instability. The presence of a layer of contrast medium between the humeral head and the anteroinferior glenohumeral ligament (AIGHL) (crescent sign) combined with a triangular-shaped space between the humeral head, AIGHL, and glenoid (triangle sign) has a sensitivity of 86% and specificity of 94% in diagnosing MDI [20].

Computed tomography (CT) can be used for bone evaluation and in cases in which CR does not give enough information about devices positioning. CT arthrography (CTA) is a valid alternative to MRA when susceptibility artifacts are present.

1.1.2 Patient-Related Failure

Several studies have attempted to establish the prognostic factors that may increase the risk of postoperative recurrence following surgical stabilization. Young age and participation in risk activities were identified as major prognostic factors in all of them in addition to the presence of bone defects [21–25]. Age at the first dislocation and male gender have been strongly correlated with a significantly higher risk of recurrent instability after a first dislocation, approaching 80% [21, 26]. Coherently to that, young male patients are more prone to recurrence after primary stabilization [11]. In a study of over 5900 patients, those younger than 20 years had a 12.6% risk of postoperative dislocation and a 7.7% revision rate after primary stabilization, compared to 5.5% and 2.8%, respectively, in patients older than 29 years of age [14]. When compared to adults, young patients usually have higher activity level, more compliant tissue, and decreased muscle bulk. Ninety percent of patients with recurrent dislocations after arthroscopic repair are male [16, 17].

The number of dislocations before stabilization, in addition to the number of previous surgeries, negatively correlates with postsurgical

success [27]. Wasserstein et al. [26] found that patients with three or more dislocations had double the risk for revision surgery and ten times the risk of re-dislocating. Patients with more than one stabilization procedure trended toward lower functional outcomes and less overall satisfaction [28]. These results are likely related to progressive damage tissue.

Collision athletes and contact overhead athletes are more frequently subject to higher energy trauma that can lead to shoulder dislocation and other injuries. In addition, postoperative return to collision sports is associated to a higher risk of new trauma and re-dislocation. Cho et al. [29] and Rhee et al. [30] reported higher instability recurrence rate in active athletes (17.2%) after arthroscopic Bankart repair. Even higher rates are reported in patients who practice collision sports (25–28%). Uhorchak et al. [31] reported outcomes of open Bankart repair, and they found a recurrence of 12% in collision and contact sports athletes. Castagna et al. [32] analyzed the effectiveness of arthroscopic Bankart repair in adolescent athletes who practiced overhead or contact sports at competitive level and reported higher recurrence rate in very high-energy contact sports (rugby) and in high-energy contact sports associated with overhead position of the arm (water polo). Other authors associated contact sports with higher risk of recurrence, but it does not seem to be a contraindication for arthroscopic Bankart repair [33, 34].

Calvo et al. [21] evaluated prospectively 61 patients treated arthroscopically with Bankart repair for recurrent anterior shoulder instability. They developed a risk score for failure of arthroscopic Bankart repair based upon an analysis of the factors that may determine the outcomes (level of satisfaction and degree of stability). Age younger than 28 years, ligamentous laxity, the presence of a fracture of the glenoid rim involving more than 15% of the articular surface, and postoperative participation in contact or overhead sports were associated with a higher risk of recurrence and scored 1, 1.5, and 1 point, respectively. Those patients with a total score of two or more points had a relative risk of recurrence of 43% and should be treated by open surgery. Later, Balg

et al. [22] developed the instability severity index score (ISIS) to predict the success of arthroscopic Bankart repair. The ISIS score ranges from 0 to 10, with higher scores predicting a higher risk of recurrence after stabilization. Six risk factors are considered that can predict a higher recurrence rate: age at the surgery (over or below 20), degree and type of preoperative sport, hyperlaxity, and bone loss studied on CR.

Epileptic seizures can cause shoulder dislocation and instability, but these patients follow a characteristic pattern of instability with peculiar structural lesions. Bühler and Gerber [35] studied 34 shoulders in which initial dislocation had been caused by an epileptic seizure. Fifty percent of them had anterior instability and 50% posterior instability. They also found a higher recurrence rate for anterior instability comparing with posterior instability (47 versus 12%) after primary repair. Most of them were associated to poor control of epilepsy disease. Thangarajah et al. [36] followed up 49 patients with recurrent instability with epilepsy for 15 years: 73% of them showed anterior instability, 15% posterior, and 10% multidirectional instability. Eighty percent of all patients showed bone loss. They identified bone loss and persistent postoperative epileptic seizures as the principal factors for recurrent instability. Epileptic medical control and bone block procedure are associated with lower rate of recurrence.

1.1.3 Inadequate Treatment: Anatomic Abnormalities and Technique of Stabilization

Shoulder stabilization surgery should be tailored to the patient and to the specific abnormalities existing in the shoulder. In a cohort of 32 patients surgically revised for recurrent anterior dislocation of the shoulder after surgical repair, Rowe et al. [37] found that an abnormality that had not been adequately addressed and explaining the recurrence could be identified in more than 85% of the patients with postoperative shoulder instability. Moreover, Meeham and Petersen [12] proved in a similar investigation that in almost

half of the cases there is more than one lesion. Therefore, in revision instability surgery, it is crucial to study and identify the specific anatomic abnormalities responsible for the poor outcome. The most frequent abnormalities that can lead to shoulder instability surgery failure are the presence of non-repaired or medially repaired Bankart lesion (Fig. 1.1), poor capsulolabral tissue (Fig. 1.2) or hyperlaxity, and unaddressed bone defects (either on the glenoid or the humeral side) [12, 15, 37, 38].




Fig. 1.1 Left shoulder. Arthroscopic view from the posterior portal: medially repaired Bankart lesion




Fig. 1.2 Left shoulder. Arthroscopic view from the anterosuperior portal: poor capsulolabral tissue

Insufficient labral detachment followed by anatomic re-fixation of a medially healed labrum after multiple episodes of recurrence is probably the most common error during Bankart repair. Anterior labro-ligamentous periosteal sleeve avulsion (ALPSA) lesions have been identified as a risk factor for recurrence comparing with isolated Bankart lesion [27, 39] (Fig. 1.3). This lesion is present more frequently in patients with high number of dislocations. The reason of recurrence after repair may be related to the poor quality of capsulolabral tissue, due to progressive damage. Underestimation of HAGL lesions is also responsible for persistent postoperative instability (Fig. 1.4). A high index of suspicion is necessary to identify and repair this lesion, which can appear in 9% of anterior instability cases [40, 41]. Cases of first-time shoulder dislocation without Bankart lesion and no multidirectional laxity can show a high incidence of HAGL lesions [42].

Poorly positioned anchors have also been associated with recurrence of instability [43]. The number of suture anchors used for primary arthroscopic Bankart repair plays also an important role in the recurrence rate, and three or more anchors are usually recommended in most common cases of anterior shoulder instability [27, 44, 45]. With regard to the type of anchors, data showed no difference in recurrence rate between




Fig. 1.3 Right shoulder. Arthroscopic view from the anterosuperior portal: ALPSA lesion



Fig. 1.4 Right shoulder. Arthroscopic view from the posterior portal: HAGL lesion

metal or biodegradable devices [46]. However, a significant difference was found between patients in whom knot-tying and knotless suture anchors were used, with higher rate of recurrence using knotless anchors [47].

One of the most commonly known mistakes includes failure to recognize and address capsular laxity during arthroscopic repair [12, 15, 37, 38]. Hyperlaxity and instability may be coexisting conditions. The difference between instability and hyperlaxity needs to be assessed preoperatively and influences the therapeutic decision. After multiple shoulder dislocations, anterior capsular tissue may be stretched and becomes redundant [12, 28]. Bigliani et al. [48] demonstrated that anterior capsular stretching can occur with or without labral detachment. Rowe et al. [37] showed that 83% of patients with recurrent dislocations after surgical repair had significant capsular laxity.

It is known that the recurrence of instability is significantly higher in patients with anterior glenoid bone defects [21, 23]. Imaging studies are essential for the evaluation of patients with recurrent instability, since it allows the identification and quantification of glenoid bone loss and other possible articular abnormalities. While CR is considered important for bone loss assessment and many different radiographic views have been proposed, CT scan is considered the ideal method

to quantify both glenoid and humeral head bone defects. Several authors [49–51] described the glenoid osseous defect as being located anteriorly at approximately the 3 o'clock position (in the right shoulder) and extending toward inferiorly. 3D CT scan with humeral head subtraction facilitates quantification of glenoid bone defect related to the total area and depth of the defect. Glenoid bone defect over 20% have been strongly associated with high risk of recurrence of instability after Bankart repair [45, 52], but Calvo et al. [21] demonstrated that a glenoid bone defect involving 15% of the articular surface represented a higher risk of postoperative failure. Yamamoto et al. [53] introduced the concept of the “glenoid track” determining whether a bipolar lesion was significant. The “glenoid track” concept offers the surgeon the possibility to predict engagement, based on size and morphology lesions [49]. The critical size of a Hill-Sachs lesion is thought to be a volume over 250 mm³, defined as “large Hill-Sachs lesion” [54]. Recent clinical evidence supports the “on-track” versus “off-track” model in predicting failure of isolated Bankart repair in shoulders with bipolar bone loss [55, 56].

Bone augmentation procedures are preferred to address bone defects, and Latarjet is regarded as the gold standard technique for this condition [57, 58]. Walch et al. [9] conducted a study of 68 shoulders after open Latarjet and reported a recurrence rate of 5.9% after a mean 20-year follow-up. Young and Rockwood [59] studied a population of 39 patients with painful instability after shoulder stabilization performed with an open Bristow procedure and attributed the recurrences to the presence of capsular redundancy in 23 (59%) cases. Other investigations have also found labral defects and capsular elongation at arthroscopic revision of recurrent instability in patients previously operated with bone block procedures [60, 61]. Arthroscopic examination was considered extremely useful in identifying these abnormalities, and labral re-fixation with capsular plication was recommended to stabilize the shoulder [61]. However, other authors have attributed postoperative instability after Latarjet to complications related to the coracoid

graft, either due to malposition, malunion, or nonunion [62]. Gasbarro et al. [63] analyzed the reasons for failure after coracoid transfer procedures in a cohort of 83 patients and considered too inferior or too medial graft placement to be a risk factor for recurrence, as well as single screw fixation of the coracoid graft. Nonunion is a well-known complication after Latarjet procedure that can involve over 9% of the patients, but it has not been clearly associated with a higher risk of recurrence [64]. The coracoid graft can also show osteolysis at its upper half, but this complication does not seem to be correlated with postoperative recurrence either [65–67].

Eden-Hybinette, either open or arthroscopic, has been regarded as the elective technique for failed Latarjet, especially in patients with bone defects [68, 69]. Lunn et al. [70] reported the first series of the Eden-Hybinette revision procedure in a cohort of 46 patients with failed Latarjet and found different risk factors for recurrence such as malposition, lysis, or avulsion of the coracoid graft. Interestingly, the authors identified that ligamentous laxity was present in 14 patients and for the first time incriminated subscapularis weakness as a reason for failure 10 patients (5 patients had a complete rupture of the subscapularis tendon). Calvo et al. [71] reported a series of 11 patients who underwent revision surgery for recurrent instability after Latarjet stabilization. The technique used was based on the specific anatomic abnormalities found at arthroscopy: the coracoid graft inadequately positioned was repositioned with open surgery in three cases; extraarticular capsular reinforcement was performed in four shoulders that showed hyperlaxity or poor capsulolabral tissue and no severe bone defect, while arthroscopic Eden-Hybinette was used in four shoulders with humeral or glenoid bone defects and a nonviable coracoid graft.

Boileau et al. [44] pointed out the role of certain Hill-Sachs defects in the recurrence following surgical stabilization. Recently, Locher et al. [72] assessed the impact of “off-track” Hill-Sachs lesions in a study of 254 patients with anterior instability managed with a Bankart repair. The

authors demonstrated that Hill-Sachs “off-track” lesions constitute an important risk factor for recurrence of instability after arthroscopic Bankart repair and need of revision surgery compared to “on-track” defects. “Remplissage” is regarded as the procedure of choice for those patients with “off-track” defects, albeit Latarjet procedure could be a valid alternative in shoulders with “off-track” Hill-Sachs lesions by increasing the articular surface area. However, Millet et al. [11] demonstrated that the presence of “off-track” Hill-Sachs lesions increases the risk for persistent engagement after surgery also after stabilization with the Latarjet technique.

Based on the few comparative studies reported, there is no evidence on the superiority of open or arthroscopic stabilization in terms of recurrence, and the fact that arthroscopic stabilization represents an independent risk for recurrence cannot be sustained. Mohtadi et al. [73] carried out a prospective study of 196 patients randomized to undergo open or arthroscopic soft tissue stabilization and concluded that although there were no differences concerning postoperative quality of life, the recurrence rate was superior after arthroscopic surgery. However, Fabbriani et al. [74], in a study with a similar design, failed to find differences between the two therapeutic approaches and noticed that the group treated arthroscopically showed superior postoperative mobility over the open group. Moreover, Archetti Netto et al. [75] reported lower failure rates, higher mobility, and fewer complications after arthroscopic Bankart stabilization. With regard to coracoid transfer procedures, there are not published studies on the superiority of the arthroscopic versus the traditional open approach. Arthroscopic surgery is very helpful in identifying articular abnormalities to be amended, and arthroscopic revision stabilization provides satisfactory results [62]. The technique allows direct visualization of the pathology that may be responsible for recurrence, including unexpected causes that can be corrected during the same procedure, such as loose bodies, rotator cuff tears, or chondral lesions [76, 77].

1.1.4 New Trauma

Traumatic injuries to the surgically repaired shoulder are one of the biggest contributors to recurrence. As the majority of those affected are young with initial injuries often due to athletic activities, return to collision sport or overhead throwing sports predisposes this population to reinjury. Tauber et al. [38] reviewed 41 patients and found that 85% of initial shoulder dislocations and 59% of re-dislocations after surgical stabilization were traumatic.

Conclusion

Key factors for successful surgical shoulder stabilization are adequate patient selection, precise surgical technique selection and fulfillment, identification and correction of all joint abnormalities, and integration of patient and surgeon expectations. For this purpose, we must be able to correctly answer the following questions: what are the characteristics of the patient? Which shoulder injuries should be treated? Did the patient have a new trauma responsible for recurrence? Despite all known risk factors for recurrence of instability, there are cases in which it is not possible to establish the cause of primary repair failure.

References

- Zacchilli MA, Owens BD. Epidemiology of shoulder dislocations presenting to emergency Departments in the United States. *J Bone Joint Surg Am.* 2010;92:542–9. <https://doi.org/10.2106/JBJS.I.00450>.
- Dumont GD, Russell RD, Robertson WJ. Anterior shoulder instability: a review of pathoanatomy, diagnosis and treatment. *Curr Rev Musculoskelet Med.* 2011;4:200–7. <https://doi.org/10.1007/s12178-011-9092-9>.
- Bankart ASB. Recurrent or habitual dislocation of the shoulder-joint. *Br Med J.* 1923;2:1132–3.
- Rosenberg BN, Richmond JC, Levine WN. Long-term followup of Bankart reconstruction. Incidence of late degenerative glenohumeral arthrosis. *Am J Sports Med.* 1995;23:538–44. <https://doi.org/10.1177/036354659502300504>.
- Hobby J, Griffin D, Dunbar M, Boileau P. Is arthroscopic surgery for stabilisation of chronic shoulder instability as effective as open surgery? A systematic review and meta-analysis of 62 studies including 3044 arthroscopic operations. *J Bone Joint Surg Br.* 2007;89:1188–96. <https://doi.org/10.1302/0301-620X.89B9.18467>.
- Freedman KB, Smith AP, Romeo AA, et al. Open Bankart repair versus arthroscopic repair with trans-glenoid sutures or bioabsorbable tacks for recurrent anterior instability of the shoulder: a meta-analysis. *Am J Sports Med.* 2004;32:1520–7. <https://doi.org/10.1177/0363546504265188>.
- Franceschi F, Longo UG, Ruzzini L, et al. Arthroscopic salvage of failed arthroscopic Bankart repair: a prospective study with a minimum follow-up of 4 years. *Am J Sports Med.* 2008;36:1330–6. <https://doi.org/10.1177/0363546508314403>.
- Millar NL, Murrell GAC. The effectiveness of arthroscopic stabilisation for failed open shoulder instability surgery. *J Bone Joint Surg Br.* 2008;90:745–50. <https://doi.org/10.1302/0301-620X.90B6.20018>.
- Walch G. Chronic anterior instability of the shoulder in adults. Summary. *Rev Chir Orthop Reparatrice Appar Mot.* 2000;86(Suppl 1):147–9.
- Nourissat G, Neyton L, Metais P, et al. Functional outcomes after open versus arthroscopic Latarjet procedure: a prospective comparative study. *Orthop Traumatol Surg Res.* 2016;102:S277–9. <https://doi.org/10.1016/j.otsr.2016.08.004>.
- Millett PJ, Clavert P, Warner JJP. Open operative treatment for anterior shoulder instability: when and why? *J Bone Joint Surg Am.* 2005;87:419–32. <https://doi.org/10.2106/JBJS.D.01921>.
- Meehan RE, Petersen SA. Results and factors affecting outcome of revision surgery for shoulder instability. *J Shoulder Elbow Surg.* 2005;14:31–7. <https://doi.org/10.1016/j.jse.2004.05.005>.
- Sisto DJ. Revision of failed arthroscopic bankart repairs. *Am J Sports Med.* 2007;35:537–41. <https://doi.org/10.1177/0363546506296520>.
- Huber H, Gerber C. Voluntary subluxation of the shoulder in children. A long-term follow-up study of 36 shoulders. *J Bone Joint Surg Br.* 1994;76:118–22.
- Zabinski SJ, Callaway GH, Cohen S, Warren RF. Revision shoulder stabilization: 2- to 10-year results. *J Shoulder Elbow Surg.* 1999;8:58–65.
- Johansson K. Multidirectional instability of the glenohumeral joint: an unstable classification resulting in uncertain evidence-based practice. *Br J Sports Med.* 2016;50:1105–6. <https://doi.org/10.1136/bjsports-2016-096094>.
- Kim S-H, Park J-C, Park J-S, Oh I. Painful jerk test: a predictor of success in nonoperative treatment of posteroinferior instability of the shoulder. *Am J Sports Med.* 2004;32:1849–55.
- Sachs RA, Williams B, Stone ML, et al. Open Bankart repair: correlation of results with postoperative subscapularis function. *Am J Sports Med.* 2005;33:1458–62. <https://doi.org/10.1177/0363546505275350>.
- Probyn LJ, White LM, Salonen DC, et al. Recurrent symptoms after shoulder instability repair: direct MR arthrographic assessment—correlation with second-look surgical evaluation. *Radiology.* 2007;245:814–23. <https://doi.org/10.1148/radiol.2453061329>.

20. Schaeffeler C, Waldt S, Bauer JS, et al. MR arthrography including abduction and external rotation images in the assessment of atraumatic multidirectional instability of the shoulder. *Eur Radiol.* 2014;24:1376–85. <https://doi.org/10.1007/s00330-014-3133-x>.
21. Calvo E, Granizo JJ, Fernández-Yruegas D. Criteria for arthroscopic treatment of anterior instability of the shoulder: a prospective study. *J Bone Joint Surg Br.* 2005;87:677–83. <https://doi.org/10.1302/0301-620X.87B5.15794>.
22. Balg F, Boileau P. The instability severity index score. A simple pre-operative score to select patients for arthroscopic or open shoulder stabilisation. *J Bone Joint Surg Br.* 2007;89:1470–7. <https://doi.org/10.1302/0301-620X.89B11.18962>.
23. Ahmed I, Ashton F, Robinson CM. Arthroscopic Bankart repair and capsular shift for recurrent anterior shoulder instability: functional outcomes and identification of risk factors for recurrence. *J Bone Joint Surg Am.* 2012;94:1308–15. <https://doi.org/10.2106/JBJS.J.01983>.
24. Voos JE, Livermore RW, Feeley BT, et al. Prospective evaluation of arthroscopic bankart repairs for anterior instability. *Am J Sports Med.* 2010;38:302–7. <https://doi.org/10.1177/0363546509348049>.
25. Flinkkilä T, Hyvönen P, Ohtonen P, Leppilähti J. Arthroscopic Bankart repair: results and risk factors of recurrence of instability. *Knee Surg Sports Traumatol Arthrosc.* 2010;18:1752–8. <https://doi.org/10.1007/s00167-010-1105-5>.
26. Wasserstein D, Dwyer T, Veillette C, et al. Predictors of dislocation and revision after shoulder stabilization in Ontario, Canada, from 2003 to 2008. *Am J Sports Med.* 2013;41:2034–40. <https://doi.org/10.1177/0363546513492952>.
27. Randelli P, Ragone V, Carminati S, Cabitza P. Risk factors for recurrence after Bankart repair a systematic review. *Knee Surg Sports Traumatol Arthrosc.* 2012;20:2129–38. <https://doi.org/10.1007/s00167-012-2140-1>.
28. Marquardt B, Garmann S, Schulte T, et al. Outcome after failed traumatic anterior shoulder instability repair with and without surgical revision. *J Shoulder Elbow Surg.* 2007;16:742–7. <https://doi.org/10.1016/j.jse.2007.02.132>.
29. Cho NS, Hwang JC, Rhee YG. Arthroscopic stabilization in anterior shoulder instability: collision athletes versus noncollision athletes. *Arthroscopy.* 2006;22:947–53. <https://doi.org/10.1016/j.arthro.2006.05.015>.
30. Rhee YG, Ha JH, Cho NS. Anterior shoulder stabilization in collision athletes: arthroscopic versus open Bankart repair. *Am J Sports Med.* 2006;34:979–85. <https://doi.org/10.1177/0363546505283267>.
31. Uhorchak JM, Arciero RA, Huggard D, Taylor DC. Recurrent shoulder instability after open reconstruction in athletes involved in collision and contact sports. *Am J Sports Med.* 2000;28:794–9. <https://doi.org/10.1177/03635465000280060501>.
32. Castagna A, Delle Rose G, Borroni M, et al. Arthroscopic stabilization of the shoulder in adolescent athletes participating in overhead or contact sports. *Arthroscopy.* 2012;28:309–15. <https://doi.org/10.1016/j.arthro.2011.08.302>.
33. Mazzocca AD, Brown FM, Carreira DS, et al. Arthroscopic anterior shoulder stabilization of collision and contact athletes. *Am J Sports Med.* 2005;33:52–60. <https://doi.org/10.1177/0363546504268037>.
34. Grana WA, Buckley PD, Yates CK. Arthroscopic Bankart suture repair. *Am J Sports Med.* 1993;21:348–53. <https://doi.org/10.1177/036354659302100304>.
35. Bühler M, Gerber C. Shoulder instability related to epileptic seizures. *J Shoulder Elbow Surg.* 2002;11:339–44. <https://doi.org/10.1067/mse.2002.124524>.
36. Thangarajah T, Lambert S. The management of recurrent shoulder instability in patients with epilepsy: a 15-year experience. *J Shoulder Elbow Surg.* 2015;24:1723–7. <https://doi.org/10.1016/j.jse.2015.04.008>.
37. Rowe CR, Zarins B, Ciullo JV. Recurrent anterior dislocation of the shoulder after surgical repair. Apparent causes of failure and treatment. *J Bone Joint Surg Am.* 1984;66:159–68.
38. Tauber M, Resch H, Forstner R, et al. Reasons for failure after surgical repair of anterior shoulder instability. *J Shoulder Elbow Surg.* 2004;13:279–85. <https://doi.org/10.1016/S1058274604000254>.
39. Porcellini G, Campi F, Pegreffo F, et al. Predisposing factors for recurrent shoulder dislocation after arthroscopic treatment. *J Bone Joint Surg Am.* 2009;91:2537–42. <https://doi.org/10.2106/JBJS.H.01126>.
40. Wolf EM, Siparsky PN. Glenoid avulsion of the glenohumeral ligaments as a cause of recurrent anterior shoulder instability. *Arthroscopy.* 2010;26:1263–7. <https://doi.org/10.1016/j.arthro.2010.06.005>.
41. Flury M, Rickenbacher D, Audigé L. Arthroscopic treatment of anterior shoulder instability associated with a HAGL lesion—a case series. *J Shoulder Elbow Surg.* 2016;25:1989–96. <https://doi.org/10.1016/j.jse.2016.02.030>.
42. Savoie FH, Holt MS, Field LD, Ramsey JR. Arthroscopic management of posterior instability: evolution of technique and results. *Arthroscopy.* 2008;24:389–96. <https://doi.org/10.1016/j.arthro.2007.11.004>.
43. Kim S-H, Ha K-I, Kim Y-M. Arthroscopic revision Bankart repair: a prospective outcome study. *Arthroscopy.* 2002;18:469–82. <https://doi.org/10.1053/jars.2002.32230>.
44. Boileau P, Villalba M, Héry J-Y, et al. Risk factors for recurrence of shoulder instability after arthroscopic Bankart repair. *J Bone Joint Surg Am.* 2006;88:1755–63. <https://doi.org/10.2106/JBJS.E.00817>.
45. An VVG, Sivakumar BS, Phan K, Trantalis J. A systematic review and meta-analysis of clinical and patient-reported outcomes following two procedures for recurrent traumatic anterior instability of the shoulder: Latarjet procedure vs. Bankart repair. *J Shoulder Elbow Surg.* 2016;25:853–63. <https://doi.org/10.1016/j.jse.2015.11.001>.

46. Thal R. Knotless suture anchor: arthroscopic bankart repair without tying knots. *Clin Orthop*. 2001;390:42–51.
47. Cho NS, Lubis AMT, Ha JH, Rhee YG. Clinical results of arthroscopic bankart repair with knot-tying and knotless suture anchors. *Arthroscopy*. 2006;22:1276–82. <https://doi.org/10.1016/j.arthro.2006.07.005>.
48. Bigliani LU, Kelkar R, Flatow EL, et al. Glenohumeral stability. Biomechanical properties of passive and active stabilizers. *Clin Orthop*. 1996;330:13–30.
49. Griffith JF, Antonio GE, Tong CWC, Ming CK. Anterior shoulder dislocation: quantification of glenoid bone loss with CT. *AJR Am J Roentgenol*. 2003;180:1423–30. <https://doi.org/10.2214/ajr.180.5.1801423>.
50. Saito H, Itoi E, Sugaya H, et al. Location of the glenoid defect in shoulders with recurrent anterior dislocation. *Am J Sports Med*. 2005;33:889–93. <https://doi.org/10.1177/0363546504271521>.
51. Ji J-H, Kwak D-S, Yang P-S, et al. Comparisons of glenoid bony defects between normal cadaveric specimens and patients with recurrent shoulder dislocation: an anatomic study. *J Shoulder Elbow Surg*. 2012;21:822–7. <https://doi.org/10.1016/j.jse.2011.10.016>.
52. Shibata H, Gotoh M, Mitsui Y, et al. Risk factors for shoulder re-dislocation after arthroscopic Bankart repair. *J Orthop Surg*. 2014;9:53. <https://doi.org/10.1186/s13018-014-0053-z>.
53. Yamamoto N, Itoi E, Abe H, et al. Contact between the glenoid and the humeral head in abduction, external rotation, and horizontal extension: a new concept of glenoid track. *J Shoulder Elbow Surg*. 2007;16:649–56. <https://doi.org/10.1016/j.jse.2006.12.012>.
54. Hartzler RU, Bui CNH, Jeong WK, et al. Remplissage of an off-track Hill-Sachs lesion is necessary to restore biomechanical glenohumeral joint stability in a bipolar bone loss model. *Arthroscopy*. 2016;32:2466–76. <https://doi.org/10.1016/j.arthro.2016.04.030>.
55. Di Giacomo G, Itoi E, Burkhart SS. Evolving concept of bipolar bone loss and the Hill-Sachs lesion: from “engaging/non-engaging” lesion to “on-track/off-track” lesion. *Arthroscopy*. 2014;30:90–8. <https://doi.org/10.1016/j.arthro.2013.10.004>.
56. Tokish JM, Shaha JS, Cook JB, et al. Predictive value and clinical validation of the “on-track” vs. “off-track” concept in bipolar bone loss in anterior glenohumeral instability. *Orthop J Sports Med*. 2015;3:2325967115S00089. <https://doi.org/10.1177/2325967115S00089>.
57. Anderl W, Krieglleder B, Heuberger PR. All-arthroscopic implant-free iliac crest bone grafting: new technique and case report. *Arthroscopy*. 2012;28:131–7. <https://doi.org/10.1016/j.arthro.2011.10.027>.
58. Walch G. Luxation récidivante antérieure de l'épaule. *Rev Chir Orthop*. 1991;77:177–91.
59. Young DC, Rockwood CA. Complications of a failed Bristow procedure and their management. *J Bone Joint Surg Am*. 1991;73:969–81.
60. Cuéllar A, Cuéllar R, de Heredia PB. Arthroscopic revision surgery for failure of open Latarjet technique. *Arthroscopy*. 2017;33:910–7. <https://doi.org/10.1016/j.arthro.2016.09.017>.
61. Castagna A, Garofalo R, Melito G, et al. The role of arthroscopy in the revision of failed Latarjet procedures. *Musculoskelet Surg*. 2010;94(Suppl 1):S47–55. <https://doi.org/10.1007/s12306-010-0060-0>.
62. Boileau P, Richou J, Lisai A, et al. The role of arthroscopy in revision of failed open anterior stabilization of the shoulder. *Arthroscopy*. 2009;25:1075–84. <https://doi.org/10.1016/j.arthro.2009.04.073>.
63. Gasbarro G, Giugale JM, Walch G, Lin A. Predictive surgical reasons for failure after coracoid process transfers. *Orthop J Sports Med*. 2016;4:2325967116676795. <https://doi.org/10.1177/2325967116676795>.
64. Gupta A, Delaney R, Petkin K, Lafosse L. Complications of the Latarjet procedure. *Curr Rev Musculoskelet Med*. 2015;8:59–66. <https://doi.org/10.1007/s12178-015-9258-y>.
65. Di Giacomo G, de Gasperis N, Costantini A, et al. Does the presence of glenoid bone loss influence coracoid bone graft osteolysis after the Latarjet procedure? A computed tomography scan study in 2 groups of patients with and without glenoid bone loss. *J Shoulder Elbow Surg*. 2014;23:514–8. <https://doi.org/10.1016/j.jse.2013.10.005>.
66. Di Giacomo G, Costantini A, de Gasperis N, et al. Coracoid graft osteolysis after the Latarjet procedure for anteroinferior shoulder instability: a computed tomography scan study of twenty-six patients. *J Shoulder Elbow Surg*. 2011;20:989–95. <https://doi.org/10.1016/j.jse.2010.11.016>.
67. Haeni DL, Opsomer G, Sood A, et al. Three-dimensional volume measurement of coracoid graft osteolysis after arthroscopic Latarjet procedure. *J Shoulder Elbow Surg*. 2017;26:484–9. <https://doi.org/10.1016/j.jse.2016.08.007>.
68. Giannakos A, Vezeridis PS, Schwartz DG, et al. All-arthroscopic revision eden-hybinette procedure for failed instability surgery: technique and preliminary results. *Arthroscopy*. 2017;33:39–48. <https://doi.org/10.1016/j.arthro.2016.05.021>.
69. Mizuno N, Denard PJ, Raiss P, et al. Long-term results of the Latarjet procedure for anterior instability of the shoulder. *J Shoulder Elbow Surg*. 2014;23:1691–9. <https://doi.org/10.1016/j.jse.2014.02.015>.
70. Lunn JV, Castellano-Rosa J, Walch G. Recurrent anterior dislocation after the Latarjet procedure: outcome after revision using a modified Eden-Hybinette operation. *J Shoulder Elbow Surg*. 2008;17:744–50. <https://doi.org/10.1016/j.jse.2008.03.002>.
71. Calvo E, Rojas D, Morcillo D, et al. Management of the failed Latarjet procedure. In: Boileau P, editor. *Shoulder concepts: arthroscopy, arthroplasty & fractures*. Montpellier: Lippincott Williams & Wilkins; 2012.
72. Locher J, Wilken F, Beitzel K, et al. Hill-Sachs off-track lesions as risk factor for recurrence of instability after arthroscopic bankart repair. *Arthroscopy*. 2016;32:1993–9. <https://doi.org/10.1016/j.arthro.2016.03.005>.
73. Mohtadi NGH, Chan DS, Hollinshead RM, et al. A randomized clinical trial comparing open and

- arthroscopic stabilization for recurrent traumatic anterior shoulder instability: two-year follow-up with disease-specific quality-of-life outcomes. *J Bone Joint Surg Am.* 2014;96:353–60. <https://doi.org/10.2106/JBJS.L.01656>.
74. Fabbriani C, Milano G, Demontis A, et al. Arthroscopic versus open treatment of Bankart lesion of the shoulder: a prospective randomized study. *Arthroscopy.* 2004;20:456–62. <https://doi.org/10.1016/j.arthro.2004.03.001>.
75. Archetti Netto N, Tamaoki MJS, Lenza M, et al. Treatment of Bankart lesions in traumatic anterior instability of the shoulder: a randomized controlled trial comparing arthroscopy and open techniques. *Arthroscopy.* 2012;28:900–8. <https://doi.org/10.1016/j.arthro.2011.11.032>.
76. Araghi A, Prasarn M, St Clair S, Zuckerman JD. Revision anterior shoulder repair for recurrent anterior glenohumeral instability. *Bull Hosp Jt Dis.* 2005;62:102–4.
77. Tamai K, Higashi A, Tanabe T, Hamada J. Recurrences after the open Bankart repair: a potential risk with use of suture anchors. *J Shoulder Elbow Surg.* 1999;8:37–41.



How to Manage Failed Anterior Arthroscopic Repair

2

Vito Bongiorno

2.1 Introduction

Arthroscopic Bankart repair is the most performed procedure for the treatment of post-traumatic capsulo-labral detachment [1, 2]. Although some authors still perform open Bankart repair claiming better results, the arthroscopic technique is considered the gold standard for anterior stabilization [3, 4].

Recurrence of instability after arthroscopic Bankart repair is increasing over years probably because of the large number of procedures done by surgeons performing few procedures per year (i.e., lack of adequate training) [5]. Another cause might be the excessive use of arthroscopic procedures in the attempt to treat lesions that needed to be treated differently [6, 7].

Management of failed arthroscopic repair for anterior shoulder instability needs to focus mainly on the reasons for failure. Except for new trauma occurring on the operated shoulder at least 9–12 months after primary surgery in a patient that has fully recovered, other causes may be wrong surgical technique or, more often, wrong indication. Accurate analysis of the preoperative imaging and clinical history of the patient can be very helpful in understanding the cause of failure.

2.2 Anterior Shoulder Instability: Spectrum of Disease

Traumatic anteroinferior dislocation of the shoulder causes a spectrum of lesions that determine the basis for further episodes of instability.

Instability arises from either soft tissue and/or bone damage. Capsulolabral complex is usually detached and stretched when a trauma occurs, and this lesion can be extended to the inferior labrum and in worst cases involve the whole labrum in a 360° detachment from the glenoid. Anterior glenoid rim may be involved. On the other side, posterior aspect of the humeral head is usually damaged because of the impact with the anteroinferior glenoid (Hill-Sachs lesion). This lesion can be very medial and deep on the cartilage surface, thus engaging with the anterior glenoid rim and causing dislocation. Di Giacomo et al. [8] introduced the new concept of “on-track/off-track lesion” to establish an algorithm for the treatment of the anterior instability.

The size of anteroinferior glenoid bone loss has been widely investigated to establish a treatment algorithm and any further recurrence [9]. Addressing glenoid bone loss must take into account the type of lesion. In case of glenoid fracture with a small fragment, especially in recurrent instability, fragment can undergo resorption, and bone loss is typically larger than the bone fragment as it appears on computed tomography (CT) scan [10].

V. Bongiorno
Polyclinique Notre Dame, Draguignan, France

2.3 Causes of Failure

A new high-energy trauma is a possible cause of failure if it occurs at least after 9–12 months from surgery and after total recovery and return to sport/work. Unfortunately, many times the reason of failure is related to errors in surgical technique (Fig. 2.1) or indication.

Indeed, the number of anchors, number of sutures, and position of the lower anchor are all crucial factors determining the outcome of the repair [11]. However, wrong indication is probably the most common cause of failure. According to Boileau et al. [11], factors like age at surgery (less than 20 years), type (overhead) and level (competitive) of sport participation, hyperlaxity, and bone loss either at the level of anteroinferior glenoid rim and the posteromedial humeral head are all concurrent to increased failure rate. Porcellini et al. [12] also described a higher risk of recurrence following arthroscopic Bankart repair on the basis of sex, age, and time from the first dislocation to surgery.

Randelli and Taverna [13] published a paper analyzing a young athlete after first dislocation and observed that patients with high-demanding shoulder activities had higher risk of failure, thus suggesting early surgical stabilization. Aboalata et al. [14] also described better clinical and radiological results in patients stabilized after first dislocation, and Crall et al. [15] evaluated the



Fig. 2.1 Right shoulder. Arthroscopic view from antero-superior portal. Surgical error: previous anchors were not positioned on the glenoid rim

cost-effectiveness of this strategy showing better overall results with early surgery.

2.4 Evaluation

The first thing to analyze in case of failure of anterior stabilization is preoperative clinical history and imaging. In addition, surgical report and postoperative rehabilitation protocol should be analyzed. In case of failure of arthroscopic Bankart repair, accurate assessment of soft tissues and bone loss, either on the glenoid and on the humeral head, is highly recommended. Magnetic resonance arthrography (MRA) and CT scan are the most accurate imaging studies (Fig. 2.2).

2.5 Treatment Algorithm

After a Bankart repair, failure can be diagnosed when recurrence of dislocation does occur. When regained stability is enough for a sedentary lifestyle, the patient cannot return to full sport activities but does not always complain of persistent



Fig. 2.2 Right shoulder. 2D CT scan. Anchor insertion site is usually the weakest point. Glenoid bone loss is calculated

instability. This explains some good results reported in short-term follow-up studies of untreated failed arthroscopic Bankart repair. Most of the times, patients are limited by a permanent minor instability, which can be painful and evolve toward more symptomatic instability and/or osteoarthritis (OA) over time.

Revision by new soft tissue repair to the glenoid has limited indications as a first solution for Bankart failure since soft tissue quality decreases over time so the hammock effect would not be reproduced properly by the weak ligament (Fig. 2.3). Nevertheless, in case of failure, revision by arthroscopic soft tissue repair should be always considered. If no glenoid erosion nor fractures are evident and, during arthroscopic assessment, the capsule-ligamentous structures show good appearance, thickness, and resistance under tension, a new soft tissue repair can be considered. The repair should be done with at least three double-loaded anchors by shifting the capsule and ligaments from posterior to anterior and from inferior to superior.

Humeral head Hill-Sachs “remplissage” has to be taken into account when the bone loss is deep and medial because it can be engaging and determine failure of treatment and early OA in a stabilized shoulder [16].

Coracoid transfer as described by Latarjet-Patte or Bristow, also performed arthroscopically as described by Lafosse et al. [17] and Boileau et al. [18], is a reliable solution in case of failed anterior arthroscopic repair with severe glenoid

bone loss, insufficient ligaments, and/or large Hill-Sachs lesion. Particularly, in case of failure of previous surgery, the sling effect of the conjoint tendon is necessary to achieve the needed stability for patients demanding high performances (sport or work).

A simple bone graft would not be enough without proper ligament complex. This is a necessary condition when considering only isolated bone graft glenoid augmentation. Conversely, an isolated bone graft with iliac crest (open or arthroscopic) can be the only solution in case of failure of anterior stabilization with Latarjet or Bristow procedure [19, 20].

References

1. Castagna A, Garofalo R, Conti M, Flanagan B. Arthroscopic Bankart repair: have we finally reached a gold standard? *Knee Surg Sports Traumatol Arthrosc.* 2016;24(2):398–405.
2. Browe DP, Rainis CA, McMahon PJ, Debski RE. Injury to the anteroinferior glenohumeral capsule during anterior dislocation. *Clin Biomech (Bristol, Avon).* 2013;28(2):140–5.
3. Virk MS, Manzo RL, Cote M, Ware JK, Mazzocca AD, Nissen CW, Shea KP, Arciero RA. Comparison of time to recurrence of instability after open and arthroscopic Bankart repair techniques. *Orthop J Sports Med.* 2016;4(6):2325967116654114.
4. Wang L, Liu Y, Su X, Liu S. A meta-analysis of arthroscopic versus open repair for treatment of Bankart lesions in the shoulder. *Med Sci Monit.* 2015;21:3028–35.
5. Middleton RM, Vo A, Ferguson J, Judge A, Alvand A, Price AJ, Rees JL. Can surgical trainees achieve arthroscopic competence at the end of training programs? A cross-sectional study highlighting the impact of working time directives. *Arthroscopy.* 2017;33(6):1151–8.
6. Ho AG, Gowda AL, Michael Wiater J. Evaluation and treatment of failed shoulder instability procedures. *J Orthop Traumatol.* 2016;17(3):187–97.
7. Walch G, Boileau P, Levigne C, Mandrino A, Neyret P, Donell S. Arthroscopic stabilization for recurrent anterior shoulder dislocation: results of 59 cases. *Arthroscopy.* 1995;11(2):173–9.
8. Di Giacomo G, Itoi E, Burkhart SS. Evolving concept of bipolar bone loss and the hill-Sachs lesion: from “engaging/non-engaging” lesion to “on-track/off-track” lesion. *Arthroscopy.* 2014;30(1):90–8.
9. Baudi P, Righi P, Bolognesi D, Rivetta S, Rossi Urtoler E, Guicciardi N, Carrara M. How to identify and calculate glenoid bone deficit. *Chir Organi Mov.* 2005;90(2):145–52.




Fig. 2.3 Right shoulder. Arthroscopic view from posterior portal: soft tissue insufficiency