

Nitish Kumar *Editor*

Biotechnological Approaches for Medicinal and Aromatic Plants

Conservation, Genetic Improvement and
Utilization

 Springer

Biotechnological Approaches for Medicinal and Aromatic Plants

Nitish Kumar
Editor

Biotechnological Approaches for Medicinal and Aromatic Plants

Conservation, Genetic Improvement
and Utilization

 Springer

Editor
Nitish Kumar
Department of Biotechnology
Central University of South Bihar
Panchanpur, Gaya, Bihar, India

ISBN 978-981-13-0534-4 ISBN 978-981-13-0535-1 (eBook)
<https://doi.org/10.1007/978-981-13-0535-1>

Library of Congress Control Number: 2018953756

© Springer Nature Singapore Pte Ltd. 2018

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Singapore Pte Ltd.
The registered company address is: 152 Beach Road, #21-01/04 Gateway East, Singapore 189721, Singapore

Preface

Medicinal and aromatic plants are the most important source of life-saving drugs for the majority of the world's population. The biotechnological tools are important to select, multiply and conserve the critical genotypes of medicinal plants. In vitro regeneration holds tremendous potential for the production of high-quality plant-based medicine. Plant tissue culture and traditional methods provide an opportunity for conservation of endangered medicinal and aromatic plants. In vitro production of secondary metabolites in plant cell suspension cultures has been reported from various medicinal plants. Genetic transformation may be a powerful tool for enhancing the productivity of novel secondary metabolites, especially by *Agrobacterium rhizogenes*-induced hairy roots.

Biotechnological Approaches for Medicinal and Aromatic Plants-Conservation, Genetic Improvement and Utilization discusses the applications of plant biotechnology for enhancement of secondary metabolite production in vitro from medicinal and aromatic plants. This book contains 29 chapters divided into 3 parts.

Part I: In vitro production of secondary metabolite

Part II: In vitro propagation, genetic transformation and germplasm conservation

Part III: Conventional and molecular approach

Gaya, Bihar, India

Nitish Kumar

Contents

Part I In Vitro Production of Secondary Metabolite

- 1 Production of Plant Secondary Metabolites: Current Status and Future Prospects** 3
P. Silpa, K. Roopa, and T. Dennis Thomas
- 2 The Effects of *rol* Genes of *Agrobacterium rhizogenes* on Morphogenesis and Secondary Metabolite Accumulation in Medicinal Plants** 27
Sayantika Sarkar, Ipshta Ghosh, Dipasree Roychowdhury, and Sumita Jha
- 3 Conventional and Biotechnological Approaches to Enhance Steviol Glycosides (SGs) in *Stevia rebaudiana* Bertoni** 53
Arpan Modi and Nitish Kumar
- 4 Effect of Chemical Elicitors on Pentacyclic Triterpenoid Production in In Vitro Cultures of *Achyranthes aspera* L.** 63
L. Sailo, Vinayak Upadhyya, Poornananda M. Naik, Neetin Desai, Sandeep R. Pai, and Jameel M. Al-Khayri
- 5 The Current Status and Future Applications of Hairy Root Cultures.** 87
Nisha Dhiman, Vanita Patial, and Amita Bhattacharya
- 6 In Vitro Culture and Production of Secondary Metabolites in *Centella asiatica*** 157
Shweta Kumari, Shashikant, Nitish Kumar, and Maheshwar Prasad Trivedi
- 7 Characterization of a Secondary Metabolite from *Aegle marmelos* (Vilva Tree) of Western Ghats.** 175
Vellingiri Manon Mani and Arockiam Jeyasundar Parimala Gnana Soundari

8	Role of Secondary Metabolites for the Mitigation of Cadmium Toxicity in Sorghum Grown Under Mycorrhizal Inoculated Hazardous Waste Site	199
	Prasann Kumar, Shweta Pathak, Mukul Kumar, and Padmanabh Dwivedi	
9	In Vitro Production of Some Important Secondary Metabolites from <i>Zingiber</i> Species	213
	Sanatombi Rajkumari and K. Sanatombi	
10	Hairy Root Culture for In Vitro Production of Secondary Metabolites: A Promising Biotechnological Approach	235
	Ravi Shankar Singh, Tirthartha Chattopadhyay, Dharamsheela Thakur, Nitish Kumar, Tribhuvan Kumar, and Prabhash Kumar Singh	
11	<i>Ocimum gratissimum</i>: A Review on Ethnomedicinal Properties, Phytochemical Constituents, and Pharmacological Profile	251
	Chaudhary Priyanka, Sharma Shivika, and Sharma Vikas	
12	Phytochemical Analysis with Special Reference to Leaf Saponins in <i>Gnidia glauca</i> (Fresen.) Gilg.	271
	Torankumar Sannabommaji, Vadlapudi Kumar, D. V. Poornima, Hari Gajula, J. Rajashekar, T. Manjunatha, and Giridhara Basappa	
13	In Vitro Production of Bacosides from <i>Bacopa monnieri</i>	289
	Praveen Nagella, Poornananda M. Naik, and Jameel M. Al-Khayri	
14	Production of the Anticancer Compound Camptothecin in Root and Hairy Root Cultures of <i>Ophiorrhiza mungos</i> L.	303
	B. Wetterauer, E. Wildi, and M. Wink	
Part II In Vitro Propagation, Genetic Transformation and Germplasm Conservation		
15	In Vitro Approaches for Conservation and Sustainable Utilization of <i>Butea monosperma</i> (Lam.) Taub. Var. Lutea (Witt.) Maheshwari: A Highly Valuable Medicinal Plant	345
	Rajesh Yarra, Ramesh Mushke, and Madhu Velmala	
16	Cryopreservation of Medicinal Herbs: Major Breakthroughs, Hurdles and Future.	353
	Suprabuddha Kundu, Umme Salma, and Saikat Gantait	
17	Ethnobotany: A Bridge Between Traditional Knowledge and Biotechnological Studies on Medicinal and Aromatic Plants	383
	A. S. Vishwanathan	

18	In Vitro Manipulations for Value Addition in Potent Herbal Insecticidal Activities of <i>Chrysanthemum cinerariaefolium</i>	395
	Shamshad A. Khan, Priyanka Verma, Varsha A. Parasharami, and Laiq Ur Rahman	
19	Biotechnological Approaches for Genetic Improvement of Fenugreek (<i>Trigonella foenum-graceum</i> L.)	417
	M. Aasim, F. S. Baloch, A. Bakhsh, M. Sameeullah, and K. M. Khawar	
20	Biotechnological Advancement in an Important Medicinal Plant, <i>Withania coagulans</i>: An Overview and Recent Updates	445
	Mangal S. Rathore, Kusum Khatri, Jasminkumar Kheni, and Narpat S. Shekhawat	
21	Tissue Culture in Mulberry (<i>Morus</i> spp.) Intending Genetic Improvement, Micropropagation and Secondary Metabolite Production: A Review on Current Status and Future Prospects	467
	Tanmoy Sarkar, Thallapally Mogili, S. Gandhi Doss, and Vankadara Sivaprasad	
22	In Vitro Conservation Strategies for <i>Gloriosa superba</i> L.: An Endangered Medicinal Plant	489
	Ritu Mahajan, Pallavi Billowaria, and Nisha Kapoor	
23	Somaclonal Variations and Their Applications in Medicinal Plant Improvement.	503
	Frédéric Ngezahayo	
Part III Conventional and Molecular Approach		
24	Genetic Improvement of Medicinal and Aromatic Plants Through Haploid and Double Haploid Development	523
	Sweta Sharma, Kshitij Vasant Satardekar, and Siddhivinayak S. Barve	
25	Role of Molecular Marker in the Genetic Improvement of the Medicinal and Aromatic Plants.	557
	Anubha Sharma, Nitish Kumar, and Iti Gontia Mishra	
26	Genetic Engineering Potential of Hairy Roots of Poppy (<i>Papaver</i> spp.) for Production of Secondary Metabolites, Phytochemistry, and In Silico Approaches	569
	Mala Trivedi, Aditi Singh, Parul Johri, Rachana Singh, and Rajesh K. Tiwari	
27	Advances in Genetic Engineering of <i>Ajuga</i> Species	599
	Waqas Khan Kayani, Humna Hasan, and Bushra Mirza	

- 28 Laurel (*Laurus nobilis* L.): A Less-Known Medicinal Plant to the World with Diffusion, Genomics, Phenomics, and Metabolomics for Genetic Improvement. 631**
Muhammad Azhar Nadeem, Muhammad Aasim, Saliha Kırıcı, Ünal Karık, Muhammad Amjad Nawaz, Abdurrahim Yılmaz, Hasan Maral, Khalid Mahmood Khawar, and Faheem Shehzad Baloch
- 29 Biological Databases for Medicinal Plant Research 655**
Sonu Kumar and Asheesh Shanker

About the Editor

Nitish Kumar has over 8 years of teaching and research experience in Plant Biotechnology and currently working as an Assistant Professor at Department of Biotechnology, Central University of South Bihar, Gaya, Bihar, India. He received his PhD degree in Botany from Council for Scientific and Industrial Research (CSIR)-Central Salt & Marine Chemicals Research Institute (CSMCRI)/Bhavnagar University, Bhavnagar, Gujarat, India. He has been the recipient of the Outstanding Faculty in Biological Sciences Award in the year of 2017 by the Venus International Foundation, Chennai, Tamil Nadu, India. He has received many awards/fellowships during his academic career from different organizations like CSIR, Department of Biotechnology (DBT), Indian Council of Agricultural Research (ICAR), and Department of Science & Technology (DST). He also got the Fast Track Young Scientist Award from DST, Government of India, in 2013. Dr. Kumar has published more than 40 research papers in peer-reviewed journals of national and international repute. He holds membership of several academic bodies/societies. He is also an Associate Editor of the journal *Gene*.

Part I
In Vitro Production of Secondary
Metabolite

Chapter 1

Production of Plant Secondary Metabolites: Current Status and Future Prospects



P. Silpa, K. Roopa, and T. Dennis Thomas

Abstract Plants are the prime life-supporting system on earth. Despite its use as food, it is also utilized as a source of life-saving drugs for majority of the population in the world. Many plants yield phytochemicals known as secondary metabolites, which are pharmaceutically important and are extracted directly from the plants collected from natural habitat. Regardless of conventional methods, biotechnological approaches especially plant tissue culture techniques play a unique role in producing and extracting secondary metabolites at industrial level. This book chapter discusses the various strategies adopted for secondary metabolite production in plants.

Keywords Secondary metabolites · Tissue culture · Metabolic engineering · Abiotic stress

1.1 Introduction

Plants have been the go-to resource for most of man's needs from aboriginal times. Recently, there is an increased interest in medicinal plants due to the rise in the use of herbal medicine and its therapeutic effects. Plants being a great source of bioactive secondary metabolites play a vital role in the field of drug development (Jose and Thomas 2014). Thus, the alternation of these active components through tissue culture and other biotechnical methods acts as a pillar in drug research. However, the requirement of high quantity of raw material is the main hurdle in using plant material as a key resource in drug development. Modifying the genetic makeup through metabolic engineering and high biomass production through tissue culture of these plants would lead to quality and quantity efficient production of bioactive compounds required for drug research.

P. Silpa · K. Roopa · T. Dennis Thomas (✉)
Department of Plant Science, Riverside Transit Campus, Central University of Kerala,
Kasaragod, Kerala, India

1.1.1 Plant Secondary Metabolites

Plants synthesize a variety of organic compounds, mainly classified as primary and secondary metabolites. Primary metabolites are required for essential biochemical processes such as growth, development, photosynthesis and respiration. Secondary metabolites are mainly involved in defence and protect against environmental stresses and also give specific colour, odour and tastes to the plants. Bioactive compounds are also used in agriculture field to protect crops from pests and also in signal transduction to attract seed dispersal and pollination (Wink 2003). Plant secondary metabolites have no key role in maintenance of the life processes in plants. It forms a significant source for pharmaceuticals, insecticides, flavouring agents, drugs (morphine, codeine, cocaine, quinine, etc.) and many other important biochemicals. In addition to this, certain plant secondary metabolites like phenolics, terpenoids, flavonoids and sulphur and nitrogen derivatives play a critical role to prevent many human diseases (Leicach and Chludil 2014). Biotic and abiotic stresses have a pivotal role in accumulation of secondary metabolites in various plant species (Pavarini et al. 2012). Plant secondary metabolites have been divided into three classes, namely, terpenes, phenolic and nitrogen-containing compounds (Taiz and Zeiger 2004).

1.1.2 Terpenes

Terpenes are one of the largest classes of secondary metabolites, and it is built up from isoprene units. Terpene is derived from the word “turpentine”. Terpenes are further divided into monoterpenes (e.g. carvone, perillyl alcohol, geraniol, limonene), sesquiterpene, diterpene (retinoic acid and retinol), triterpene (lupeol, betulinic acid, olealonic acid) and tetraterpene (lycopene, β -carotene, α -carotene, lutein) based on the number of isoprene units (Thoppil and Bishayee 2011). In higher plants, terpene biosynthesis occurs through two pathways, namely, mevalonic acid pathway and MEP pathway. Minor changes in the enzyme terpene synthases can trigger new catalytic properties easily and thereby induce terpene production (Keeling et al. 2008). Terpenes provide a number of esteemed functions, to attract pollinators, signalling compounds in metabolic pathway, plant-pathogen interactions, environmental stresses and other plant defences (Chen et al. 2011). Certain reports have shown that an endophytic fungus *Hypericum perforatum* helps in the production of terpenes in some plants (Zwenger 2008).

1.1.3 Phenolic Compounds

Tannins, flavonoids, stilbenes, lignans, quinines, phenolic acids, coumarins, etc. are the major phenolic compounds from medicinal herbs. Phenolic compounds form an aromatic ring and contain one or more hydroxyl groups. It possesses various bioactivities like anti-inflammatory, antioxidant, anti-carcinogenic and anti-mutagenic effects (Huang et al. 2010). Phenolics are responsible for the colour of red fruits, wines, and juices and also act as flavouring agents (Cheynier 2012). It can be divided into simple and complex phenolic compounds. These biologically active compounds mainly involved in plant defences during stressed conditions. Flavones and flavanols, isoflavonoids, tannins, anthocyanins, and lignin are included in the complex phenolic compounds. These compounds have significant role in many physiological events like flower and root differentiation, characterization of developmental stages, gene activity determination and growth vigour (Akillioglu 1994).

1.1.4 Nitrogen-Containing Compounds

Alkaloids are heterogenous group of secondary metabolites, and in an estimate, about 12,000 alkaloids are isolated from various plants (Ziegler and Facchini 2008). The three major alkaloids, cyanogenic glycosides and glucosinolates and nonprotein amino acids, are the three groups of nitrogen-containing compounds. They mainly act as growth regulators, provide protection from predators and also maintain ionic balance. Cyanogenic glycosides and glucosinolates send out volatile poisons and also play a key role in defence mechanisms (Taiz and Zeiger 2004).

1.2 Production of Plant Secondary Metabolites

Plant secondary metabolites are antibiotic, antifungal or antiviral agents that have the primary function of protecting the plants from disease-causing organisms or pathogens. Plants are mainly exploited for pharmaceuticals, food colours, flavours, fragrances and sweeteners. In this high-tech era, man mainly depends on plant bioactive components for the development of drugs and plant-derived drugs, and intermediates constitute about 25% of the total prescription drugs. Biotechnological approaches such as plant tissue culture techniques have great potential as an alternative for production of useful medicinal compounds like alkaloids, terpenoids, steroids, saponins, phenolics, flavonoids and amino acids. Some commercially available secondary metabolites which are available in the market include shikonin and taxol. In natural conditions, uniform availability of secondary metabolites is not possible due to several reasons. However, cultivation of plant cells or tissues in

aseptic conditions in a bioreactor often leads to consistent production of secondary metabolites with improved quality and yield (Fowler 1985).

For improving the production of secondary metabolites, a number of strategies like screening and selection of high-yielding cell lines; culture of cells from various organs such as shoots, roots, leaves, callus, etc.; suspension culture; induction by elicitors; metabolic engineering; and optimization of media and plant growth regulators were adopted (Anand 2010).

Plant cell and tissue culture holds great potential for controlled production of numerous secondary metabolites which could be useful for various purposes. Secondary metabolite production under controlled condition is devoid of any environmental fluctuations and is cost-effective (Rao and Ravishankar 2002). Plant cell and tissue culture helps to produce important bioactive compounds, and advances in this area may enhance the production of these compounds. The secondary metabolites produced by plant tissue and organ culture are similar to secondary metabolites produced by intact plants. Commercially produced shikonin and taxol are now available in markets. About 20 recombinant proteins like enzymes, antibodies, growth factors and edible vaccines have been produced from tissue culture techniques. Large-scale production of secondary metabolites can be achieved through cell suspension culture by transferring friable callus to suitable medium. The advantage of using cell suspension culture for secondary metabolite production over field-grown plant is that it is devoid of production interfering compounds (Filova 2014).

Production of secondary metabolites in higher plants can be achieved through different explants under sterile condition. Plant cell and tissue culture technique is routinely employed to extract secondary metabolites from various plant species (Table 1.1). Many secondary metabolites are synthesized from primary metabolites in higher plants. About 100,000 secondary metabolites have been isolated from higher plants (Jeong and Park 2006). Its production is usually in lesser quantities, and it is determined by developmental stage and physiology of plants. Medium optimization is required to enhance the production. In many cases, suspension culture, organ culture, embryo culture and callus culture have been successful in increasing the yield of secondary metabolites. The interested metabolites may sometimes synthesize in plant from specialized tissue or organs. For example, saponin is produced from the root of *Panax ginseng*, and hence for large scale, in vitro production of saponin from *P. ginseng* requires root culture. At present several medicinal plants have been utilized to establish various culture systems like callus culture, organ culture, hairy root culture and suspension culture. There are many biotechnological approaches to produce bioactive components and are briefly described below.

Table 1.1 Recent studies on secondary metabolites production by using plant cell/tissue cultures

Plant	Compound/s	References
<i>Anoectochilus roxburghii</i>	Kinsenoside	Jin et al. (2017)
<i>Withania somnifera</i>	Withanolides	Ahlawat and Abdin (2017)
<i>Stevia rebaudiana</i>	Antioxidants	Ahmad et al. (2016)
<i>Vitis</i> species	Monoterpene and sesquiterpene	Alonso et al. (2015)
<i>Dysoxylum binectariferum</i>	Rohitukine	Mahajan et al. (2015)
<i>Prunella vulgaris</i>	Antioxidants	Fazal et al. (2014)
<i>Withania somnifera</i>	Withanolide	Sabir et al. (2013)
<i>Vitis</i> species	Nerolidol	Escoriza et al. (2013)
<i>Solanum tuberosum</i>	Chlorogenic acid (phenolics)	Navarre et al. (2013)
<i>Humulus lupulus</i>	Prenylflavonoid	Matousek et al. (2012)
<i>Ruta graveolens</i>	Umbelliferone	Vialart et al. (2012)
<i>Nicotiana tabacum</i>	Anthocyanins	Huang et al. (2012)
<i>Senecio</i> species	Pyrrrolizidine alkaloid	Karam et al. (2011)
<i>Medicago sativa</i>	Saponins	Szakiel et al. (2011)
<i>Brachiaria</i> species	Protodioscin	Barbosa-Ferreria et al. (2011)
<i>Lavandula officinalis</i>	Tetrahydrofurate (THF) acetate derivative	Patel et al. (2011)
<i>Malus</i> species	Anthocyanins	Lin-Wang et al. (2011)
<i>Eugenia uniflora</i>	Tannins and flavonoids	Santos et al. (2011)
<i>Crotalaria retusa</i>	Monocrotaline and pyrrolizidine	Anjos et al. (2010)
<i>Echium plantagineum</i>	Pyrrrolizidine (alkaloid)	Lucena et al. (2010)
<i>Vernonia</i> species	Lactones and flavonoids	Keles et al. (2010)
<i>Lychnophora</i> species	Lactones and flavonoids	Gobbo-Neto et al. (2010)
<i>Angelica gigas</i>	Decursin, decursinol angelate	Rhee et al. (2010)
<i>Lavandula officinalis</i>	Deoxyartemisinin	Patel et al. (2010)
<i>Lavandula pedunculata</i>	Essential oils	Zuzarte et al. (2010)
<i>Lavandula vera</i>	Volatiles	Georgiev et al. (2010)
<i>Lithospermum erythrorhizon</i>	Shikonin	Zhang et al. (2010)
<i>Argemone mexicana</i>	Sanguinarine	Trujillo – Villanueva et al. (2010)
<i>Picrorhiza kurroa</i>	Picoside –I	Sood and Chauhan (2010)
<i>Thevetia peruviana</i>	Peruvoside	Zabala et al. (2010)
<i>Abrus precatorius</i>	Glycyrrhizin	Karwasara et al. (2010)
<i>Silybum marianum</i>	Silymarin	Khalili et al. (2010)
<i>Artemisia</i> species	Artemisin	Brown (2010)
<i>Lavandula pedunculata</i>	Camphor and 1,8- cineole	Zuzarte et al. (2010)

1.2.1 *Callus Culture*

Callus is the mass of undifferentiated cells containing meristematic loci (Bhojwani and Dantu 2013). For induction of calli from explants, 2, 4-D is the most preferred auxin. However, a combination of auxin and cytokinin or high concentration of auxin alone may be used by various workers for callus induction (Filova 2014). For secondary metabolite production, usually non-embryogenic calli were selected which have homogenous mass of dedifferentiated cells. For callus growth and multiplication, auxin is generally preferred, whereas it is omitted for secondary metabolite production. Screening and selection of high-yielding cell lines and standardization of media for optimum secondary metabolite production are some important strategies to improve secondary metabolite production. Suspension culture is an alternative strategy to obtain high level of secondary metabolite production. Small clumps of calli are transferred to liquid medium in flasks, and this will be followed by continuous agitation on an orbital shaker. Agitation often exerts a pressure on the cell clumps resulting in the breaking of larger clumps into smaller aggregates. Agitation also helps in the uniform distribution of cells and better aeration of cells inside medium. High rate of cell division can be achieved in suspension culture than normal callus culture.

1.2.2 *Hairy Root Culture*

Plant hairy root culture is the most promising technique among root culture for the production of secondary metabolites. Fast hormone-independent growth, genetic stability, lateral branching and lack of geotropism are the major characteristics of hairy roots. Inoculation of *Agrobacterium rhizogenes* helps to synthesize secondary metabolites in hairy roots (Karuppusamy 2009; Palazon et al. 1997).

A. rhizogenes have root-inducing plasmid (Ri plasmid) which contain a T-DNA; during the time of infection, T-DNA is further divided into TL and TR region in some strains (strain A4) of *A. rhizogenes* in induction processes. Two sets of PRi genes, *aux genes* (in TR region) and *rol genes* (in TL region) are involved (Jouanin 1984). Elicitation of hairy root promotes secondary metabolite production and also arrests feedback inhibition, preventing degradation of metabolites in the culture medium (Chandra and Chandra 2011).

In many plants, hairy root become green by culturing in continuous exposure of light. It generates photo-oxidative stress in hairy roots (Mukherjee et al. 2014). It produces excess H₂O₂ in the root (Behnke et al. 2010). Phenolics to volatile terpenoids shift occur in green hairy roots of carrot, due to the redirection of primary metabolites towards synthesis of volatile isoprenoid synthesis (Mukherjee et al. 2016).

1.2.3 Organ Culture

Organ culture techniques help the rapid propagation of plants. It also gets a high quantity of bioactive compounds and higher growth compared to plants grown in natural habitats. In *Fritillaria unibracteata*, rapid propagation is achieved through small bulb, and the content of alkaloids is higher in this bulb culture (Gao et al. 1999). Tropane alkaloids hyoscyamine and scopolamine were produced in high quantity in root culture (Fazilatun et al. 2004). Many valuable medicinal compounds were obtained from root culture (Pence 2011; Li et al. 2002). Secondary metabolites produced in plant aerial parts are produced from root culture (Bourgaud et al. 2001; Nogueira and Romano 2002; Smith et al. 2002; Kaimoyo et al. 2008). Organ culture exhibits less sensitivity of shear stress, but in biomass production, they show a high degree of spatial heterogeneity. Ginseng roots are the only example for commercially production of secondary metabolites by organ culture (Hibino and Ushiyama 1999).

1.2.4 Elicitation

Elicitors are substances which produce signals against pathogenic attack resulting in the accumulation of secondary metabolites in plants. Further it can improve the biosynthesis of specific compounds when introduced into a living cell system (Radman et al. 2003). Elicitation is the process in which the living cells will be treated with biotic or abiotic elicitors to obtain enhanced rate of secondary metabolites (Rao and Ravishankar 2002). The most commonly employed biotic elicitors include polysaccharides, glycoproteins, yeast extract and some fungi like *Fusarium oxysporum*, *Aspergillus niger* and *Rhizopus oryzae* (Dornenburg and Knorr 1995). The abiotic elicitors are non-biological origin and are mostly inorganic salts, jasmonic acid, salicylic acid, high pH and environmental stress conditions such as heavy metals, UV radiation, osmotic shock, etc. (Naik and Al-Khayri 2015). High stilbenes accumulation in root cultures of *Cayratia trifolia* was observed by Arora et al. (2009). The addition of alar (N-dimethylamino succinamic acid) along with the elicitor salicylic acid enhanced the stilbenes content up to 12-folds (Arora et al. 2009). In *Centella asiatica*, the presence of triterpenes in callus suspension culture derived from leaves showed an increase after incorporating amino acids (Kim et al. 2004). Adding the amino acid isoleucine at 2 mM in the medium enhanced the production of hyperforin in *Hypericum perforatum* (Karppinen et al. 2007). Hence elicitation is considered as one of the most effective methods for enhancing the secondary metabolite production in cultures (Oksman-Caldentey and Inze 2004). Heavy metals and increasing temperature affect the production of secondary metabolites. In *Robinia pseudoacacia* seedlings exposed to elevated carbon dioxide, high temperature and heavy metals (Pb-Cd) enhanced the production of secondary metabolites (Zhao et al. 2016).

1.2.5 Endophytes

Microbes, such as bacteria or fungus which lives inside the plant without making any indication of disease, are called endophytes. Many useful compounds were obtained from the plants with endosymbionts. For example, *taxol*, an anticancer agent, is obtained from *Taxus brevifolia* when infected with a fungi *Taxomyces andreanae* (Strobel et al. 1993). Accumulation of several valuable metabolites in plants is due to the synergistic effect of both plants and endophytes (Engels et al. 2008). The endophytic fungi of *Pinus sylvestris* and *Rhododendron tomentosum* produce useful secondary metabolites having various antibacterial and antioxidant activities (Kajula et al. 2010). It is reported that some endophytic fungi produce a number of beneficial phytochemicals in Leguminosae family (Wink 2013). In some plants, due to the presence of endophytic fungi, an alkaloid called indolizidine is produced (Ralphs et al. 2008). The endophytic fungus *Phoma medicaginis* produces a compound, hydroxy-6-methylbenzoic acid, which shows a noticeable antimicrobial activity (Yang et al. 1994). Phenylpropanoids, lignins, phenol and phenolic compounds, alkaloids, steroids, etc. are isolated from many mycoendophytes (Herre et al. 2007). Novel metabolites from the endophytes have a vital role in the treatment of many infectious diseases (Rai et al. 2012). It is reported that several genes and protein present in plants were seen in fungi and bacteria. This suggests that horizontal gene transfer may take place from endosymbiotic bacteria and fungi (Wink and Schimmer 2010).

Inoculation of arbuscular mycorrhizal fungi (AMF) increases the production of secondary metabolites in economically important plants (Maier et al. 1995). The presence of AMF can considerably enhance the growth and biomass of a plant (Silva et al. 2004). It can also improve the capability to absorb the micronutrients and macronutrients (Chu et al. 2001; Matsubara et al. 2009). Recently, there is an increase in research about the efficiency of AMF in improving the secondary metabolite production (Table 1.2). There is a qualitative and quantitative increase in secondary metabolite production in many plants due to the presence of AMF (Ponce et al. 2004; Ceccarelli et al. 2010). In addition to these, there is an increasing edible vegetable quality triggered by AMF (Baslam et al. 2013).

1.2.6 Nitric Oxide

Nitric oxide plays a crucial role in the production of some important phytochemicals. By the induction of some stresses, accumulation of nitric oxide takes place. Several studies showed that nitric oxide plays a significant role in the development, growth and defence responses of the plant (Flores et al. 2008; Hong et al. 2008). Pharmaceutically important secondary metabolites can be produced by elicitor-induced nitric oxide response (Xu and Dong 2008). Therefore, the significance of nitric oxide can be applied in various biotechnological processes resulting in the production of target secondary metabolites.

Table 1.2 Recent report on influence of inoculation of arbuscular mycorrhizal fungi (AMF) on secondary metabolite production

Host plant	Plant organ	Evaluated phytochemicals	AMF	Effects	References
<i>Moringa oleifera</i>	Leaves	Glucosinolates	<i>Rhizophagus intraradices</i>	+ ve	Cosme et al. (2014)
			<i>Funneliformis mosseae</i>	+ ve	
<i>Helianthus annuus</i>	Seeds	Fixed oil	<i>Funneliformis mosseae</i>	+ve	Heidari and Karami (2014)
<i>Passiflora alata</i>	Leaves	Total phenols and total flavonoids	<i>Gigaspora albida</i>	+ ve	Oliveira et al. (2015)
<i>Stevia rebaudiana</i>	Leaves	Stevioside and rebaudioside A	<i>Rhizophagus fasciculatus</i>	+ ve	Mandal et al. (2013)
<i>Anadenanthera colubrina</i>	Leaves	Total phenols, total flavonoids and total tannins	<i>Acaulospora longula</i> + <i>Gigaspora albida</i>	+ ve	Pedone-Bonfim et al. (2013)
<i>Cucumis sativus</i>	Leaves	Phenols, flavonoids and lignin	<i>Funneliformis mosseae</i>	+ ve	Chen et al. (2013)

1.2.7 Abiotic Stress

Abiotic stresses are important for the production of secondary metabolites. Water stress is one of the prominent abiotic stresses which can influence secondary metabolite production. Light has a crucial role in the induction of both primary and secondary metabolites. Light-grown suspension culture displays an increase in phenolic production, in antioxidant activity and also in the total plant metabolite production (Ali and Abbasi 2014). It is reported that there is a link between antioxidant activity and total phenolic content in the suspension culture of *Artemisia absinthium* (Ali and Abbasi 2014). High blue light ratio increases all phenolic acids and flavonoids in some plants (Ouzounis et al. 2014).

1.2.8 Bioreactor

Bioreactors were designed for the commercial production of secondary metabolites. Bubble column bioreactor (Huang and McDonald 2009) and stirred tanks (Su 2006) are the widely used bioreactors for the culture of plant cell. Plants like *Eurycoma longifolia* grow well in bioreactor, and it has a rapid growth compared to that of flask cultures (Lulu et al. 2015). Exposure to UV light brings out secondary metabolite production in bioreactor and also in flask cultures. In *Lavandula vera* cells, rosmarinic acid production showed a 32-fold increase in bioreactors as compared to normal shake flask cultures (Pavlov et al. 2005). Production of secondary metabolites from cells of *Digitalis lantana*, *Catharanthus roseus*, *Hypericum perforatum*,

Panax ginseng, *Sophora flavescens*, etc. has been cultured in various bioreactors (Filova 2014). Sharma et al. (2011) studied the puerarin accumulation in *Pueraria tuberosa* during shoot cultures in static and liquid medium with or without aeration. Shoots were grown in growtek bioreactor with different aeration, and the maximum puerarin content was 1484 µg/g dry weight, which was about 2.3-fold higher than puerarin content recorded in control cultures (Sharma et al. 2011). The genes involving in withanolides production in *Withania somnifera* were upregulated in bioreactor. About 1.5-folds increase in the production of withanolides was found in bioreactor as compared to shake flask (Ahlawat and Abdin 2017). Different bioreactors like continuous immersion bioreactor with net (CIB-N), continuous immersion bioreactor (CIB), temporary immersion bioreactor (TIB) and temporary immersion bioreactor with net (TIB-N) were used for the production of bioactive compounds (Jang et al. 2016). Of these, CIB system was found to be the most efficient bioreactor for the large-scale production of metabolites (Jang et al. 2016). Bioreactor culture of *Anoectochilus roxburghii* accumulated highest level of kinsenoside and other polysaccharides (Jin et al. 2017).


1.2.9 Metabolic Engineering

Metabolic engineering has proven to be a valuable tool for large-scale production of several biomedically as well as industrially relevant secondary metabolites from plants. Current studies employing transgenic and/or recombinant technologies have opened up opportunities for metabolic engineering/metabolomics, leading to the manufacturing of high-value secondary metabolites, even at the commercial level. Metabolic engineering mainly aims at the increase in the production of desired products, brings down the production of unwanted compounds and scales up the yield of novel compounds (Ludwing-Muller et al. 2014). Many of the medicinally and economically important terpenoids are produced through metabolic engineering (Elgar 2017). Patel et al. (2016) reported that the incorporation of squalene synthase gene in isoprenoid pathway of *Withania somnifera* increased the production of bioactive compounds.

1.2.10 Immobilization

Immobilization enhances the secondary metabolite production by entrapping plant cells in suitable matrix, which can protect cells from liquid shear forces and allow better cell-to-cell contacts. The viability of immobilized cells extended over a prolonged period of time (Brodelius 1985). The benefit of immobilized plant cell is that it can extend its production time, making the cells catalyse the same reaction almost indefinitely. The immobilized cells can overcome the task of isolating the compounds from the biomass; rather the products will be delivered in the medium itself.

Fig. 1.1 Chemical structure of taxol



It can also perform multienzyme operations, and by using high-yielding cell lines, the productivity can be enhanced significantly (Smetanska 2008).

1.3 Production of Valuable Pharmaceutical Compounds Through In Vitro Culture Techniques

Most of the new therapeutics was evolved from secondary metabolites from plants. Advancement in the plant cell and tissue culture enhanced the production of several pharmaceutically active compounds, and some of such key compounds are described below.

1.3.1 Taxol

Taxol (Fig. 1.1) is a compound extracted from the bark of the Pacific yew tree (*Taxus brevifolia*) which possesses anticancer properties (Oksman-Caldentey and Inze 2004). In polymerized form of microtubules, taxol stabilizes it and thereby causes the death of cells. Due to the huge commercial use of taxol, *Taxus* species have been massively explored. In some studies, it was found that addition of certain amino acids like phenylalanine in the medium yielded maximum taxol in *T. cuspidata* (Long and Croteau 2005). The effect of both biotic and abiotic elicitors positively influenced the yield and accumulation of taxol in some species (Pavarini et al. 2012).

1.3.2 Morphine and Codeine

Morphine and codeine (Fig. 1.2a, b) are the pain-relieving drugs obtained from the members of the family Papaveraceae. Both these compounds occur naturally in Poppy plant (*Papaver somniferum*). Both morphine and codeine were commercially produced in cultures using callus and suspension culture (Yoshikawa and Furuya 1985). The optimum quantity of codeine and morphine was observed in cultures devoid of exogenous hormone (Furuya et al. 1972).

Fig. 1.2 Chemical structure of morphine (a) and codeine (b)





Fig. 1.3 Chemical structure of ginsenoside



1.3.3 Ginsenosides

The active component of the plant *Panax ginseng* is referred to as ginsenosides which are essential for various physiological activities. Chemically ginsenosides are a group of triterpene saponins (Fig. 1.3). The addition of spermidine in the medium enhanced the production of ginsenosides in cultures (Marsik et al. 2014). Further, an elicitor, casein hydrolyzate enhanced ginsenosides production without suppressing the biomass (Marsik et al. 2014). In root culture of *Panax ginseng*, jasmonic acid improves the ginsenosides production (Lambert et al. 2011).

1.3.4 L-DOPA

A nonprotein amino acid, L-DOPA (L-3, 4-dihydroxy phenylalanine; Fig. 1.4), is a potent drug obtained from the plant (*Mucuna hassjoo*) and is mainly used to cure Parkinson's disease (Brain and Lockwood 1976). It is the precursor of many secondary metabolites like alkaloids, melanin and betalain (Daxenbichler et al. 1971). The requirement of large quantities of L- DOPA led to the development of cell

Fig. 1.4 Chemical structure of L-DOPA




Fig. 1.5 Chemical structure of diosgenin





Fig. 1.6 Chemical structure of capsaicin



culture techniques to obtain optimum production of this compound. *Mucuna hass-joo* cells were cultured in MS medium supplemented with kinetin for obtaining optimum quantity of L-DOPA (Vanisree et al. 2004).

1.3.5 Diosgenin

Diosgenin (Fig. 1.5) is a biologically active metabolite, intermediate to various steroid drugs (Tal et al. 1984). Due to its high demand in the market, the production of diosgenin was enhanced by the application of in vitro techniques, and thereby it is beneficial to modern system of medicine. The optimum accumulation of diosgenin in culture was greatly influenced by the carbon and nitrogen level in the medium (Tal et al. 1984).

1.3.6 Capsaicin

The alkaloid, capsaicin (Fig. 1.6), is obtained from *Capsicum* species and mainly serves as a food additive (Ravishankar et al. 2003). The quantity of capsaicin differs significantly in suspension culture and immobilization technique. The amount of capsaicin was comparatively low in suspension culture, whereas it was increased substantially to about 100-fold in immobilization technique (Lindsey 1995). Further, the addition of isocaproic acid in the medium enhanced the production of capsaicin (Lindsey 1995).

Fig. 1.7 Chemical structure of berberine





Fig. 1.8 Chemical structure of camptothecin



1.3.7 Berberine

The isoquinoline alkaloid berberine (Fig. 1.7) is obtained from the cell culture of *Coptis japonica* (Vanisree et al. 2004), *Thalictrum* species (Nakagawa et al. 1986) and *Berberis* species (Morimoto et al. 1988). In order to increase the berberine yield in cultures, several elicitors were employed by Funk et al. (1987). Further, Nakagawa et al. (1984) and Morimoto et al. (1988) standardized the nutrient medium for optimum berberine production.

1.3.8 Camptothecin

Camptothecin (Fig. 1.8), a potent antitumor alkaloid, is isolated from the plant *Camptotheca acuminata* (Padmanabha et al. 2006). The production of camptothecin from *C. acuminata* cells in cultures was optimum on MS medium supplemented with 4.0 mg/l NAA (Thengane et al. 2003).

1.3.9 Vincristine and Vinblastine

The plant *Catharanthus roseus* (also known as *Vinca rosea*) contains the vinblastine and vincristine (Fig. 1.9a, b) which are used in chemotherapy (Noble 1990). Due to its irreplaceable medicinal properties, application of biotechnological tools




Fig. 1.9 Chemical structure of vinblastine (a) and vincristine (b)

especially plant tissue culture techniques was employed to produce large quantity of these compounds (Oksman-Caldentey and Inze 2004). Several chemicals such as oxalate, maleate, ferric chloride and sodium borohydride were added in the medium to increase the production of vinblastine (Verma et al. 2007). The other factors which influenced the production of these alkaloids include various stresses such as salinity, drought, heavy metals (Pandey 2017), UV stress (Binder et al. 2009) and presence of elicitors and addition of bioregulators (Zhao et al. 2001).

1.4 Conclusions

Plants are potent source of various useful phytochemicals. The secondary metabolites can be extracted from various parts of the plant. In recent years, there is an increased use of biotechnological tools to obtain continuous and reliable source of secondary metabolites. Among the various techniques, plant cell and tissue culture technology plays a crucial role in secondary metabolite production. The major advantage of using this technology is that it can provide bioactive secondary metabolites in controlled conditions irrespective of season and soil conditions. However, many challenges like difficulties in scaling-up, sustainability of culture and phytochemical recovery, etc. still remain to be challenging factors in this area. Continuous refinements of these techniques are necessary to overcome these limitations.

References

- Ahlatw, S., & Abdin, M. Z. (2017). Comparative study of withanolide production and the related transcriptional responses of biosynthetic genes in fungi elicited cell suspension culture of *Withania somnifera* in shake flask and bioreactor. *Plant Physiology and Biochemistry*, *114*, 19–28.
- Ahmad, N., Rab, A., & Ahmad, N. (2016). Light-induced biochemical variations in secondary metabolite production and antioxidant activity in callus cultures of *Stevia rebaudiana* (Bert). *Journal of Photochemistry and Photobiology. B*, *154*, 51–56.
- Akillioglu M (1994) *Zeytin agaclarinda dogel fenolik bilesiklerin mevsimsel degisimi Cizerinde arastirmalar* [Investigations on the seasonal changes of natural phenolic compounds in olive tree]. PhD thesis, Ege Univ. Fen Bilimleri Enst. Bahce Bitkileri Anabilim Dalı, Izmir.
- Ali, M., & Abbasi, B. H. (2014). Light-induced fluctuations in biomass accumulation, secondary metabolites production and antioxidant activity in cell suspension cultures of *Artemisia absinthium* L. *Journal of Photochemistry and Photobiology B: Biology*, *140*, 223–227.
- Alonso, R., Berli, F. J., Bottini, R., & Piccoli, P. (2015). Acclimation mechanisms elicited by sprayed abscisic acid, solar UV-B and water deficit in leaf tissues of field-grown grapevines. *Plant Physiology and Biochemistry*, *91*, 56–60.
- Anand. (2010). Various approaches for the secondary metabolite production through plant tissue culture. *Pharmacia*, *1*, 1–7.
- Anjos, B. L., Nobre, V. M., Dantas, A. F., Medeiros, R. M., Neto, T. S. O., Molyneux, R. J., & RietCorrea, F. (2010). Poisoning of sheep by seeds of *Crotalaria retusa*: Acquired resistance by continuous administration of low doses. *Toxicon*, *55*, 28–32.
- Arora, J., Roat, C., Goyal, S., & Ramawat, K. G. (2009). High stilbenes accumulation in root cultures of *Cayratia trifolia* (L.) Domin grown in shake flasks. *Acta Physiologiae Plantarum*, *31*, 1307–1312.
- Barbosa-Ferreria, M., Brum, K. B., Oliveira, N. M., do Valle, C. B., Ferreira, V. B., Garcez, V. S., Riet-Correa, F., & de Lemos, R. A. (2011). Steroidal saponin protodioscin concentration in different species and cultivars of *Brachiaria spp.* *Veterinaria e Zootecnia*, *18*, 98–103.
- Baslam, M., Esteban, R., García-Plazaola, J. I., & Goicoechea, N. (2013). Effectiveness of arbuscular mycorrhizal fungi (AMF) for inducing the accumulation of major carotenoids, chlorophylls and tocopherol in green and red leaf lettuces. *Applied Microbiology and Biotechnology*, *97*, 3119–3128.
- Behnke, K., Kaiser, A., Zimmer, I., Brüggemann, N., Janz, D., Polle, A., Hampp, R., Hahnisch, R., Popko, J., Schmitt-Kopplin, P., Ehrling, B., Rennenberg, H., Barta, C., Loreto, F., & Schnitzler, J. (2010). RNAi-mediated suppression of isoprene emission in poplar transiently impacts phenolic metabolism under high temperature and highlight intensities: A transcriptomic and metabolomic analysis. *Plant Molecular Biology*, *74*, 61–75.
- Bhojwani, S. S., & Dantu, P. K. (2013). Micropropagation. In *Plant tissue culture: An introductory text* (pp. 245–274). London: Springer.
- Binder, B. Y., Peebles, C. A., Shanks, J. V., & San, K. Y. (2009). The effects of UV-B stress on the production of terpenoid indole alkaloids in *Catharanthus roseus* hairy roots. *Biotechnology Progress*, *25*, 861–865.
- Bourgau, F., Gravot, A., Milesi, S., & Gontier, E. (2001). Production of plant secondary metabolites: A historical perspective. *Plant Science*, *161*, 839–851.
- Brain, K. R., & Lockwood, G. B. (1976). Hormonal control of steroid levels in tissue cultures from *Trigonella foenumgraecum*. *Phytochemistry*, *15*, 1651–1654.
- Brodellius, P. (1985). The potential role of immobilization in plant cell biotechnology. *Trends in Biotechnology*, *3*, 280–285.
- Brown, G. D. (2010). The biosynthesis of artemisinin (Qinghaosu) and the phytochemistry of *Artemisia annua* L.(Qinghao). *Molecules*, *15*, 7603–7698.

- Ceccarelli, N., Curadi, M., Martelloni, L., Sbrana, C., Picciarelli, P., & Giovannetti, M. (2010). Mycorrhizal colonization impacts on phenolic content and antioxidant properties of artichoke leaves and flower heads two years after field transplant. *Plant and Soil*, 335, 311–323.
- Chandra, S., & Chandra, R. (2011). Engineering secondary metabolite production in hairy roots. *Phytochemistry Reviews*, 10, 371–395.
- Chen, F., Tholl, D., Bohlmann, J., & Pichersky, E. (2011). The family of terpene synthase in plants: A mid-size family of genes for specialised metabolism that is highly diversified throughout the kingdom. *The Plant Journal*, 66, 212–229.
- Chen, S., Jin, W., Liu, A., Zhang, S., Liu, D., Wang, F., Lin, X., & He, C. (2013). Arbuscular mycorrhizal fungi (AMF) increase growth and secondary metabolism in cucumber subjected to low temperature stress. *Scientia Horticulturae*, 160, 222–229.
- Cheyrier, V. (2012). Phenolic compounds: From plants to foods. *Phytochemistry Reviews*, 11, 153–177.
- Chu, E. Y., Möller, M. D., & Carvalho, J. G. (2001). Effects of mycorrhizal inoculation on sour-sop seedlings in fumigated and not fumigated soil. *Pesquisa Agropecuária Brasileira*, 36, 671–680.
- Cosme, M., Franken, P., Mewis, I., Baldermann, S., & Wurst, S. (2014). Arbuscular mycorrhizal fungi affect glucosinolate and mineral element composition in leaves of *Moringa oleifera*. *Mycorrhiza*, 24, 565–570.
- Daxenbichler, M. E., VanEtten, C. H., Hallinan, E. A., Earle, F. R., & Barclay, A. S. (1971). Seeds as sources of L-DOPA. *Journal of Medicinal Chemistry*, 14, 463–465.
- Dörnenburg, H., & Knorr, D. (1995). Strategies for the improvement of secondary metabolite production in plant cell cultures. *Enzyme and Microbial Technology*, 17, 674–684.
- Elgar, S. M. (2017). *Metabolic engineering for the production of functionalized terpenoids in heterologous hosts*. Doctoral dissertation, Massachusetts Institute of Technology, USA.
- Engels, B., Dahm, P., & Jennewein, S. (2008). Metabolic engineering of taxadiene biosynthesis in yeast as a first step towards Taxol (Paclitaxel) production. *Metabolic Engineering*, 10, 201–206.
- Escoriza, G., Sansberro, P., Garcia-Lampasona, S., Gatica, M., Bottini, R., & Piccoli, P. (2013). In vitro cultures of *Vitis vinifera* L. cv. Chardonnay synthesize the phytoalexin nerolidol upon infection by *Phaeoacremonium parasiticum*. *Phytopathologia Mediterranea*, 52, 289–297.
- Fazal, H., Abbasi, B. H., & Ahmad, N. (2014). Optimization of adventitious root culture for production of biomass and secondary metabolites in *Prunella vulgaris* L. *Applied Biochemistry and Biotechnology*, 74, 2086–2095.
- Fazilatun, N., Normisah, M., & Zhari, I. (2004). Superoxide radical scavenging properties of extracts and flavonoids isolated from the leaves of *Blumea balsamifera*. *Pharmaceutical Biology*, 42, 404–408.
- Filová, A. (2014). Production of secondary metabolites in plant tissue cultures. *Research Journal of Agricultural Science*, 46, 236–245.
- Flores, T., Todd, C. D., Tovar-Mendez, A., Dhanoa, P. K., Correa-Aragunde, N., Hoyos, M. E., & Polacco, J. C. (2008). Arginase-negative mutants of *Arabidopsis* exhibit increased nitric oxide signaling in root development. *Plant Physiology*, 147, 1936–1946.
- Fowler, M. W. (1985). Problems in commercial exploitation of plant tissue cultures. In K. H. Neumann, W. Barz, & E. Reinhardt (Eds.), *Primary and secondary metabolism of plant cell cultures* (pp. 362–378). Berlin: Springer.
- Funk, C., Gügler, K., & Brodelius, P. (1987). Increased secondary product formation in plant cell suspension cultures after treatment with a yeast carbohydrate preparation (elicitor). *Phytochemistry*, 26, 401–405.
- Furuya, T., Ikuta, A., & Syono, K. (1972). Alkaloids from callus cultures of *Papaver somniferum*. *Phytochemistry*, 11, 3041–3044.
- Gao, S. L., Zhu, D. N., Cai, Z. H., Jiang, Y., & Xu, D. R. (1999). Organ culture of a precious Chinese medicinal plant – *Fritillaria unibracteata*. *Plant Cell, Tissue and Organ Culture*, 59, 197–201.

- Georgiev, M., Georgiev, V., Penchev, P., Antonova, D., Pavlov, A., Ilieva, M., & Popov, S. (2010). Volatile metabolic profiles of cell suspension cultures of *Lavandula vera*, *Nicotiana tabacum* and *Helianthus annuus*, cultivated under different regimes. *Engineering in Life Sciences*, *10*, 148–157.
- Gobbo-Neto, L., Guaratini, T., Pessoa, C., Moraes, M. O. D., Costa-Lotufo, L. V., Vieira, R. F., & Lopes, N. P. (2010). Differential metabolic and biological profiles of *Lychnophora ericoides* mart. (Asteraceae) from different localities in the Brazilian “campos rupestres”. *Journal of the Brazilian Chemical Society*, *21*, 750–759.
- Heidari, M., & Karami, V. (2014). Effects of different mycorrhiza species on grain yield, nutrient uptake and oil content of sunflower under water stress. *Journal of the Saudi Society of Agricultural Sciences*, *13*, 9–13.
- Herre, E. A., Mejia, I. C., Kylo, D. A., Rojas, E., Maynard, Z., Butler, L., & Van bael, S. A. (2007). Ecological implications of anti-pathogen effects of tropical fungal endophytes and mycorrhizae. *Ecology*, *88*, 550–558.
- Hibino, K., & Ushiyama, K. (1999). Commercial production of ginseng by plant tissue culture technology. In T.-J. Fu, G. Singh, & W. R. Curtis (Eds.), *Plant cell and tissue culture for the production of food ingredients* (Vol. 30, pp. 215–224). Dordrecht: Kluwer Academic Publishers.
- Hong, J. K., Yun, B. W., Kang, J. G., Raja, M. U., Kwon, E., Sorhagen, K., Chu, C., Wang, Y., & Loake, G. J. (2008). Nitric oxide function and signalling in plant disease resistance. *Journal of Experimental Botany*, *59*, 147–154.
- Huang, T. K., & McDonald, K. A. (2009). Bioreactor engineering for recombinant protein production in plant cell suspension cultures. *Biochemical Engineering Journal*, *45*, 168–184.
- Huang, W. Y., Cai, Y. Z., & Zhang, Y. (2010). Natural phenolic compounds from medicinal herbs and dietary plants: Potential use for cancer prevention. *Nutrition and Cancer*, *62*, 1–20.
- Huang, Z. A., Zhao, T., Fan, H. J., Wang, N., Zheng, S. S., & Ling, H. Q. (2012). The upregulation of NtAN2 expression at low temperature is required for anthocyanin accumulation in juvenile leaves of Lc-transgenic tobacco (*Nicotiana tabacum* L.). *Journal of Genetics and Genomics*, *39*, 149–156.
- Jang, H. R., Lee, H. J., Shohael, A. M., Park, B. J., Paek, K. Y., & Park, S. Y. (2016). Production of biomass and bioactive compounds from shoot cultures of *Rosa rugosa* using a bioreactor culture system. *Horticulture, Environment and Biotechnology*, *57*, 79–87.
- Jeong, G. T., & Park, D. H. (2006). Enhanced secondary metabolite biosynthesis by elicitation in transformed plant root system. In M. M. JD, W. S. Adney, J. R. Mielenz, & T. Klasson (Eds.), *Applied biochemistry and biotechnology* (pp. 436–446). New York: Humana Press.
- Jin, M. Y., Han, L., Li, H., Wang, H. Q., Piao, X. C., & Lian, M. L. (2017). Kinsenoside and polysaccharide production by rhizome culture of *Anoectochilus roxburghii* in continuous immersion bioreactor systems. *Plant Cell, Tissue and Organ Culture*, *131*, 527. <https://doi.org/10.1007/s11240-017-1302-8> (online first).
- Jose, S., & Thomas, T. D. (2014). Comparative phytochemical and antibacterial studies of two indigenous medicinal plants *Curcuma caesia* Roxb. and *Curcuma aeruginosa*. *International Journal of Green Pharmacy*, *8*, 65–71.
- Jouanin, L. (1984). Restriction map of an agropine-type Ri plasmid and its homologies with Ti plasmids. *Plasmid*, *12*, 91–102.
- Kaimoyo, E., Farag, M. A., Sumner, L. W., Wasmann, C., Cuello, J. L., & VanEtten, H. (2008). Sub-lethal levels of electric current elicit the biosynthesis of plant secondary metabolites. *Biotechnology Progress*, *24*, 377–384.
- Kajula, M., Tejesvi, M. V., Kolehmainen, S., Mäkinen, A., Hokkanen, J., Mattila, S., & Pirttilä, A. M. (2010). The siderophore ferricrocin produced by specific foliar endophytic fungi in vitro. *Fungal Biology*, *114*, 248–254.
- Karam, F. S., Haraguchi, M., & Gardner, D. (2011). Seasonal variation in pyrrolizidine alkaloid concentration and plant development in *Senecio madagascariensis* Poir. (Asteraceae) in Brazil. In F. Riet-Correa (Ed.), *Poisoning by plants, mycotoxins and related toxins* (pp. 179–185). Wallingford: CAB International.

- Karppinen, K., Hokkanen, J., Tolonen, A., Mattila, S., & Hohtola, A. (2007). Biosynthesis of hyperforin and adhyperforin from amino acid precursors in shoot cultures of *Hypericum perforatum*. *Phytochemistry*, *68*, 1038–1045.
- Karuppusamy, S. (2009). A review on trends in production of secondary metabolites from higher plants by in vitro tissue, organ and cell cultures. *Journal of Medicinal Plant Research*, *3*, 1222–1239.
- Karwasara, V. S., Jain, R., Tomar, P., & Dixit, V. K. (2010). Elicitation as yield enhancement strategy for glycyrrhizin production by cell cultures of *Abrus precatorius* Linn. *In Vitro Cellular and Development Biology – Plant*, *46*, 354–362.
- Keeling, C. I., Weisshar, S., Lin, R. P., & Bohlmann, J. (2008). Functional plasticity of paralogous diterpene synthases involved in conifer defense. *Proceedings of the National Academy of Sciences of the United States of America*, *105*, 1085–1090.
- Keles, L. C., Melo, N. I. D., Aguiar, G. D. P., Wakabayashi, K. A. L., Carvalho, C. E. D., Cunha, W. R., & Lopes, N. P. (2010). Lychnophorinae (Asteraceae): A survey of its chemical constituents and biological activities. *Quim Nova*, *33*, 2245–2260.
- Khalili, M., Hasanloo, T., & Kazemi Tabar, S. K. (2010). Ag {+} enhanced silymarin production in hairy root cultures of '*Silybum marianum*' (L.) gaertn. *Plant Omics*, *3*, 109.
- Kim, O. T., Kim, M. Y., Hong, M. H., Ahn, J. C., & Hwang, B. (2004). Stimulation of asiaticoside accumulation in the whole plant cultures of *Centella asiatica* (L.) Urban by elicitors. *Plant Cell Reports*, *23*, 339–344.
- Lambert, E., Faizal, A., & Geelen, D. (2011). Modulation of triterpene saponin production: In vitro cultures, elicitation, and metabolic engineering. *Applied Biochemistry and Biotechnology*, *164*, 220–237.
- Leicach, S. R., & Chludil, H. D. (2014). Plant secondary metabolites: Structure–activity relationships in human health prevention and treatment of common diseases. In Atta-ur-Rahman (Ed.), *Studies in natural products chemistry* (pp. 267–270). Amsterdam: Elsevier.
- Li, W., Koike, K., Asada, Y., Hirotsani, M., Rui, H., Yoshikawa, T., & Nikaido, T. (2002). Flavonoids from *Glycyrrhiza pallidiflora* hairy root cultures. *Phytochemistry*, *60*, 351–355.
- Lindsey, K. (1995). Manipulation by nutrient limitation of the biosynthetic activity of immobilized cells of *Capsicum frutescens* Mill. *ev. annum. Planta*, *165*, 126–133.
- LIN-WANG, K. U., Micheletti, D., Palmer, J., Volz, R., Lozano, L., Espley, R., & Iglesias. (2011). High temperature reduces apple fruit colour via modulation of the anthocyanin regulatory complex. *Plant, Cell & Environment*, *34*, 1176–1190.
- Long, R. M., & Croteau, R. (2005). Preliminary assessment of the C13-side chain 2'-hydroxylase involved in Taxol biosynthesis. *Biochemical and Biophysical Research Communications*, *338*, 410–417.
- Lucena, R. B., Rissi, D. R., Maia, L. A., Flores, M. M., Dantas, A. F. M., Nobre, V. M. D. T., & Barros, C. S. (2010). Poisoning by pyrrolizidine alkaloids in ruminants and horses in Brazil. *Pesquisa Veterinaria Brasileira*, *30*, 447–452.
- Ludwing-Muller, J., Jahn, L., Lippert, A., Puschel, J., & Walter, A. (2014). Improvement of hairy root cultures and plants by changing biosynthetic pathways leading to pharmaceutical metabolites: Strategies and applications. *Biotechnology Advances*, *32*, 1168–1179.
- Lulu, T., Park, S. Y., Ibrahim, R., & Paek, K. Y. (2015). Production of biomass and bioactive compounds from adventitious roots by optimization of culturing conditions of *Eurycoma longifolia* in balloon-type bubble bioreactor system. *Journal of Bioscience and Bioengineering*, *119*, 712–717.
- Mahajan, V., Sharma, N., Kumar, S., Bhardwaj, V., Ali, A., Khajuria, R. K., Bedi, Y. S., Vishwakarma, R. A., & Gandhi, S. G. (2015). Production of rohitukine in leaves and seeds of *Dysoxylum binectariferum*: An alternate renewable resource. *Pharmaceutical Biology*, *53*, 446–450.
- Maier, W., Peipp, H., Schimidt, J., Wray, V., & Strack, D. (1995). Levels of terpenoid glycoside (blumenin) and cell wall-bound phenolics in some cereal mycorrhizas. *Plant Physiology*, *109*, 465–470.