


Association for Women in Mathematics Series

Alyson Deines · Daniela Ferrero  
Erica Graham · Mee Seong Im  
Carrie Manore · Candice Price *Editors*

# Advances in the Mathematical Sciences

AWM Research Symposium, Los Angeles, CA,  
April 2017



# **Association for Women in Mathematics Series**

---

## **Volume 15**

---

### **Series Editor**

Kristin Lauter  
Microsoft Research  
Redmond, Washington, USA

## Association for Women in Mathematics Series

---

Focusing on the groundbreaking work of women in mathematics past, present, and future, Springer's Association for Women in Mathematics Series presents the latest research and proceedings of conferences worldwide organized by the Association for Women in Mathematics (AWM). All works are peer-reviewed to meet the highest standards of scientific literature, while presenting topics at the cutting edge of pure and applied mathematics. Since its inception in 1971, The Association for Women in Mathematics has been a non-profit organization designed to help encourage women and girls to study and pursue active careers in mathematics and the mathematical sciences and to promote equal opportunity and equal treatment of women and girls in the mathematical sciences. Currently, the organization represents more than 3000 members and 200 institutions constituting a broad spectrum of the mathematical community, in the United States and around the world.

More information about this series at <http://www.springer.com/series/13764>

Alyson Deines • Daniela Ferrero • Erica Graham  
Mee Seong Im • Carrie Manore • Candice Price  
Editors

# Advances in the Mathematical Sciences

AWM Research Symposium, Los Angeles,  
CA, April 2017



*Editors*

Alyson Deines  
Center for Communications Res  
San Diego, CA, USA

Erica Graham  
Bryn Mawr College  
Ardmore, PA, USA

Carrie Manore  
Los Alamos National Laboratory  
Los Alamos, NM, USA

Daniela Ferrero  
Department of Mathematics  
Texas State University  
San Marcos, TX, USA

Mee Seong Im  
United States Military Academy  
New York, NY, USA

Candice Price  
Department of Mathematics  
University of San Diego  
San Diego, CA, USA

ISSN 2364-5733

ISSN 2364-5741 (electronic)

Association for Women in Mathematics Series

ISBN 978-3-319-98683-8

ISBN 978-3-319-98684-5 (eBook)

<https://doi.org/10.1007/978-3-319-98684-5>

Library of Congress Control Number: 2018959855

Mathematics Subject Classification: 05E05, 05E10, 14Q99, 16G20, 18E10, 20F65, 20G40, 34G10, 57M27, 92B25, 92C20, 92D20, 92E20, 97D40

© The Author(s) and the Association for Women in Mathematics 2018

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors, and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG  
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland



# Preface

This volume highlights the mathematical research presented at the 2017 Association for Women in Mathematics (AWM) Research Symposium. This event, fourth in the biennial series of AWM Research Symposia, was held at the University of California Los Angeles (UCLA) on April 8–9, 2017. The objective of the AWM Research Symposia Series is to increase awareness of the mathematical achievements of women in academia, industry, and government, as well as to provide a supportive environment for female mathematicians, at all stages of their careers, to share their research. Additionally, these symposia promote research collaboration, as they facilitate the creation of new networks of women researchers and support the research collaboration networks already existing in several areas of mathematics. The symposia also include social events to enable networking among women in different career paths, or at different career stages, while promoting the discussion of prospects, visibility, and recognition.

## About the 2017 AWM Research Symposium

The 2017 AWM Research Symposium was organized by Raegan Higgins, Kristin Lauter, Magnhild Lien, Ami Radunskaya, Tatiana Toro, Luminita Vese, and Carol Woodward. The Department of Mathematics at UCLA and the Institute for Pure and Applied Mathematics (IPAM) hosted the event that featured 4 plenary talks by distinguished women mathematicians (Table 1), 19 special sessions on a broad range of areas in mathematics (Table 2), a poster sessions for graduate students and recent Ph.D.s, a job panel, a reception, a banquet, and a student chapter event.

Eight of the 19 special sessions were organized by research networks supported by the AWM ADVANCE grant: Women in Numbers (WIN), Women in Math Biology (WIMB), Women in Noncommutative Algebra and Representation Theory (WINART), Women in Numerical Analysis and Scientific Computing (WINASC), Women in Computational Topology (WinCompTop), Women in Topology (WIT),

**Table 1** 2017 AWM Research Symposium: plenary talks

Speaker	Title
Ruth Charney	Searching for hyperbolicity
Svitlana Marboroda	The hidden landscape of localization of eigenfunctions
Linda Petzold	Inference of the functional network controlling circadian rhythm
Mariel Vazquez	Understanding DNA topology

Women in Shape (WiSh), and Algebraic Combinatorixx (ACxx). For more details about these and other Research Collaboration Networks for Women visit: <https://awmadvance.org/research-networks/>.

The keynote speaker at the banquet was Maria Helena Noronha, who in a very inspirational address described her journey from Brazil to Southern California, as a researcher and as a mentor. During the banquet, the second AWM Presidential Award was presented to Deanna Haunsperger, in recognition of her contribution to advance the goals of AWM through her work in the Summer Math Program (SMP) at Carleton College.

The symposium also included a full-day session titled *Wikipedia edit-a-thon*, during which participants at the 2017 AWM Research Symposium took turns writing Wikipedia entries to enhance the visibility of women in mathematics and their contributions.

## About This Volume

The first chapter in this volume corresponds to the opening plenary talk at the symposium, *Searching for Hyperbolicity* by Ruth Charney. Her chapter is an excellent introduction to geometric group theory, by a renowned expert in the field, whose research contributed to the consolidation of geometric group theory as a mathematical area.

The following three chapters comprise mathematical results presented at the *WINART Special Session: Representations of Algebras*. Mee Seong Im and Angela Wu present their work on representation theory of the generalized iterated wreath product of cyclic groups (chapter “Generalized Iterated Wreath Products of Cyclic Groups and Rooted Trees Correspondence”) and symmetric groups (chapter “Generalized Iterated Wreath Products of Symmetric Groups and Generalized Rooted Trees Correspondence”). Chapter “Conway–Coxeter Friezes and Mutation: A Survey” consists of a survey on Conway–Coxeter friezes and mutation, in which Karin Baur, Eleonore Faber, Sira Gratz, Khrystyna Serhiyenko, and Gordana Todorov connect Conway–Coxeter friezes, introduced in combinatorics in the 1970s, and cluster combinatorics, arising from the introduction of cluster algebras in the early 2000s.

**Table 2** 2017 AWM Research Symposium: special sessions

Title	Organizers
WIN—Work from Women in Numbers	Beth Malmskog, Katherine Stange
WinCompTop: Applications of Topology and Geometry	Emilie Purvine, Radmila Sazdanovic, Shirley Yap
WIMB—From Cells to Landscapes: Modeling Health and Disease	Erica Graham, Carrie Manore
ACxx: Algebraic Combinatorics	Hélène Barcelo, Gizem Karaali
WINASC: Recent Research Development on Numerical Partial Differential Equations and Scientific Computing	Chiu-Yen Kao, Yekaterina Epshteyn
WINART: Representations of Algebras	Susan Montgomery, Maria Vega
WiSh: Shape Modeling and Applications	Asli Genctav, Kathryn Leonard
WIT—Topics in Homotopy Theory	Julie Bergner, Angelica Osorno
Women in Sage Math	Alyson Deines, Anna Haensch
Women in Government Labs	Cindy Phillips, Carol Woodward
EDGE-y Mathematics: A Tribute to Dr. Sylvia Bozeman and Dr. Rhonda Hughes	Alejandra Alvarado, Candice Price
SMPosium: A Celebration of the Summer Mathematics Program for Women	Alissa S. Crans, Pamela A. Richardson
The Many Facets of Statistics Applied, Pure and BIG	Monica Jackson, Jo Hardin
History of Mathematics	Janet Beery
Commutative Algebra	Alexandra Seceleanu, Emily Witt
Biological Oscillations Across Time Scales	Tanya Leise, Stephanie Taylor
Geometric Group Theory	Pallavi Dani, Tullia Dymarz, Talia Fernos
Recent Progress in Several Complex Variables	Purvi Gupta, Loredana Lanzani
Research in Collegiate Mathematics Education	Shandy Hauk, Pao-sheng Hsu

Chapter “Orbit Decompositions of Unipotent Elements in the Generalized Symmetric Spaces of  $SL_2(\mathbb{F}_q)$ ” features the work of Catherine Buell, Vicky Klima, Jennifer Schaefer, Carmen Wright, and Ellen Ziliak, who jointly studied the orbit decompositions of unipotent elements in the generalized symmetric spaces of  $SL_2(F_q)$ . Their results were presented in the special session *EDGE-y Mathematics: A Tribute to Dr. Sylvia Bozeman and Dr. Rhonda Hughes*. Chapter “A Characterization of the  $U(\Omega, m)$  Sets of a Hyperelliptic Curve as  $\Omega$  and  $m$  Vary” is a paper on computational algebraic geometry by Christelle Vincent, and it was part of the special session *Women in Sage*. In “A First Step Toward Higher Order Chain Rules in Abelian Functor Calculus”, Christine Osborne and Amelia Tebbe take a first step toward higher-order chain rules in abelian functor calculus, by proving the second-order directional derivative chain rule using concrete computational techniques. Their work was presented in the *WIT-Topics in Homotopy Theory*.

In “DNA Topology Review” the volume transitions to mathematical biology, as Garrett Jones and Candice Price survey how topology can be applied to model biological processes, such as actions of proteins on DNA. Their work introduces essential concepts together with a description of fundamental literature, offering an excellent guide for undergraduate or graduate students, as well as for scholars, interested in learning the basics of DNA topology. Chapter “Structural Identifiability Analysis of a Labeled Oral Minimal Model for Quantifying Hepatic Insulin Resistance” features a paper presented at the special session *From Cells to Landscapes: Modeling Health and Disease*. Jacqueline Simens, Melanie Cree-Green, Bryan Bergman, Kristen Nadeau, and Cecilia Diniz Behn present a structural identifiability analysis of a labeled oral minimal model for quantifying hepatic insulin resistance. Their research contributes to the understanding of aging, trauma, and many diseases, such as obesity and type 2 diabetes.

The following two chapters correspond to the special session *Biological Oscillations Across Time Scales*. Chapter “Spike-Field Coherence and Firing Rate Profiles of CA1 Interneurons During an Associative Memory Task” features a study of the spike-field coherence and firing rate profiles of CA1 interneurons during an associative memory task, by Pamela Riviere and Lara Rangel. Their work analyzes whether inhibitory interneurons from the CA1 region of the hippocampus contain information about task dimensions in their firing rates. Following this work is a computational study of learning-induced sequence reactivation during sharp-wave ripple activity, as observed during sleep states, by Paola Malerba, Katya Tsimring, and Maxim Bazhenov. The authors use a model of spiking neuron networks of excitatory and inhibitory neurons in the CA3 and CA1 regions of the hippocampus to study the firing behavior of neurons during sharp-wave-ripple activity.

Chapter “Learning-Induced Sequence Reactivation During Sharp-Wave Ripples: A Computational Study” corresponds to a presentation in the special session *WINASC: Recent Research Development on Numerical Partial Differential Equations and Scientific Computing*. It consists of the work by Beatrice Riviere and Xin Yang using a DG method for the simulation of CO<sub>2</sub> storage in a saline aquifer. Their work has important practical applications, since porous media, such as saline aquifers or oil and gas reservoirs, are a major cause of the excessive amount of carbon dioxide in the atmosphere.

In “A DG Method for the Simulation of CO<sub>2</sub> Storage in Saline Aquifer”, Beth M. Campbell Hetrick presents a study of regularization for an ill-posed inhomogeneous Cauchy problem, extending previous results for the homogeneous problem to the inhomogeneous case. While results and numerical experiments in Hilbert space are plentiful, this chapter contains regularization results for inhomogenous ill-posed problems for true Banach space, where little is known and many exciting problems await solutions.

The 2017 AWM Research Symposium included a special session on mathematics education comprising six presentations on various theoretical perspectives on the nature of human cognition and knowledge structures. Research methods ranged from individual interview and classroom observation to national survey and in-depth

study of a particular instance or case. The span of topics covered calculus, combinatorics, linear algebra, foundations of proof, application of mathematics to teaching, and development of future teachers. This volume concludes with a chapter in which Shandy Hauk, Chris Rasmussen, Nicole Engelke Infante, Elise Lockwood, Michelle Zandieh, Stacy Brown, Yvonne Lai, and Pao-sheng Hsu offer highlights of the six presentations in this session.

## Acknowledgments

All chapters in this volume have been peer-reviewed. The editors thank the authors who submitted their manuscripts and the referees who carefully and thoroughly reviewed them. Their highly valuable work was essential to ensure the quality of this volume meets the high standards that characterize the AWM Springer Series.

The editors express their gratitude to the organizers of the symposium and the hosting institutions, whose dedication and hard work were central to the success of the 2017 AWM Research Symposium. They also extend their appreciation to the National Science Foundation (NSF), the National Security Agency (NSA), and the Mathematical Sciences Research Institute (MSRI) for their funding to support special sessions. The editors extend their appreciation to Microsoft for their help funding the student event, to Wolfram and Maple for sponsoring prizes for best poster contest, and to Springer, Oxford University Press, Basic Books, AMS, and *Mathematical Reviews* for presenting booths in the exhibit area.

Special thanks to Kristin Lauter, editor of the AWM Springer Series, for trusting us with the task of publishing this volume and to Jeffrey Taub and Dimana Tzvetkova, our editors at Springer, for their guidance and help while editing these proceedings.

San Diego, CA, USA  
San Marcos, TX, USA  
Ardmore, PA, USA  
New York, NY, USA  
Los Alamos, NM, USA  
San Diego, CA, USA  
July 2018

Alyson Deines  
Daniela Ferrero  
Erica Graham  
Mee Seong Im  
Carrie Manore  
Candice Price

# Contents

<b>Searching for Hyperbolicity</b> .....	1
Ruth Charney	
<b>Generalized Iterated Wreath Products of Cyclic Groups and Rooted Trees Correspondence</b> .....	15
Mee Seong Im and Angela Wu	
<b>Generalized Iterated Wreath Products of Symmetric Groups and Generalized Rooted Trees Correspondence</b> .....	29
Mee Seong Im and Angela Wu	
<b>Conway–Coxeter Friezes and Mutation: A Survey</b> .....	47
Karin Baur, Eleonore Faber, Sira Gratz, Khrystyna Serhiyenko, and Gordana Todorov	
<b>Orbit Decompositions of Unipotent Elements in the Generalized Symmetric Spaces of <math>SL_2(\mathbb{F}_q)</math></b> .....	69
Catherine Buell, Vicky Klima, Jennifer Schaefer, Carmen Wright, and Ellen Ziliak	
<b>A Characterization of the <math>U(\Omega, m)</math> Sets of a Hyperelliptic Curve as <math>\Omega</math> and <math>m</math> Vary</b> .....	79
Christelle Vincent	
<b>A First Step Toward Higher Order Chain Rules in Abelian Functor Calculus</b> .....	97
Christina Osborne and Amelia Tebbe	
<b>DNA Topology Review</b> .....	121
Garrett Jones and Candice Renée Price	
<b>Structural Identifiability Analysis of a Labeled Oral Minimal Model for Quantifying Hepatic Insulin Resistance</b> .....	145
Jacqueline L. Simens, Melanie Cree-Green, Bryan C. Bergman, Kristen J. Nadeau, and Cecilia Diniz Behn	

<b>Spike-Field Coherence and Firing Rate Profiles of CA1 Interneurons During an Associative Memory Task .....</b>	161
Pamela D. Rivière and Lara M. Rangel	
<b>Learning-Induced Sequence Reactivation During Sharp-Wave Ripples: A Computational Study .....</b>	173
Paola Malerba, Katya Tsimring, and Maxim Bazhenov	
<b>A DG Method for the Simulation of CO<sub>2</sub> Storage in Saline Aquifer .....</b>	205
Beatrice Riviere and Xin Yang	
<b>Regularization Results for Inhomogeneous Ill-Posed Problems in Banach Space .....</b>	233
Beth M. Campbell Hetrick	
<b>Research in Collegiate Mathematics Education .....</b>	245
Shandy Hauk, Chris Rasmussen, Nicole Engelke Infante, Elise Lockwood, Michelle Zandieh, Stacy Brown, Yvonne Lai, and Pao-sheng Hsu	
<b>Index .....</b>	269

# Searching for Hyperbolicity



Ruth Charney

**Abstract** This paper is an expanded version of a talk given at the AWM Research Symposium 2017. It is intended as a gentle introduction to geometric group theory with a focus on the notion of hyperbolicity, a theme that has inspired the field from its inception to current-day research. The last section includes a discussion of some current approaches to extending techniques from hyperbolic groups to more general classes of groups.

## 1 Introduction

This paper is an expanded version of a talk given at the AWM Research Symposium 2017. It is intended as a gentle introduction to geometric group theory for the non-expert, with a focus on the notion of hyperbolicity. Geometric group theory came into its own in the 1990s, in large part due to a seminal paper of Mikhail Gromov [14]. While the field has grown considerably since that time, hyperbolicity remains a central theme and continues to drive much current research in the field.

As the name suggests, geometric group theory provides a bridge between groups viewed as algebraic objects and geometry. Groups arise in all areas of mathematics and can be described in many different ways. Some arise purely algebraically (such as certain matrix groups), others have combinatorial descriptions (via presentations), and still others are defined topologically (such as fundamental groups of topological spaces). Geometric group theory is based on the principle that if a group acts as symmetries of some geometric object, then one can use geometry to better understand the group.

For many groups, it is easy to find such an action. The symmetric group on  $n$ -letters acts by symmetries on an  $n$ -simplex and the dihedral group of order  $2n$  is the symmetry group of a regular  $n$ -gon. This is also the case for some infinite groups.

---

R. Charney (✉)  
Brandeis University, Waltham, MA, USA  
e-mail: [charney@brandeis.edu](mailto:charney@brandeis.edu)

The free abelian group  $\mathbb{Z}^n$  acts by translation on  $\mathbb{R}^n$  (preserving the Euclidean metric), and the free group on  $n$ -generators acts by translation on a regular tree of valence  $2n$  with edges of length one.

So the first question one might ask is, when can one find such an action? Given an abstract group  $G$ , can we always realize  $G$  as a group of symmetries of some geometric object? As we will see below, the answer is yes. However, some geometric objects are more useful than others for this purpose. In the early 1900s, Max Dehn was interested in groups arising as fundamental groups of hyperbolic surfaces. These groups act by isometries on the hyperbolic plane  $\mathbb{H}^2$ . Dehn used the geometry of the hyperbolic plane to prove some amazing properties of these groups. We will discuss one of these results in Sect. 3 below. Decades later, these ideas motivated Gromov, who introduced the notion of a hyperbolic metric space. He showed that the properties that Dehn deduced held more generally for any group acting nicely on such a space.

Gromov's notion of hyperbolic spaces and hyperbolic groups have been much studied since that time. Many well-known groups, such as mapping class groups and fundamental groups of surfaces with cusps, do not meet Gromov's criteria, but nonetheless display some hyperbolic behavior. In recent years, there has been much interest in capturing and using this hyperbolic behavior wherever and however it occurs.

In this paper, I will review some basic notions in geometric group theory, discuss Dehn's work and Gromov's notion of hyperbolicity, then introduce the reader to some recent developments in the search for hyperbolicity. My goal is to be comprehensible, not comprehensive. For those interested in learning more about the subject, I recommend [1] and [6] for a general introduction to geometric group theory and [7] and [13] for more detail about hyperbolic groups.

## 2 Geodesic Metric Spaces, Isometries and Quasi-Isometries

We begin with some basic definitions. Let  $X$  be a metric space with distance function  $d : X \times X \rightarrow \mathbb{R}$ . A *geodesic* in  $X$  is a distance preserving map from an interval  $I \subset \mathbb{R}$  into  $X$ , that is, a map  $\alpha : I \rightarrow X$  such that for all  $t_1, t_2 \in I$ ,

$$d(\alpha(t_1), \alpha(t_2)) = |t_1 - t_2|.$$

The interval  $I$  may be finite or infinite. This definition is analogous to the notion of a geodesic in a Riemannian manifold. In particular, a geodesic between two points in  $X$  is a length-minimizing path.

A *geodesic metric space* is a metric space  $X$  in which any two points are connected by a geodesic. For such metric spaces, the distance is intrinsic to the space; the distance between any two points is equal to the minimal length of a path connecting them. Often, we also require that our metric space be *proper*, that is, closed balls in  $X$  are compact.

*Example 1* Consider the unit circle  $S^1$  in the plane. There are two natural metrics we could put on  $S^1$ . The first is the induced Euclidean metric: the distance between two points is the length of the straight line in  $\mathbb{R}^2$  between them. The other is the arc length metric: the distance between two points is the length of the (shortest) circular arc between them. The first of these is not a geodesic metric (since, for example, there is no path in  $S^1$  of length 2 connecting a pair of antipodal points) whereas the second one is geodesic.

*Example 2* Suppose  $\Gamma$  is a connected graph. There is a natural geodesic metric on  $\Gamma$  obtained by identifying each edge with a copy of the unit interval  $[0, 1]$  and defining the distance between any two points in  $\Gamma$  to be the length of the shortest path between them. This metric is proper if and only if each vertex has finite valence.

*Example 3* Let  $M$  be a complete Riemannian manifold. Then the usual distance function given by minimizing path lengths is a proper, geodesic metric on  $M$ .

A map between two metric spaces  $f : X \rightarrow Y$  is an *isometry* if it is bijective and preserves distances. In lay terms, an isometry of  $X$  to itself is a “symmetry” of  $X$ . These symmetries form a group under composition.

Now suppose we are given a group  $G$ . Our first goal is to find a nice metric space on which  $G$  acts as a group of symmetries. Sometimes, such an action arises naturally. For example, suppose  $G$  is the fundamental group of a Riemannian manifold  $M$ . Then passing to the universal cover  $\tilde{M}$ , we get an action of  $G$  by deck transformations on  $\tilde{M}$ . This action is distance preserving since it takes geodesic paths to geodesic paths.

More generally, the same works for the fundamental group of any geodesic metric space  $X$  that admits a universal cover. The universal cover  $\tilde{X}$  inherits a geodesic metric such that the projection to  $X$  is a local isometry, and the deck transformations act isometrically on  $\tilde{X}$ .

*Example 4* Consider the free group on two generators  $F_2$ . This group is the fundamental group of a wedge of two circles,  $S^1 \vee S^1$ , so it acts by isometries on the universal cover, namely the regular 4-valent tree.

In general, a group  $G$  can act by isometries on a variety of different geodesic metric spaces. Some of these actions, however, are not helpful in studying the group. For example, any group acts on a single point! To have any hope that the geometry of the space will produce information about the group, we will need some extra conditions on the action.

**Definition 1** A group  $G$  is said to act *geometrically* on a metric space  $X$  if the action satisfies the following three properties.

- *isometric*: Each  $g \in G$  acts as an isometry on  $X$ .
- *proper*: For all  $x \in X$ , there exists  $r > 0$  such that  $\{g \in G \mid B(x, r) \cap gB(x, r) \neq \emptyset\}$  is finite, where  $B(x, r)$  denotes the ball of radius  $r$  centered at  $x$ .
- *cocompact*: There exists a compact set  $K \subset X$  whose translates by  $G$  cover  $X$  (or equivalently,  $X/G$  is compact).

In particular, the fundamental group of a compact metric space  $X$  acts geometrically on the universal covering space  $\tilde{X}$ . But now suppose that our group  $G$  arises purely algebraically. How can we find a metric space  $X$  on which  $G$  acts geometrically? The following is a construction that works for any finitely generated group.


Choose a finite generating set  $S$  for  $G$ . Define the *Cayley graph* for  $G$  with respect to  $S$  to be the graph  $\Gamma_S(G)$  whose vertices are in one-to-one correspondence with the elements of  $G$  and for each  $s \in S$ ,  $g \in G$ , there is an edge (labelled by  $s$ ) connecting the vertex  $g$  to the vertex  $gs$ .

$G$  acts on  $\Gamma_S(G)$  by left multiplication on the vertices. That is,  $h \in G$  maps the vertex labelled  $g$  to the vertex labelled  $hg$ . Note that  $h$  takes edges to edges, the edge connecting  $g$  to  $gs$  maps to the edge connecting  $hg$  to  $hgs$ . Thus, if we put the path-length metric on  $\Gamma_S(G)$  as described above, then this action preserves the metric and is easily seen to be geometric.


*Example 5* Consider the free abelian group  $\mathbb{Z}^2$  with generating set  $S = \{(1, 0), (0, 1)\}$ . Viewing  $\mathbb{Z}^2$  as the set of integer points in the plane  $\mathbb{R}^2$ , we can identify the Cayley graph of  $\mathbb{Z}^2$  with the square grid connecting these points. Distances are measured by path lengths in this grid, so the distance from  $(0, 0)$  to  $(n, m)$  is  $|n| + |m|$ . Note that there are, in general, many geodesic paths between any two points. See Fig. 1.

*Example 6* Let  $F_2$  be the free group with generating set  $S = \{a, b\}$ . View  $F_2$  as the fundamental group of the wedge of two circles labelled  $a$  and  $b$ . Lifting these labels to the universal covering space, we get a 4-valent tree, as in Fig. 2, with every red edge labelled  $a$  and every blue edge labelled  $b$  (Metrically, you should picture every edge as having the same length.) This tree is precisely the Cayley graph  $\Gamma_S(F_2)$ . To

**Fig. 1** Geodesics from  $x$  to  $y$  in the Cayley graph of  $\mathbb{Z}^2$



**Fig. 2** Cayley graph of  $F_2$



see this, choose a base vertex  $v$  and identify each vertex with the element of  $F_2$  that translates  $v$  to that vertex.

Clearly, the Cayley graph depends on the choice of generating set. For example, were we to add a third generator  $(1, 1)$  to our generating set for  $\mathbb{Z}^2$ , the Cayley graph would get additional edges which cross the grid diagonally and would hence change the distance between vertices.

This could be a cause for concern; we seek geometric properties of the Cayley graph that are intrinsic to the group, so they should not be dependent on a choice of generating set. Luckily, in the case of a finitely generated group, replacing one finite generating set by another does not distort distances too badly. This leads to a fundamental concept in geometric group theory.

**Definition 2** A map  $f : X \rightarrow Y$  between two metric spaces is a *quasi-isometric embedding* if there exists constants  $K, C$  such that for all  $x, z \in X$

$$\frac{1}{K} d_X(x, z) - C \leq d_Y(f(x), f(z)) \leq K d_X(x, z) + C.$$


If in addition, every point in  $Y$  lies within  $C$  of some point in  $f(X)$ , then  $f$  is a *quasi-isometry*. In this case we write  $X \sim_{QI} Y$ .

It can be shown that any quasi-isometry has a “quasi-inverse,” so the relation  $X \sim_{QI} Y$  is an equivalence relation. We remark that quasi-isometries need not be continuous maps.

*Example 7* Consider the inclusion of the integer grid into the plane  $\mathbb{R}^2$ . This is a quasi-isometry. The quasi-inverse is a discontinuous map sending the interior of each square to its boundary.

*Example 8* Consider the graph in Fig. 3. Collapsing each of the triangles to a point gives a quasi-isometry of this graph onto the 3-valent tree. We call such a graph a quasi-tree.

It is an easy exercise to show that if  $G$  is a finitely generated group, then the Cayley graphs with respect to different finite generating sets are all quasi-isometric. (In fact, they are bi-Lipschitz, i.e., we can take the maps to be continuous and the constant  $C$  to be zero.) Thus, identifying  $G$  with the vertex set in its Cayley graph, we can view  $G$  itself as a metric space and this metric is well-defined up to quasi-isometry. In fact, we have the following more general statement.



**Fig. 3** A graph quasi-isometric to a tree

**Proposition 1 (Milnor-Švarc Lemma)** *Suppose  $G$  acts geometrically on a geodesic metric space  $X$ . Then  $G$  is finitely generated and for any choice of basepoint  $x_0 \in X$ , the map  $G \rightarrow X$  taking  $g \mapsto gx_0$  is a quasi-isometry.*

As a result, the notion of quasi-isometry is a fundamental concept in geometric group theory. The properties of a group that one can hope to glean from its action on a metric space are generally properties preserved by quasi-isometries. Moreover, since any finite index subgroup  $H < G$  is quasi-isometric to  $G$ , often the strongest statement we can make regarding a property  $(P)$  is that our group  $G$  is *virtually*  $(P)$ , meaning that some finite index subgroup of  $G$  satisfies property  $(P)$ .

The classification of finitely generated groups up to quasi-isometry is a meta-problem in the field. It is easy to see that any two groups that are commensurable (i.e., they contain subgroups of finite index that are isomorphic) are quasi-isometric. So a related problem is the question of rigidity: for a given group  $G$ , is every group quasi-isometric to  $G$  also commensurable to  $G$ ?

### 3 Hyperbolic Groups

Once we have our group acting geometrically on a metric space, we can ask how geometric properties of the space are reflected in algebraic or combinatorial properties of the group. The classical example of this comes from the work of Max Dehn [12]. Dehn was interested in fundamental groups of surfaces. A closed orientable surface of genus  $g \geq 2$  (i.e. a torus with  $g$  holes) can be given a Riemannian metric of constant curvature  $-1$  and its universal covering space can be identified with the hyperbolic plane  $\mathbb{H}^2$ . This gives a geometric action of the fundamental group on  $\mathbb{H}^2$ . Using geometric properties of the hyperbolic plane, Dehn proved some very strong combinatorial properties for these groups. I will describe two of his results here.

One way to describe a group is by means of a presentation. Given a set of generators  $S$  for a group  $G$ , there is a natural surjection from the free group  $F(S)$  onto  $G$ . The kernel  $K$  of this map is a normal subgroup of  $F(S)$ . Formally, a *presentation* of  $G$  consists of a generating set  $S$ , together with a set  $R \subset F(S)$  such that  $R$  generates  $K$  as a normal subgroup, or in other words,  $G$  is the quotient of  $F(S)$  by the normal closure of  $R$ . The elements of  $R$  are called *relators*. We denote such a presentation for  $G$  by writing

$$G = \langle S \mid R \rangle.$$

In practice, we usually indicate  $R$  by a set of equations that hold in  $G$ . Viewing elements of  $F(S)$  as “words” in the alphabet  $S \cup S^{-1}$ , the elements in  $R$  are words that are equal to the identity in  $G$ . However, many other words, such as products and conjugates of those in  $R$ , are also equal to the identity in  $G$ . The idea is that *all* relations among the generators that hold in  $G$  should be consequences of the ones listed in the presentation.

Here are some examples. The cyclic group of order  $n$  has presentation

$$\mathbb{Z}/n\mathbb{Z} = \langle s \mid s^n = 1 \rangle$$

while the free abelian group on two generators has presentation

$$\mathbb{Z}^2 = \langle a, b \mid ab = ba \rangle.$$

Can you recognize the following group?

$$G = \langle u, v \mid u^4 = 1, u^2 = v^3 \rangle.$$

It turns out that this group is isomorphic to the special linear group  $SL(2, \mathbb{Z})$ . The isomorphism is given by identifying

$$u = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \quad v = \begin{bmatrix} 0 & -1 \\ 1 & 1 \end{bmatrix}$$

Every group can be described by a presentation, though in general  $S$  and  $R$  need not be finite. Presentations can be extremely useful, and are the starting point for combinatorial group theory. On the other hand, presentations can sometimes be very mysterious and frustratingly difficult to decipher. For example, consider the following questions.

1. *The Word Problem:* Given a finite presentation  $\langle S \mid R \rangle$  and a word  $w$  in  $F(S)$ , is there an algorithm to decide whether  $w$  represents the identity element in  $G$ ?
2. *The Isomorphism Problem:* Given two finite presentations  $\langle S \mid R \rangle$  and  $\langle S' \mid R' \rangle$ , is there an algorithm to decide whether the groups they represent are isomorphic?

It turns out that there are groups for which no such algorithms exist. In this case we say the Word Problem (or the Isomorphism Problem) is unsolvable. What Dehn showed, using the geometry of hyperbolic space, was that for fundamental groups of hyperbolic surfaces, both of these problems are solvable. Moreover, he showed that for an appropriate choice of presentation, the word problem has a particularly nice solution. Namely, any word  $w \in F(S)$  that represents the identity element in  $G$  must contain more than half of a relator  $r$ , and hence can be shortened by applying the equation  $r = 1$ . It follows that the word problem is solvable in linear time, that is, the time it takes to decide whether  $w \in F(S)$  represents the identity element in  $G$ , is linear in the length of  $w$ . An algorithm of this type is now known as a Dehn's algorithm.

Some 75 years later, Mikhail Gromov made a startling observation: the only property of the hyperbolic plane that Dehn really needed to derive his results was the fact that triangles in  $\mathbb{H}^2$ , no matter how far apart their vertices may be, are always “thin.” And from this observation, the modern field of hyperbolic geometry (and more generally geometric group theory) was born.

What do we mean by “thin”? Let  $X$  be any geodesic metric space. A triangle  $T(a, b, c)$  in  $X$  consists of three vertices  $a, b, c \in X$  together with a choice of geodesics connecting them.

**Definition 3** Let  $X$  be a geodesic metric space and let  $\delta \geq 0$ . We say a triangle  $T(a, b, c)$  in  $X$  is  $\delta$ -thin if each side of  $T$  lies in the union of the  $\delta$ -neighborhoods of the other two sides. (See Fig. 4.)

One can show that in  $\mathbb{H}^2$ , every triangle, even those with vertices at infinity (ideal triangles) are  $\delta$ -thin for  $\delta = \ln(1 + \sqrt{2})$ . This fact was crucial to Dehn’s work.

This brings us finally to Gromov’s notion of hyperbolicity.


**Definition 4** A geodesic metric space  $X$  is  $\delta$ -hyperbolic if every triangle in  $X$  is  $\delta$ -thin. We say  $X$  is *hyperbolic* if it is  $\delta$ -hyperbolic for some  $\delta$ . A finitely generated group  $G$  is *hyperbolic* if it acts geometrically on a hyperbolic metric space.

One can show that if two spaces are quasi-isometric and one of them is hyperbolic, then so is the other (though the constant  $\delta$  may change). In particular, a finitely generated group  $G$  is hyperbolic if and only if some (hence any) Cayley graph of  $G$  is hyperbolic.


*Example 9* We begin with a trivial example. Any bounded metric space  $X$  is  $\delta$ -hyperbolic where  $\delta$  is the diameter of  $X$ , and hence any finite group is hyperbolic.

*Example 10* Let  $X$  be an infinite tree. Then for any three points  $a, b, c$  in  $X$ , the triangle connecting them degenerates into a tripod and is hence 0-hyperbolic! (See Fig. 5.) Since the Cayley graph of a free group is a tree, it follows that finitely generated free groups are hyperbolic.

**Fig. 4** A  $\delta$ -thin triangle



**Fig. 5** A triangle in a tree is 0-thin



*Example 11* Recall the presentation of  $SL(2, \mathbb{Z})$  given above. The center of  $SL(2, \mathbb{Z})$  is the order two subgroup generated by  $u^2 = v^3$ . Modding out by this subgroup gives the group  $PSL(2, \mathbb{Z})$  with presentation

$$PSL(2, \mathbb{Z}) = \langle u, v \mid u^2 = v^3 = 1 \rangle.$$

The Cayley graph of  $PSL(2, \mathbb{Z})$  with respect to this generating set is the quasi-tree drawn in Fig. 3 (continued out to infinity), with the edges of triangles labelled  $v$  and the remaining edges labeled  $u$ .  $SL(2, \mathbb{Z})$  also acts geometrically on this quasi-tree (with the center acting trivially), so  $SL(2, \mathbb{Z})$  and  $PSL(2, \mathbb{Z})$  are both hyperbolic groups.

*Example 12* Here is a non-example. Let  $\mathbb{R}^2$  be the plane with the standard Euclidean metric. Taking larger and larger isosceles right triangles, we can see that there is no bound on “thinness.” Since the Cayley graph of  $\mathbb{Z}^2$  is quasi-isometric to  $\mathbb{R}^2$ ,  $\mathbb{Z}^2$  is not hyperbolic. Indeed, it is a theorem that a hyperbolic group cannot contain a copy of  $\mathbb{Z}^2$ .

Now suppose that we are given a hyperbolic group  $G$ . What does the geometry tell us about the group? Here is a list of some consequences of hyperbolicity. We refer the reader to [3] and [7] for proofs and additional references.

1.  $G$  has a finite presentation.
2.  $G$  has a Dehn’s algorithm, hence a linear time solution to the Word Problem.
3. The Isomorphism Problem is solvable for the class of hyperbolic groups.
4. The centralizer of every element of  $G$  is virtually cyclic.
5.  $G$  has at most finitely many conjugacy classes of torsion elements.
6. For any finite set of elements  $g_1, \dots, g_k$  in  $G$ , there exists  $n > 0$  such that the set  $\{g_1^n \dots g_k^n\}$  generates a free subgroup of rank at most  $k$ .
7. For  $n$  sufficiently large,  $H^n(G; \mathbb{Q}) = 0$ .
8. If  $G$  is torsion-free, it has a finite  $K(G, 1)$ -space (i.e., a finite CW-complex with fundamental group  $G$  and contractible universal covering space).

The proofs of these properties are beyond the scope of this paper, but the conclusion should be clear: geometry can have strong implications for algebraic and combinatorial properties of a group.

## 4 Beyond Hyperbolicity

Classical hyperbolic geometry and Gromov’s generalization to  $\delta$ -hyperbolic spaces have provided powerful tools for studying hyperbolic groups. But this class of groups is very special. For example, any group containing a subgroup isomorphic to  $\mathbb{Z}^2$  cannot be hyperbolic (see property (4) above). In recent years, there has been much interest in generalizing some of these techniques to broader classes of groups. Gromov himself introduced a notion of “non-positive curvature” for geodesic metric

spaces, called CAT(0) spaces, and groups acting on these spaces have also been extensively studied. CAT(0) geometry, for example, played a major role in the recent work of Agol, Wise, and others leading to a proof of the Virtual Haken Conjecture, the last remaining piece in Thurston's program to classify three-manifolds. But this is a topic for another day.


Other approaches to generalizing the theory of hyperbolic groups involve looking at groups that act on hyperbolic spaces, but where the actions are not geometric; instead, they satisfy some weaker conditions. This includes, for example, “relatively hyperbolic groups,” “acylindrically hyperbolic groups,” and “hierarchically hyperbolic groups.” The first of these is modeled on fundamental groups of hyperbolic manifolds with cusps; the others are inspired by mapping class groups and their actions on curve complexes. Some very nice introductions to these topics can be found in [2, 16, 17].

My own work in this area has focused on a somewhat different approach to capturing hyperbolic behavior in more general spaces and groups. Let's begin with an example.

*Example 13* Let  $Z$  be the space obtained by gluing a circle and a torus together at a single point,

$$Z = S^1 \vee T^2.$$

Let  $\tilde{Z}$  be its universal cover. In  $\tilde{Z}$ , the inverse image of the torus consists of infinitely many copies of the Euclidean plane (which we refer to as “flats”) and emanating from each lattice point in each of these planes is a line segment which projects to the circle. The result is a tree-like configuration of planes and lines (see Fig. 6), which we will refer to as the “tree of flats.” Triangles lying in a single flat can



**Fig. 6** Tree of flats

be arbitrarily “fat” whereas triangles whose sides lie mostly along the vertical lines are “thin.” Thus, in  $\tilde{Z}$ , we have hyperbolic-like directions, and non-hyperbolic directions. Intuitively, the more we travel vertically, the more hyperbolic it feels.

We are interested in identifying geodesics in a metric space  $X$  that behave like geodesics in a hyperbolic space. A good way to encode such geodesics is by means of a boundary. In general, unbounded metric spaces do not come equipped with a boundary. For example, in the hyperbolic plane (or the Euclidean plane) one can travel forever in any direction. To create a boundary for such a space, we need to add a point for each “direction to infinity.” In the case of the hyperbolic or Euclidean plane, this space of directions forms a circle. Adding this circle to the plane “at infinity” compactifies the space.

It turns out that this idea generalizes nicely to any hyperbolic metric space. For a  $\delta$ -hyperbolic space  $X$ , we define the boundary as follows. A *ray* in  $X$  is an isometric embedding  $\alpha : [0, \infty) \rightarrow X$ . As a set, the boundary of  $X$  is defined to be

$$\partial X = \{\alpha \mid \alpha : [0, \infty) \rightarrow X \text{ is a ray}\} / \sim$$

where  $\alpha \sim \beta$  if  $\alpha$  and  $\beta$  remain bounded distance from each other. In the Euclidean plane, for example, two rays are equivalent if and only if they are parallel.

To topologize  $\partial X$ , think of two rays as representing nearby points in the boundary if they remain close to each other for a long time. More precisely, define a neighborhood  $N(\alpha, R)$  of a ray  $\alpha$  to be the set of rays  $\beta$  such that  $\beta(t)$  lies within  $2\delta$  of  $\alpha(t)$  for  $0 \leq t \leq R$ . As  $R$  increases, these neighborhoods get smaller and smaller, and together, they form a neighborhood basis for a topology on  $\partial X$ .

For example, the boundary of the hyperbolic plane  $\mathbb{H}^2$  is a circle while the boundary of an infinite tree is a Cantor set.

From the geometric group theory viewpoint, a key property of the boundary of a hyperbolic space is quasi-isometry invariance.

**Theorem 1** *Let  $f : X \rightarrow Y$  be a quasi-isometry between two hyperbolic metric spaces. Then  $f$  induces a homeomorphism  $\partial f : \partial X \cong \partial Y$ . In particular, a hyperbolic group  $G$  has a well-defined boundary, namely the boundary of a Cayley graph of  $G$ .*

These boundaries have many nice properties and applications. The boundary gives rise to a compactification of  $X$ ,  $\overline{X} = X \cup \partial X$ , and it provides a powerful tool for studying the dynamics of groups actions, rigidity theorems, geodesic flows, etc.

In the quest to extend the techniques of hyperbolic geometry to more general spaces and groups, it is natural to ask whether analogous boundaries can be defined in more general contexts. Certainly we can consider equivalence classes of geodesic rays in any geodesic metric space  $X$ . However, if  $X$  is not hyperbolic, many things can go wrong. In some cases, it is not even clear how to define a topology on this set as the neighborhoods described above need not satisfy the requirements for a neighborhood basis.

Moreover, even when there is a nice topology on  $\partial X$ , other fundamental properties of hyperbolic boundaries can fail to hold. Consider, for example, the boundary of the Euclidean plane. This boundary is a circle and each point on the boundary can be represented by a unique ray based at the origin. As observed above, it provides a compactification of the plane. That's the good news. Here is some bad news.

- Many isometries (in particular all translations) act trivially on the boundary.
- The only pairs of points on the boundary that can be joined by a bi-infinite geodesic are pairs of antipodal points.
- A quasi-isometry of the plane to itself need not extend to a map on the boundary. For example, the map  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  taking  $re^{i\theta} \mapsto re^{i(\theta+\ln(r))}$  is a quasi-isometry which twists each ray emanating from the origin into a spiral.

In short, many of the properties of hyperbolic boundaries that permit applications to dynamics, rigidity, etc. fail to hold for this boundary.

Most significantly from the point of view of geometric group theory, quasi-isometry invariance fails miserably for non-hyperbolic boundaries. There are examples of groups that act geometrically on two CAT(0) spaces (spaces of non-positive curvature) whose boundaries are not homeomorphic [11]. Thus, we don't have a well-defined notion of a boundary for these groups.

What goes wrong is the failure of the Morse property. A quasi-isometry  $f : X \rightarrow Y$  of hyperbolic spaces takes a geodesic ray in  $X$  to a quasi-geodesic ray in  $Y$ , that is, a quasi-isometric embedding of the half-line  $\mathbb{R}^+ = [0, \infty)$  into  $Y$ . The Morse property guarantees that this quasi-geodesic ray lies close to some geodesic ray and hence determines a well-defined point at infinity.

**Definition 5** A ray (or bi-infinite geodesic)  $\alpha$  in  $X$  is *Morse* if there exists a function  $N : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that for any  $(K, C)$ -quasi-geodesic  $\beta$  with endpoints on  $\alpha$ ,  $\beta$  lies in the  $N(K, C)$ -neighborhood of  $\alpha$ . The function  $N$  is called a *Morse gauge* for  $\alpha$  and we say that  $\alpha$  is  $N$ -Morse.

If  $X$  is hyperbolic, then there exists a Morse gauge  $N$  such that every ray in  $X$  is  $N$ -Morse. This property is the key to proving quasi-isometry invariance for boundaries of hyperbolic spaces, and it plays a key role in the proofs of many other properties of hyperbolic spaces as well. If  $X$  is not hyperbolic, the Morse property may fail for some (or perhaps all) rays in  $X$ . On the other hand, many non-hyperbolic spaces contain a large number of Morse rays. Consider, for example, our “tree of flats” described above. It can be shown that a ray that spends a uniformly bounded amount of time in any flat is Morse. We view Morse rays as “hyperbolic-like” directions in  $X$ . Indeed, it can be shown that these rays share many other nice properties with rays in hyperbolic space [4, 10]. For example, if two sides of a triangle are  $N$ -Morse, then the triangle is  $\delta$ -thin where  $\delta$  depends only on the Morse gauge  $N$ .

This brings us to the Morse boundary. The *Morse boundary*,  $\partial_M X$  can be defined for any proper geodesic metric space  $X$ . As a set, it consists of equivalence classes of Morse rays,

$$\partial_M X = \{\alpha \mid \alpha : [0, \infty) \rightarrow X \text{ is a Morse ray}\} / \sim$$

where the equivalence  $\sim$  is defined as before. The topology is more subtle. For a sequence of rays  $\{\alpha_i\}$  to converge to  $\alpha$  in this topology, they must not only converge pointwise, they must also be uniformly Morse, that is, there exists a Morse gauge  $N$  such that all of the  $\alpha_i$  are  $N$ -Morse.

This boundary was first introduced for CAT(0) spaces by myself and Harold Sultan in [4] and shown to be quasi-isometry invariant. This was then generalized to arbitrary proper geodesic metric spaces by Matt Cordes in [8].

**Theorem 2** *Let  $f : X \rightarrow Y$  be a quasi-isometry between two proper geodesic metric spaces. Then  $f$  induces a homeomorphism  $\partial f : \partial_M X \rightarrow \partial_M Y$ . In particular,  $\partial_M G$  is well-defined for any finitely generated group.*

*Example 14*

- (1) If  $X$  is hyperbolic, then all rays are  $N$ -Morse for some fixed  $N$ . So in this case,  $\partial_M X = \partial X$ , the usual hyperbolic boundary. For example,  $\partial_M \mathbb{H}^2$  is a circle.
- (2) For  $X = \mathbb{R}^2$  the Euclidean plane, there are no Morse rays at all, so  $\partial_M X = \emptyset$ .
- (3) Let  $\tilde{Z}$  be the “tree of flats” from Example 13. Then the Morse geodesics in  $\tilde{Z}$  are those that spend a uniformly bounded amount of time in any flat and the maximum time spent in a flat is determined by the Morse gauge. Thus, for a given Morse gauge  $N$ , the  $N$ -Morse rays emanating from a fixed basepoint  $z_0$  lie in a subspace quasi-isometric to a tree. It follows that these rays determine a Cantor set in the boundary and the Morse boundary of  $\tilde{Z}$  is the direct limit of these Cantor sets. Note that since  $\tilde{Z}$  is the universal cover of  $S^1 \vee T^2$ , the fundamental group  $\mathbb{Z} * \mathbb{Z}^2 = \pi_1(S^1 \vee T^2)$  acts geometrically on  $\tilde{Z}$ . Hence by the theorem above, the Morse boundary of the group  $\mathbb{Z} * \mathbb{Z}^2$  is homeomorphic to the Morse boundary of  $\tilde{Z}$ .

The Morse boundary was designed to capture hyperbolic-like behavior in non-hyperbolic metric spaces and to give a well-defined notion of a boundary for a finitely generated group  $G$ . When the Morse boundary is non-trivial, it provides a new tool for studying these spaces and groups. It can be used, for example, to study the dynamics of isometries [15] and to determine when two groups are quasi-isometric [5]. It can also be used to study geometric properties of subgroups  $H < G$  [10]. For a survey of recent results on Morse boundaries, see [9].

Geometric group theory is a broad and growing area of mathematics. This article is intended only as a snapshot of some themes that run through the field. I invite you to investigate further!

**Acknowledgements** R. Charney was partially supported by NSF grant DMS-1607616.

## References

1. B. Bowditch, *A Course on Geometric Group Theory*. MSJ Memoirs, vol. 16 (Mathematical Society of Japan, Tokyo, 2006)
2. B. Bowditch, Relatively hyperbolic groups. *Int. J. Algebra Comput.* **22**(3), 1250016 [66 pp.] (2012)
3. M. Bridson, A. Haefliger, *Metric Spaces of Non-positive Curvature*. Grundlehren der Mathematischen Wissenschaften, vol. 319 (Springer, Berlin, 1999)
4. R. Charney, H. Sultan, Contracting boundaries of CAT(0) spaces. *J. Topol.* **8**, 93–117 (2013)
5. R. Charney, M. Cordes, D. Murray, Quasi-mobius homeomorphisms of Morse boundaries (Jan 2018). arXiv:1801.05315
6. M. Clay, D. Margalit (eds.), *Office Hours with a Geometric Group Theorist* (Princeton University Press, Princeton, 2017)
7. M. Coornaert, T. Delzant, A. Papadopoulos, *Géométrie et théorie des groupes: Les groupes hyperboliques de Gromov*. Lecture Notes in Mathematics, vol. 1441 (Springer, Berlin, 1990)
8. M. Cordes, Morse boundaries of proper geodesic metric spaces. *Groups Geom. Dyn.* **11**(4), 1281–1306 (2017)
9. M. Cordes, A survey on Morse boundaries and stability, in *Beyond Hyperbolicity*. LMS Lecture Notes. arXiv:1704.07598 (to appear)
10. M. Cordes, D. Hume, Stability and the Morse boundary. *J. Lond. Math. Soc.* (2) **95**(3), 963–988 (2017)
11. C. Croke, B. Kleiner, Spaces with nonpositive curvature and their ideal boundaries. *Topology* **39**, 549–556 (2000)
12. M. Dehn, *Papers on Group Theory and Topology*. Translated from the German and with introductions and an appendix by J. Stillwell (Springer, New York, 1987)
13. E. Ghys, P. de la Harpe, *Sur les groupes hyperboliques d'après Mikhael Gromov* (Bern, 1988). Progress in Mathematics, vol. 83 (Birkhäuser, Boston, 1990)
14. M. Gromov, Hyperbolic groups, in *Essays in Group Theory*. Mathematical Sciences Research Institute Publications, vol. 8 (Springer, New York, 1987), pp. 75–263
15. D. Murray, Topology and dynamics of the contracting boundary of cocompact CAT(0) spaces. *Pac. J. Math.* arXiv:1509.09314 (to appear)
16. D. Osin, Acylindrically hyperbolic groups. *Trans. Am. Math. Soc.* **368**(2), 851–888 (2016)
17. A. Sisto, What is a hierarchically hyperbolic space? (July 2017). arXiv:1707.00053

# Generalized Iterated Wreath Products of Cyclic Groups and Rooted Trees Correspondence



Mee Seong Im and Angela Wu

**Abstract** Consider the generalized iterated wreath product  $\mathbb{Z}_{r_1} \wr \mathbb{Z}_{r_2} \wr \dots \wr \mathbb{Z}_{r_k}$  where  $r_i \in \mathbb{N}$ . We prove that the irreducible representations for this class of groups are indexed by a certain type of rooted trees. This provides a Bratteli diagram for the generalized iterated wreath product, a simple recursion formula for the number of irreducible representations, and a strategy to calculate the dimension of each irreducible representation. We calculate explicitly fast Fourier transforms (FFT) for this class of groups, giving the literature's fastest FFT upper bound estimate.

**Keywords** Iterated wreath products · Cyclic groups · Rooted trees · Irreducible representations · Fast Fourier transform · Bratteli diagrams

**AMS Subject Classification** Primary 20C99, 20E08; Secondary 65T50, 05E25

## 1 Introduction

Representations of groups appear naturally in nature, more often than groups themselves. They appear in the form of a linear representation, a permutation representation, and automorphisms of an algebra, a group, a variety or scheme, or a manifold. For example, one can study functions on the circle  $S^1$ , which could be thought of as a group under addition, which form representations of  $S^1$ . Such functions could also be thought of as periodic functions on the set  $\mathbb{R}$  of real numbers, and the decomposition of the space of functions on  $S^1$  is known as the theory of Fourier series. One can also study the additive group  $\mathbb{R}$  of real numbers acting on

---

M. S. Im (✉)

Department of Mathematical Sciences, United States Military Academy,  
West Point, NY, USA

A. Wu

Department of Mathematics, University of Chicago, Chicago, IL, USA  
e-mail: [wu@math.uchicago.edu](mailto:wu@math.uchicago.edu)