

LEARNING MADE EASY

3rd Edition



Algebra II

WORKBOOK

WITH
ONLINE
PRACTICE



for
dummies[®]
A Wiley Brand

Find FREE quizzes
for every chapter online

Master Algebra II with
hundreds of practice problems

Review key math
concepts

Mary Jane Sterling

Author of *Algebra I For Dummies*
and *Algebra II For Dummies*



This book comes with access to more content online.

Test your knowledge with a quiz
for every chapter!

Register your book or ebook at
www.dummies.com/go/getaccess.

Select your product, and then follow the prompts
to validate your purchase.

You'll receive an email with your PIN and instructions.



Algebra II Workbook

3rd Edition

by Mary Jane Sterling

for
dummies[®]
A Wiley Brand

Algebra II Workbook For Dummies®, 3rd Edition

Published by: **John Wiley & Sons, Inc.**, 111 River Street, Hoboken, NJ 07030-5774, www.wiley.com

Copyright © 2019 by John Wiley & Sons, Inc., Hoboken, New Jersey

Published simultaneously in Canada

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning or otherwise, except as permitted under Sections 107 or 108 of the 1976 United States Copyright Act, without the prior written permission of the Publisher. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008, or online at www.wiley.com/go/permissions.

Trademarks: Wiley, For Dummies, the Dummies Man logo, Dummies.com, Making Everything Easier, and related trade dress are trademarks or registered trademarks of John Wiley & Sons, Inc. and may not be used without written permission. All other trademarks are the property of their respective owners. John Wiley & Sons, Inc. is not associated with any product or vendor mentioned in this book.

LIMIT OF LIABILITY/DISCLAIMER OF WARRANTY: THE PUBLISHER AND THE AUTHOR MAKE NO REPRESENTATIONS OR WARRANTIES WITH RESPECT TO THE ACCURACY OR COMPLETENESS OF THE CONTENTS OF THIS WORK AND SPECIFICALLY DISCLAIM ALL WARRANTIES, INCLUDING WITHOUT LIMITATION WARRANTIES OF FITNESS FOR A PARTICULAR PURPOSE. NO WARRANTY MAY BE CREATED OR EXTENDED BY SALES OR PROMOTIONAL MATERIALS. THE ADVICE AND STRATEGIES CONTAINED HEREIN MAY NOT BE SUITABLE FOR EVERY SITUATION. THIS WORK IS SOLD WITH THE UNDERSTANDING THAT THE PUBLISHER IS NOT ENGAGED IN RENDERING LEGAL, ACCOUNTING, OR OTHER PROFESSIONAL SERVICES. IF PROFESSIONAL ASSISTANCE IS REQUIRED, THE SERVICES OF A COMPETENT PROFESSIONAL PERSON SHOULD BE SOUGHT. NEITHER THE PUBLISHER NOR THE AUTHOR SHALL BE LIABLE FOR DAMAGES ARISING HEREFROM. THE FACT THAT AN ORGANIZATION OR WEBSITE IS REFERRED TO IN THIS WORK AS A CITATION AND/OR A POTENTIAL SOURCE OF FURTHER INFORMATION DOES NOT MEAN THAT THE AUTHOR OR THE PUBLISHER ENDORSES THE INFORMATION THE ORGANIZATION OR WEBSITE MAY PROVIDE OR RECOMMENDATIONS IT MAY MAKE. FURTHER, READERS SHOULD BE AWARE THAT INTERNET WEBSITES LISTED IN THIS WORK MAY HAVE CHANGED OR DISAPPEARED BETWEEN WHEN THIS WORK WAS WRITTEN AND WHEN IT IS READ.

For general information on our other products and services, please contact our Customer Care Department within the U.S. at 877-762-2974, outside the U.S. at 317-572-3993, or fax 317-572-4002. For technical support, please visit <https://hub.wiley.com/community/support/dummies>.

Wiley publishes in a variety of print and electronic formats and by print-on-demand. Some material included with standard print versions of this book may not be included in e-books or in print-on-demand. If this book refers to media such as a CD or DVD that is not included in the version you purchased, you may download this material at <http://booksupport.wiley.com>. For more information about Wiley products, visit www.wiley.com.

Library of Congress Control Number: 2018962324

ISBN 978-1-119-54311-4 (pbk); ISBN 978-1-119-54313-8 (ebk); ISBN 978-1-119-54312-1 (ebk)

Manufactured in the United States of America

10 9 8 7 6 5 4 3 2 1

Contents at a Glance

Introduction	1
Part 1: Getting Started with Algebra II	5
CHAPTER 1: Going Beyond Beginning Algebra	7
CHAPTER 2: Handling Quadratic (and Quadratic-Like) Equations and Inequalities	21
CHAPTER 3: Rooting Out the Rational, the Radical, and the Negative	43
CHAPTER 4: Graphing for the Good Life	61
Part 2: Functions	89
CHAPTER 5: Formulating Functions	91
CHAPTER 6: Specializing in Quadratic Functions	107
CHAPTER 7: Plugging in Polynomials	119
CHAPTER 8: Acting Rationally with Functions	143
CHAPTER 9: Exposing Exponential and Logarithmic Functions	163
Part 3: Conics and Systems of Equations	189
CHAPTER 10: Any Way You Slice It: Conic Sections	191
CHAPTER 11: Solving Systems of Linear Equations	221
CHAPTER 12: Solving Systems of Nonlinear Equations and Inequalities	237
Part 4: Other Good Stuff: Lists, Arrays, and Imaginary Numbers	251
CHAPTER 13: Getting More Complex with Imaginary Numbers	253
CHAPTER 14: Getting Squared Away with Matrices	263
CHAPTER 15: Going Out of Sequence with Sequences and Series	279
CHAPTER 16: Everything You Ever Wanted to Know about Sets and Counting	293
Part 5: The Part of Tens	309
CHAPTER 17: Basic Graphs	311
CHAPTER 18: Ten Special Sequences and Their Sums	317
INDEX	323

Table of Contents

INTRODUCTION

About This Book	1
Foolish Assumptions	2
Icons Used in This Book	2
Beyond the Book	3
Where to Go from Here	3

PART 1: GETTING STARTED WITH ALGEBRA II. 5

CHAPTER 1: **Going Beyond Beginning Algebra**7

Good Citizenship: Following the Order of Operations and Other Properties.	7
Specializing in Products and FOIL	10
Variables on the Side: Solving Linear Equations	11
Dealing with Linear Absolute Value Equations	12
Greater Math Skills: Equalizing Linear Inequalities.	14
Answers to Problems on Going Beyond Beginning Algebra	16

CHAPTER 2: **Handling Quadratic (and Quadratic-Like) Equations and Inequalities**.21

Finding Reasonable Solutions with Radicals	22
UnFOILed Again! Successfully Factoring for Solutions.	23
Your Bag of Tricks: Factoring Multiple Ways	25
Keeping Your Act Together: Factoring by Grouping	26
Resorting to the Quadratic Formula	27
Solving Quadratics by Completing the Square	29
Working with Quadratic-Like Equations.	30
Checking Out Quadratic Inequalities	32
Answers to Problems on Quadratic (and Quadratic-Like) Equations and Inequalities	34

CHAPTER 3: **Rooting Out the Rational, the Radical, and the Negative** . . .43

Doing Away with Denominators with an LCD	44
Simplifying and Solving Proportions.	46
Wrangling with Radicals	48
Changing Negative Attitudes toward Negative Exponents	49
Divided Powers: Solving Equations with Fractional Exponents	51
Answers to Problems on Rooting Out the Rational, the Radical, and the Negative	53

CHAPTER 4: **Graphing for the Good Life**61

Coordinating Axes, Coordinates of Points, and Quadrants.	62
Crossing the Line: Using Intercepts and Symmetry to Graph.	64
Graphing Lines Using Slope-Intercept and Standard Forms.	67
Graphing Basic Polynomial Curves	69

Grappling with Radical and Absolute Value Functions.	71
Enter the Machines: Using a Graphing Calculator	73
Answers to Problems on Graphing for the Good Life	77

PART 2: FUNCTIONS 89

CHAPTER 5: Formulating Functions.	91
Evaluating Functions	91
Determining the Domain and Range of a Function	93
Recognizing Even, Odd, and One-to-One Functions.	94
Composing Functions and Simplifying the Difference Quotient.	96
Solving for Inverse Functions.	99
Answers to Problems on Formulating Functions	101
CHAPTER 6: Specializing in Quadratic Functions	107
Finding Intercepts and the Vertex of a Parabola.	108
Applying Quadratics to Real-Life Situations.	109
Graphing Parabolas.	111
Answers to Problems on Quadratic Functions	113
CHAPTER 7: Plugging in Polynomials	119
Finding Basic Polynomial Intercepts	120
Digging up More-Difficult Polynomial Roots with Factoring	122
Determining Where a Function Is Positive or Negative	123
Graphing Polynomials.	125
Possible Roots and Where to Find Them: The Rational Root Theorem and Descartes's Rule	127
Getting Real Results with Synthetic Division and the Remainder Theorem	130
Connecting the Factor Theorem with a Polynomial's Roots	132
Answers to Problems on Plugging in Polynomials	134
CHAPTER 8: Acting Rationally with Functions	143
Determining Domain and Intercepts of Rational Functions	144
Introducing Vertical and Horizontal Asymptotes	145
Getting a New Slant with Oblique Asymptotes	147
Removing Discontinuities.	148
Going the Limit: Limits at a Number and Infinity	149
Graphing Rational Functions	151
Answers to Problems on Rational Functions.	156
CHAPTER 9: Exposing Exponential and Logarithmic Functions	163
Evaluating e-Expressions and Powers of e.	164
Solving Exponential Equations.	165
Making Cents: Applying Compound Interest and Continuous Compounding	167
Checking out the Properties of Logarithms.	169
Presto-Chango: Expanding and Contracting Expressions with Log Functions.	171
Solving Logarithmic Equations.	173
They Ought to Be in Pictures: Graphing Exponential and Logarithmic Functions.	175
Answers to Problems on Exponential and Logarithmic Functions.	179

PART 3: CONICS AND SYSTEMS OF EQUATIONS	189
CHAPTER 10: Any Way You Slice It: Conic Sections	191
Putting Equations of Parabolas in Standard Form	192
Shaping Up: Determining the Focus and Directrix of a Parabola	194
Back to the Drawing Board: Sketching Parabolas	196
Writing the Equations of Circles and Ellipses in Standard Form	198
Determining Foci and Vertices of Ellipses	201
Rounding Out Your Sketches: Circles and Ellipses	203
Hyperbola: Standard Equations and Foci	205
Determining the Asymptotes and Intercepts of Hyperbolas	206
Sketching the Hyperbola	208
Answers to Problems on Conic Sections	211
CHAPTER 11: Solving Systems of Linear Equations	221
Solving Two Linear Equations Algebraically	221
Using Cramer's Rule to Defeat Unruly Fractions	223
A Third Variable: Upping the Systems to Three Linear Equations	225
A Line by Any Other Name: Writing Generalized Solution Rules	227
Decomposing Fractions Using Systems	229
Answers to Problems on Systems of Equations	231
CHAPTER 12: Solving Systems of Nonlinear Equations and Inequalities	237
Finding the Intersections of Lines and Parabolas	237
Crossing Curves: Finding the Intersections of Parabolas and Circles	239
Appealing to a Higher Power: Dealing with Exponential Systems	240
Solving Systems of Inequalities	242
Answers to Problems on Solving Systems of Nonlinear Equations and Inequalities	245
PART 4: OTHER GOOD STUFF: LISTS, ARRAYS, AND IMAGINARY NUMBERS	251
CHAPTER 13: Getting More Complex with Imaginary Numbers	253
Simplifying Powers of i	254
Not Quite Brain Surgery: Doing Operations on Complex Numbers	255
"Dividing" Complex Numbers with a Conjugate	256
Solving Equations with Complex Solutions	257
Answers to Problems on Imaginary Numbers	259
CHAPTER 14: Getting Squared Away with Matrices	263
Describing Dimensions and Types of Matrices	263
Adding, Subtracting, and Doing Scalar Multiplication on Matrices	265
Trying Times: Multiplying Matrices by Each Other	267
The Search for Identity: Finding Inverse Matrices	268
Using Matrices to Solve Systems of Equations	272
Answers to Problems on Matrices	274

CHAPTER 15: Going Out of Sequence with Sequences and Series	279
Writing the Terms of a Sequence	279
Differences and Multipliers: Working with Special Sequences	282
Backtracking: Constructing Recursively Defined Sequences	283
Using Summation Notation	284
Finding Sums with Special Series	286
Answers to Problems on Sequences and Series	289
CHAPTER 16: Everything You Ever Wanted to Know about Sets and Counting	293
Writing the Elements of a Set from Rules or Patterns	294
Get Together: Combining Sets with Unions, Intersections, and Complements	295
Multiplication Countdowns: Simplifying Factorial Expressions	297
Checking Your Options: Using the Multiplication Property	298
Counting on Permutations When Order Matters	300
Mixing It Up with Combinations	301
Raising Binomials to Powers: Investigating the Binomial Theorem	303
Answers to Problems on Sets and Counting	304
PART 5: THE PART OF TENS	309
CHAPTER 17: Basic Graphs	311
Putting Polynomials in Their Place	311
Lining Up Front and Center	312
Being Absolutely Sure with Absolute Value	313
Graphing Reciprocals of x and x^2	313
Rooting Out Square Root and Cube Root	314
Growing Exponentially with a Graph	315
Logging In on Logarithmic Graphing	316
CHAPTER 18: Ten Special Sequences and Their Sums	317
Adding as Easy as One, Two, Three	317
Summing Up the Squares	318
Finding the Sum of the Cubes	318
Not Being at Odds with Summing Odd Numbers	319
Evening Things Out by Adding Up Even Numbers	319
Adding Everything Arithmetic	319
Geometrically Speaking	320
Easing into a Sum for e	320
Signing In on the Sine	321
Powering Up on Powers of 2	322
Adding Up Fractions with Multiples for Denominators	322
INDEX	323

Introduction

Here you are, pencil in hand, ready to take on the challenges of working on Algebra II problems. How did you get here? Are you taking an Algebra II class and just not getting enough homework assigned? Or have you found a few gaps in the instruction and want to fill them in before you end up with a flood of questions? Maybe you've been away from algebra for a while and you want a review. Or perhaps you're getting ready to tackle another mathematics course, such as calculus. If you're looking for some good-natured, clear explanations on how to do some standard and challenging algebra problems, then you've come to the right place.

I hope you can find everything you need in this book to practice the concepts of Algebra II. You'll find some basic (to get you in the mood) and advanced algebra topics. But not all the basics are here — that's where Algebra I comes in. The topics that aren't here are referenced for your investigation or further study.

Calculus and other, more advanced math drive Algebra II. Algebra is the passport to studying calculus and trigonometry and number theory and geometry and all sorts of good mathematics and science. Algebra is basic, and the algebra here can help you grow in your skills and knowledge.

About This Book

You don't have to do the problems in this book in the order in which they're presented. You can go to the topics you want or need and refer back to earlier problems if necessary. You can jump back and forth and up and down, if so inclined (but please, not on the furniture). The organization allows you to move freely about and find what you need.

Use this book as a review or to supplement your study of Algebra II. Each section has a short explanation and an example or two — enough information to allow you to do the problems.

If you want more background or historical information on a topic, you can refer to the companion book, *Algebra II For Dummies*, where I go into more depth on what's involved with each type of problem. (If you need more-basic information, you can try *Algebra I For Dummies* and *Algebra I Workbook For Dummies*). In this workbook, I get to the point quickly but with enough detail to see you through. The answers to the problems, at the end of each chapter, provide even more step-by-step instruction.

Foolish Assumptions

You're interested in doing algebra problems. Is that a foolish thing for me to assume? No! Of course, you're interested and excited and, perhaps, just a slight bit tentative. No need to worry. In this book, I assume that you have a decent background in the basics of algebra and want to investigate further. If so, this is the place to be. I take those basic concepts and expand your horizons in the world of algebra.

Are you a bit rusty with your algebra skills? Then the worked-out solutions in this book will act as refreshers as you investigate the different topics. You may be preparing for a more advanced mathematics course such as trigonometry or calculus. Again, the material in this book will be helpful.

Or maybe it's just my first assumption that fits your situation: You're interested in doing algebra and couldn't pass up doing the problems in this book!

Icons Used in This Book

Throughout this book, I highlight some of the most important information with icons. Here's what the icons mean:



ALGEBRA
RULES

You can read the word *rules* as a noun or a verb. Sometimes it's hard to differentiate. But usually, in this book, *rules* is a noun. This icon marks a formula or theorem or law from algebra that pertains to the subject at hand. The rule applies at that moment and at any moment in algebra.



EXAMPLE

You see this icon when I present an example problem whose solution I walk you through step by step. You get a problem and a detailed answer.



REMEMBER

This icon refers back to information that I think you may already know. It needs to be pointed out or repeated so that the current explanation makes sense.



TIP

Tips show you a quick and easy way to do a problem. Try these tricks as you're solving problems.



WARNING

There are always things that are tricky or confusing or problems that just ask for an error to happen. This icon is there to alert you, hoping to help you avoid a mathematical pitfall.

Beyond the Book

No matter how well you understand the concepts of algebra, you'll likely come across a few questions where you don't have a clue. Be sure to check out the free Cheat Sheet for a handy guide that covers tips and tricks for answering Algebra II questions. To get this Cheat Sheet, simply go to www.dummies.com and enter "Algebra II Workbook For Dummies" in the Search box.

The online practice that comes free with this book contains over 300 questions so you can really hone your Algebra II skills! To gain access to the online practice, all you have to do is register. Just follow these simple steps:

1. **Register your book or ebook at Dummies.com to get your PIN. Go to www.dummies.com/go/getaccess.**
2. **Select your product from the dropdown list on that page.**
3. **Follow the prompts to validate your product, and then check your email for a confirmation message that includes your PIN and instructions for logging in.**

If you do not receive this email within two hours, please check your spam folder before contacting us through our Technical Support website at <http://support.wiley.com> or by phone at 877-762-2974.

Now you're ready to go! You can come back to the practice material as often as you want — simply log on with the username and password you created during your initial login. No need to enter the access code a second time.

Your registration is good for one year from the day you activate your PIN.

Where to Go from Here

You may become intrigued with a particular topic or particular type of problem. Where do you find more problems like those found in a section? Where do you find the historical background of a favorite algebra process? There are many resources out there, including a couple that I wrote:

- » Do you like the applications? Try *Math Word Problems For Dummies*.
- » Are you more interested in the business-type uses of algebra? Take a look at *Business Math For Dummies*.

If you're ready for another area of mathematics, look for a couple more of my titles: *Trigonometry For Dummies* and *Linear Algebra For Dummies*.

1

Getting Started with Algebra II

IN THIS PART . . .

Find order in the order of operations and relate algebraic properties to processes used when solving equations.

Solve linear equations and inequalities and rewrite absolute value equations before solving.

Take on radical equations, rational equations, and fractional exponents.

Use one or more factorization methods to ready equations for the multiplication property of zero.

Solve equations with the quadratic formula or complete the square.

Graph basic curves using intercepts and properties of functions.

IN THIS CHAPTER

- » Applying order of operations and algebraic properties
- » Using FOIL and other products
- » Solving linear and absolute value equations
- » Dealing with inequalities

Chapter 1

Going Beyond Beginning Algebra

The nice thing about the rules in algebra is that they apply no matter what level of mathematics or what area of math you're studying. Everyone follows the same rules, so you find a nice consistency and orderliness. In this chapter, I discuss and use the basic rules to prepare you for the topics that show up in Algebra II.

Good Citizenship: Following the Order of Operations and Other Properties

The *order of operations* in mathematics deals with what comes first (much like the chicken and the egg). When faced with multiple operations, this *order* tells you the proper course of action.



ALGEBRA
RULES

The *order of operations* states that you use the following sequence when simplifying algebraic expressions:

1. **Raise to powers or find roots.**
2. **Multiply or divide.**
3. **Add or subtract.**

Special groupings can override the normal order of operations. For instance, a^{b+c} asks you to add $b + c$ before raising a to the power, which is a sum. If groupings are a part of the expression, first perform whatever's in the grouping symbol. The most common grouping symbols are parentheses, (); brackets, []; braces, { }; fraction bars, —; absolute value bars, | |; and radical signs, $\sqrt{\quad}$.

If you find more than one operation from the same level, move from left to right performing those operations.



ALGEBRA
RULES

The commutative, associative, and distributive properties allow you to rewrite expressions and not change their value. So, what do these properties say? Great question! And here are the answers:



TIP

» **Commutative property of addition and multiplication:** $a + b = b + a$, and $a \cdot b = b \cdot a$; the order doesn't matter.

Rewrite subtraction problems as addition problems so you can use the commutative (and associative) property. In other words, think of $a - b$ as $a + (-b)$.

» **Associative property of addition and multiplication:** $a + (b + c) = (a + b) + c$, and $a(b \cdot c) = (a \cdot b)c$; the order is the same, but the grouping changes.

» **Distributive property of multiplication over addition (or subtraction):**
 $a(b + c) = a \cdot b + a \cdot c$, and $a(b - c) = a \cdot b - a \cdot c$.



ALGEBRA
RULES

The *multiplication property of zero* states that if the product of $a \cdot b = 0$, then either a or b (or both) must be equal to 0.



EXAMPLE

Q. Use the order of operations and other properties to simplify the expression

$$\frac{12\left(\left(\frac{5}{6} + \frac{3}{4}\right) - \frac{1}{6}\right)}{5\sqrt{2(3)^2 + 7}}$$

A. $\frac{17}{25}$. The big fraction bar is a grouping symbol, so you deal with the numerator and denominator separately. Use the commutative and associative properties to rearrange the fractions in the numerator; square the 3 under the radical in the denominator. Next, in the numerator, combine the fractions that have a common denominator; below the fraction bar, multiply the two numbers under the radical. Reduce the first fraction in the numerator; add the numbers under the radical. Distribute the 12 over the two fractions; take the square root in the denominator. Simplify the numerator and denominator.

Here's what the process looks like:

$$\frac{12\left(\left(\frac{5}{6} - \frac{1}{6}\right) + \frac{3}{4}\right)}{5\sqrt{2(9) + 7}} = \frac{12\left(\left(\frac{4}{6}\right) + \frac{3}{4}\right)}{5\sqrt{18 + 7}} = \frac{12\left(\frac{2}{3} + \frac{3}{4}\right)}{5\sqrt{25}} = \frac{8 + 9}{5(5)} = \frac{17}{25}$$

1 Simplify: $-3[4 - 2(3^2)]$

2 Simplify: $4^3(3^2 + 11)(7^4 - 7^4)(10^{10})$

3 Simplify: $153 + 187 - 153 + 270 - 471 - 270 + 471$

4 Simplify: $(6^2 + 4)\sqrt{5^2 - 3^2}$

5 Simplify: $\frac{\sqrt[3]{4(3^2 + 7)}}{\sqrt{121} - \sqrt{49}}$

6 Simplify: $7 + 3|2(8) - 5^2|$

(For info on absolute value, see the upcoming section, “Dealing with Linear Absolute Value Equations.”)

Specializing in Products and FOIL

Multiplying algebraic expressions together can be dandy and nice or downright gruesome. Taking advantage of patterns and processes makes the multiplication quicker, easier, and more accurate.

When multiplying two binomials together, you have to multiply the two terms in the first binomial times the two terms in the second binomial — you're actually *distributing* the first terms over the second. The FOIL acronym describes a way of multiplying those terms in an organized fashion, saving space and time. FOIL refers to multiplying the two First terms together, then the two Outer terms, then the two Inner terms, and finally the two Last terms. The Outer and Inner terms usually combine. Then you add the products together by combining like terms. So, if you have $(ax + b)(cx + d)$, you can do the multiplication of the terms, or FOIL, like so:

Terms	Product
First	$ax(cx)$
Outer	$ax(d)$
Inner	$b(cx)$
Last	$b(d)$

$$= acx^2 + adx + bcx + bd$$

$$= acx^2 + (ad + bc)x + bd$$

The following examples show some multiplication patterns to use when multiplying binomials (expressions with two terms).



EXAMPLE

Q. Find the square of the binomial:
 $(2x + 3)^2$

A. $4x^2 + 12x + 9$. When squaring a binomial, you square both terms and put twice the product of the two original terms between the squares: $(a + b)^2 = a^2 + 2ab + b^2$. So, $(2x + 3)^2 = (2x)^2 + 2(2x)(3) + 3^2 = 4x^2 + 12x + 9$.

Q. Multiply the two binomials together using FOIL: $(3x - 5)(4x + 7)$

A. $12x^2 + x - 35$. Find the products: First, $3x(4x) = 12x^2$, plus Outer, $3x(7) = 21x$, plus Inner, $(-5)(4x) = -20x$, plus Last, $(-5)(7) = -35$. So, the product of $(3x - 5)(4x + 7)$ is $(3x)(4x) + (3x)(7) + (-5)(4x) + (-5)(7) = 12x^2 + 21x - 20x - 35 = 12x^2 + x - 35$.

Q. Find the product of the binomial and the trinomial: $(2x + 7)(3x^2 + x - 5)$

A. $6x^3 + 23x^2 - 3x - 35$. Distribute the $2x$ over the terms in the trinomial, and then distribute the 7 over the same terms. Combine like terms to simplify. The product of $(2x + 7)(3x^2 + x - 5)$ is $2x(3x^2 + x - 5) + 7(3x^2 + x - 5) = 6x^3 + 2x^2 - 10x + 21x^2 + 7x - 35 = 6x^3 + 23x^2 - 3x - 35$.

7 Square the binomial: $(4x - 5)^2$

8 Multiply: $(5y - 6)(5y + 6)$

9 Multiply: $(8z - 3)(2z + 5)$

10 Multiply: $(2x - 5)(4x^2 + 10x + 25)$

Variables on the Side: Solving Linear Equations

A linear equation has the general format $ax + b = c$, where x is the variable and a , b , and c are constants. When you solve a linear equation, you're looking for the value that x takes on to make the linear equation a true statement. The general game plan for solving linear equations is to isolate the term with the variable on one side of the equation and then multiply or divide to find the solution.



EXAMPLE

Q. Solve for x in the equation

$$\frac{3(x + 7) - 5}{4} = 2x + 9.$$

A. $x = -4$. First, multiply each side by 4 to get rid of the fraction. Then distribute the 3 over the terms in the parentheses. Combine the like terms on the left. Next, you want all variable terms on one side of the equation, so subtract $8x$ and 16 from each side. Finally, divide each side by -5 .

$$\begin{aligned} \cancel{4} \cdot \frac{3(x + 7) - 5}{\cancel{4}} &= 4(2x + 9) \\ 3(x + 7) - 5 &= 8x + 36 \\ 3x + 21 - 5 &= 8x + 36 \\ 3x + 16 &= 8x + 36 \\ -5x &= 20 \\ x &= -4 \end{aligned}$$

11 Solve for x : $4x - 5 = 3(3x + 10)$

12 Solve for x : $5[9 - x(x + 8)] = 5x(1 - x)$

13 Solve for x : $\frac{x}{4} + \frac{x-6}{5} = \frac{x+2}{3}$

14 Solve for x : $(x-1) + 2(x-2) + 3(x-3) + 4(x-4) + 5(x-5) = 0$

Dealing with Linear Absolute Value Equations

The absolute value of a number is the number's distance from 0. The formal definition of *absolute value* is

$$|a| = \begin{cases} a, & a \geq 0 \\ -a, & a < 0 \end{cases}$$

In other words, the absolute value of a number is exactly that number unless the number is negative; when the number is negative, its absolute value is the opposite, or a positive. The absolute value of a number, then, is the number's value without a sign; it's never negative.

When solving linear absolute value equations, you have two possibilities: one that the quantity *inside* the absolute value bars is positive, and the other that it's negative. Because you have to consider both situations, you usually get two different answers when solving absolute value equations, one from each scenario. The two answers come from setting the quantity inside the absolute value bars first equal to a positive value and then equal to a negative value.

Before setting the quantity equal to the positive and negative values, first isolate the absolute value term on one side of the equation by adding or subtracting the other terms (if you have any) from each side of the equation.



TIP

If you find more than one absolute value expression in your problem, you have to get down and dirty — consider all the possibilities. A value inside absolute value bars can be either positive or negative, so look at all the different combinations: Both values within the bars are positive, or the first is positive and the second is negative, or the first is negative and the second positive, or both are negative. Whew!



EXAMPLE

Q. Solve for x in $|3x + 7| = 11$.

A. $x = \frac{4}{3}, -6, -6$. First rewrite the absolute value equation as two separate linear equations. In the first equation, assume that the $3x + 7$ is positive and set it equal to 11. In the second equation, also equal to 11, assume that the $(3x + 7)$ is negative. For that one, negate (multiply by -1) the whole binomial, and then solve the equation.

$$3x + 7 = 11$$

$$3x = 4$$

$$x = \frac{4}{3}$$

or

$$-(3x + 7) = 11$$

$$-3x - 7 = 11$$

$$-3x = 18$$

$$x = -6$$

15 Solve for x in $|2x + 1| = 83$.

16 Solve for x in $5|4 - 3x| + 6 = 11$.

17 Solve for x in $|x + 3| + |2x - 1| = 6$.

18 Solve for x in $3|4x - 5| + 10 = 7$.

Greater Math Skills: Equalizing Linear Inequalities

A linear inequality resembles a linear equation — except for the relationship between the terms. The basic form for a linear inequality is $ax + b > c$. When $\geq$ or $<$ or $\leq$ are in the statement, the methods used to solve the inequality stay the same. When the extra bar appears under the inequality symbol, it means “or equal to,” so you read $\leq$ as “is less than or equal to.”

The main time to watch out when solving inequalities is when you multiply or divide each side of the inequality by a negative number. When you do that — and yes, you’re allowed — you have to reverse the sense or the relationship. The inequality $>$ becomes $<$, and vice versa.



ALGEBRA
RULES

When solving *absolute value inequalities* (see the preceding section for more on absolute values), you first drop the absolute value bars. Then you apply one of two separate rules for absolute value inequalities, depending on which way the inequality symbol faces:

- » Solve $|ax + b| > c$ using the two inequalities $ax + b > c$ and $ax + b < -c$.
- » Solve $|ax + b| < c$ using the single compound inequality $-c < ax + b < c$.

Two ways of writing your answers are inequality notation and interval notation:



REMEMBER

- » **Inequality notation:** This notation is just what it says it is: If your answer is all x 's greater than or equal to 3, you write $x \geq 3$. To say that the answer is all x values between -5 and $+5$, including the -5 but not the positive 5, you write $-5 \leq x < 5$.
- » **Interval notation:** Some mathematicians prefer interval notation because it's so short and sweet. You simply list the starting and stopping points of the numbers you want to use. When you see this notation, you just have to recognize that you're discussing intervals of numbers (and not, for instance, the coordinates of a point). The rule is that you use a bracket, $[$ or $]$, when you want to include the number, and use a parenthesis, $($ or $)$, when you don't want to include the number. You always use a parenthesis with ∞ or $-\infty$. For example, to write $x \geq 3$ in interval notation, you use $[3, \infty)$. Writing $-5 \leq x < 5$ in interval notation, you have $[-5, 5)$.



EXAMPLE

- Q.** Solve the inequality $8x - 15 \geq 10x + 7$. Write the answer in both inequality and interval notation.
- A.** $x \leq -11$ and $(-\infty, -11]$. First subtract $10x$ from each side; then add 15 to each side. This step moves the variable terms to the left and the constants to the right: $-2x \geq 22$. Now divide each

side by -2 . Because you're dividing by a negative number, you need to reverse the inequality sign: $x \leq -11$. That's the answer in inequality notation. The solution is that x can be any number either equal to or smaller than -11 . In interval notation, you write this as $(-\infty, -11]$.

Q. Solve the inequality $|5 - 6x| < 7$. Write the answer in both inequality and interval notation.

A. $-\frac{1}{3} < x < 2$ and $\left(-\frac{1}{3}, 2\right)$. Rewrite the absolute value inequality as the inequality $-7 < 5 - 6x < 7$. Subtract 5 from each of the three sections of the inequality to put the variable term by itself: $-12 < -6x < 2$. Now divide each section by -6 , reversing the inequality symbols. Then, after you've gone to all the trouble of reversing the inequalities, rewrite the statement again with

the smaller number on the left to correspond to numbers on the number line. This step requires reversing the inequalities again. Here are the details:

$$-12 < -6x < 2$$

$$2 > x > -\frac{1}{3}$$

$$-\frac{1}{3} < x < 2$$

In interval notation, the answer is

$$\left(-\frac{1}{3}, 2\right).$$



WARNING

This answer looks very much like the coordinates of a point $\left(-\frac{1}{3}, 2\right)$. In instances like this, be very clear about what you're trying to convey with the interval notation.

19 Solve the inequality $5x + 7 \leq 22$.

20 Solve the inequality $8(y - 4) \geq 5(3y - 12)$.

21 Solve the inequality $|3x + 7| > 4$.

22 Solve the inequality $4|5 - 2x| + 3 < 7$.

Answers to Problems on Going Beyond Beginning Algebra

This section provides the answers (in bold) to the practice problems in this chapter.

- 1 Simplify: $-3[4 - 2(3^2)]$. The answer is **42**.

$$-3[4 - 2(9)] = -3[4 - 18] = -3[-14] = 42$$

- 2 Simplify: $4^3(3^2 + 11)(7^4 - 7^4)(10^{10})$. The answer is **0**.

The third factor is 0. This makes the whole product equal to 0. Remember, the *multiplication property of zero* says that if any factor in a product is equal to 0, then the entire product is equal to 0.

- 3 Simplify: $153 + 187 - 153 + 270 - 471 - 270 + 471$. The answer is **187**.

Use the associative and commutative properties to write the numbers and their opposites together:

$$187 + 153 - 153 + 270 - 270 - 471 + 471 = 187 + 0 + 0 + 0 = 187$$

- 4 Simplify: $(6^2 + 4)\sqrt{5^2 - 3^2}$. The answer is **160**.

$$(36 + 4)\sqrt{25 - 9} = (40)\sqrt{16} = (40)(4) = 160$$

- 5 Simplify: $\frac{\sqrt[3]{4(3^2 + 7)}}{\sqrt{121} - \sqrt{49}}$. The answer is **1**.

$$\frac{\sqrt[3]{4(9 + 7)}}{\sqrt{121} - \sqrt{49}} = \frac{\sqrt[3]{4(16)}}{11 - 7} = \frac{\sqrt[3]{64}}{4} = \frac{4}{4} = 1$$

- 6 Simplify: $7 + 3|2(8) - 5^2|$. The answer is **34**.

$$7 + 3|16 - 25| = 7 + 3|-9| = 7 + 3(9) = 7 + 27 = 34$$

- 7 Square the binomial: $(4x - 5)^2$. The answer is **$16x^2 - 40x + 25$** .

When squaring a binomial, the two terms are each squared, and the term between them is twice the product of the original terms.



WARNING

A common error in squaring binomials is to forget the middle term and just use the squares of the two terms in the binomial. If you tend to forget the middle term, you can avoid the error and get the correct answer through FOIL — $(4x - 5)(4x - 5) = 16x^2 - 20x - 20x + 25 = 16x^2 - 40x + 25$.

- 8 Multiply: $(5y - 6)(5y + 6)$. The answer is **$25y^2 - 36$** .

The product of two binomials that contain the sum and difference of the same two terms results in a binomial that's the difference between the squares of the terms.

- 9 Multiply: $(8z - 3)(2z + 5)$. The answer is $16z^2 + 34z - 15$.

Using FOIL, the first term in the answer is the product of the first two terms: $(8z)(2z)$. The middle term is the sum of the products of the Outer and Inner terms: $(8z)(5)$ and $(-3)(2z)$. The final term is the product of the two last terms: $(-3)(5)$.

- 10 Multiply: $(2x - 5)(4x^2 + 10x + 25)$. The answer is $8x^3 - 125$.

Distribute the first term in the binomial $(2x)$ over the terms in the trinomial, and then distribute the second term in the binomial (-5) over the terms in the trinomial. After that, combine like terms:

$$\begin{aligned} &= 2x(4x^2 + 10x + 25) - 5(4x^2 + 10x + 25) \\ &= 8x^3 + 20x^2 + 50x - 20x^2 - 50x - 125 \\ &= 8x^3 + 20x^2 - 20x^2 + 50x - 50x - 125 \\ &= 8x^3 - 125 \end{aligned}$$

- 11 Solve for x : $4x - 5 = 3(3x + 10)$. The answer is $x = -7$.

Distribute the terms on the right to get $4x - 5 = 9x + 30$. Subtract $9x$ and add 5 to each side, which gives you $-5x = 35$. Then divide each side by -5 : $x = -7$.

- 12 Solve for x : $5[9 - x(x + 8)] = 5x(1 - x)$. The answer is $x = 1$.

First distribute the x in the bracket on the left and the $5x$ on the right. Then you can distribute the 5 outside the bracket on the left to see what the individual terms are. The squared terms disappear when you add $5x^2$ to each side of the equation. Solve for x :

$$\begin{aligned} 5[9 - x^2 - 8x] &= 5x - 5x^2 \\ 45 - 5x^2 - 40x &= 5x - 5x^2 \\ 45 - 40x &= 5x \\ 45 &= 45x \\ 1 &= x \end{aligned}$$

- 13 Solve for x : $\frac{x}{4} + \frac{x-6}{5} = \frac{x+2}{3}$. The answer is $x = 16$.

First, multiply each fraction by 60 , the least common multiple. Then distribute and simplify:

$$\begin{aligned} 15\cancel{60}\left(\frac{x}{\cancel{4}_1}\right) + 12\cancel{60}\left(\frac{x-6}{\cancel{5}_1}\right) &= 20\cancel{60}\left(\frac{x+2}{\cancel{3}_1}\right) \\ 15x + 12(x-6) &= 20(x+2) \\ 15x + 12x - 72 &= 20x + 40 \\ 7x &= 112 \\ x &= 16 \end{aligned}$$

- 14 Solve for x : $(x-1) + 2(x-2) + 3(x-3) + 4(x-4) + 5(x-5) = 0$. The answer is $x = \frac{11}{3}$.

Distribute over the terms. Combine like terms and solve for x .

$$\begin{aligned} x - 1 + 2x - 4 + 3x - 9 + 4x - 16 + 5x - 25 &= 0 \\ 15x - 55 &= 0 \\ 15x &= 55 \\ x &= \frac{11}{3} \end{aligned}$$

- 15 Solve for x in $|2x + 1| = 83$. The answer is $x = 41, -42$.

First, let the value in the absolute value be positive, and solve $2x + 1 = 83$. You get $2x = 82$, or $x = 41$. Next, let the value in the absolute value be negative, and solve $-(2x + 1) = 83$. (If you multiply each side by -1 , you don't have to distribute the negative sign over two terms. Instead, you have $2x + 1 = -83$.) Solving this, you get $2x = -84$, or $x = -42$.

- 16 Solve for x in $5|4 - 3x| + 6 = 11$. The answer is $x = 1, \frac{5}{3}$.

First subtract 6 from each side, and then divide by 5. (You can't apply the rule for changing an absolute value equation into linear equations unless the absolute value is isolated on one side of the equation.) Now you have $|4 - 3x| = 1$. Letting the expression inside the absolute value be positive, you have $4 - 3x = 1$, which means $-3x = -3$ and $x = 1$. Now, if the expression inside the absolute value is negative, you make the expression positive by negating the whole thing: $-(4 - 3x) = 1$ gives you $4 - 3x = -1$, which means $-3x = -5$ and $x = \frac{5}{3}$.

- 17 Solve for x in $|x + 3| + |2x - 1| = 6$. The answer is $x = \frac{4}{3}, -2$.

You need four different equations to solve this problem. Consider that both absolute values may be positive; then that the first is positive and the second, negative; then that the first is negative and the second, positive; and last, that both are negative.



REMEMBER

Unfortunately, not every equation gives you an answer that really works. Perhaps no value of x can make the first absolute value negative and the second positive. An extraneous solution to an equation or inequality is a false or incorrect solution. It occurs when you change the original format of the equation to a form that's more easily solved. The extraneous solution may be a solution to the new form, but it doesn't work in the original. After you change the format, simply solve each equation produced, and then check each answer in the original equation.

Here are the situations, one at a time:

» **Positive/Positive:** $(x + 3) + (2x - 1) = 6$; $x + 3 + 2x - 1 = 6$; $3x + 2 = 6$; $3x = 4$; $x = \frac{4}{3}$. When you put this answer back into the original equation, it works — you get a true statement.

» **Positive/Negative:** $(x + 3) + -(2x - 1) = 6$; $x + 3 - 2x + 1 = 6$; $-x + 4 = 6$; $-x = 2$; $x = -2$. Putting this value back into the original equation, you get a true statement, so this answer is valid, too.

» **Negative/Positive:** $-(x + 3) + (2x - 1) = 6$; $-x - 3 + 2x - 1 = 6$; $x - 4 = 6$; $x = 10$. When you put this back into the original equation, you get $13 + 19 = 6$. That isn't true, so this solution is extraneous. It doesn't work.

» **Negative/Negative:** $-(x + 3) + -(2x - 1) = 6$; $-x - 3 - 2x + 1 = 6$; $-3x - 2 = 6$; $-3x = 8$; $x = -\frac{8}{3}$.

You get another extraneous solution. When you put this value for x back into the original equation, you get

$$\left| -\frac{8}{3} + 3 \right| + \left| 2\left(-\frac{8}{3} \right) - 1 \right| \stackrel{?}{=} 6$$

$$\left| \frac{1}{3} \right| + \left| -\frac{19}{3} \right| \stackrel{?}{=} 6$$

$$\frac{20}{3} \neq 6$$

Not so! This solution doesn't work.