

Mirza Hasanuzzaman · Kamrun Nahar  
Münir Öztürk *Editors*

# Ecophysiology, Abiotic Stress Responses and Utilization of Halophytes

 Springer

# Ecophysiology, Abiotic Stress Responses and Utilization of Halophytes

Mirza Hasanuzzaman • Kamrun Nahar  
Münir Öztürk  
Editors

# Ecophysiology, Abiotic Stress Responses and Utilization of Halophytes

 Springer

*Editors*

Mirza Hasanuzzaman  
Department of Agronomy  
Sher-e-Bangla Agricultural University  
Dhaka, Bangladesh

Kamrun Nahar  
Department of Agricultural Botany  
Sher-e-Bangla Agricultural University  
Dhaka, Bangladesh

Münir Öztürk  
Botany Department & Centre for  
Environmental Studies  
Ege University  
Izmir, Turkey

ISBN 978-981-13-3761-1      ISBN 978-981-13-3762-8 (eBook)  
<https://doi.org/10.1007/978-981-13-3762-8>

Library of Congress Control Number: 2019932838

© Springer Nature Singapore Pte Ltd. 2019, corrected publication 2019

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors, and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Singapore Pte Ltd.  
The registered company address is: 152 Beach Road, #21-01/04 Gateway East, Singapore 189721, Singapore

# Foreword



Professor Chedly Abdelly

Land salinization is a major limiting factor for conventional crop production and plant diversity in natural biotopes. Climate change, affecting several regions in the world and marked mainly by the rarefaction of precipitation and the increase in temperatures, exacerbates this situation. The nonrational use of chemical fertilizers and unconventional organic fertilizers as well as pesticides has amplified soil and water salinization and caused soil contamination by heavy metals and several other pollutants. Thus, the global annual losses in agricultural production from salt-affected land are worryingly spreading and worsening. This situation is of major concern considering (i) the current global climate change tendency, and (ii) that agricultural production should increase by 50 to 70% by 2050 to match the projected population growth to 9.3 billion. Increasing yields in saline cultivated lands, creating new productive ecosystems in marginal areas, and implementing appropriate cultural practices based on using unconventional water resources to irrigate crops are the only alternatives to achieve this goal. The majority of conventional crops known for their sensitivity to salinity and all the associated stresses it generates cannot represent the main and relevant actors in these approaches. Then, what are the plant species that can combine high salt tolerance with the ability to improve the physicochemical and biological properties of salinized agricultural lands and to valorize marginal

highly saline soils and unconventional water resources through a biomass production useful for human food, for animal feed, and/or for various industrial purposes? These are halophyte species. Halophytes, plants that survive to reproduce in environments where the salt concentration is around 200 mm NaCl or more, constitute about 1% of the world's flora. They have acquired through natural selection pressure the traits required to deal with extreme environmental constraints. This explains the growing interest paid on these plants. Thus, several research institutes in the world currently focus on halophyte species. In the same context, scientific production in terms of articles in specialized journals, books, as well as invention patents valorizing the data acquired on halophytes shows currently a substantial development in terms of quantity and quality.

The book entitled *Ecophysiology, Abiotic Stress Responses and Utilization of Halophytes* edited by Prof. Mirza Hasanuzzaman, Dr. Kamrun Nahar, and Prof. Münir Öztürk, eminent international experts in plant ecophysiology, fits perfectly into the approach dealing with the characterization and valorization of halophytes. The contributions presented in this book are written by internationally reputed scientists who report relevant and adroitly positioned data in relation to recent studies on halophytes. They once again confirm the numerous (fundamental, ecological, and economic) interests of halophytes and provide to potential users of these plants new and relevant pathways to valorize highly saline lands and unconventional water resources.

At the fundamental level, the data documented show the pertinence of using a multidisciplinary approach, bringing together ecophysiological, biochemical, and molecular tools to accurately characterize halophyte responses to environmental stresses, whether separated or combined at different developmental stages of their life cycle. Importantly, contributors show that there are specific mechanisms evolved by halophytes to cope with abiotic constraints and further strengthen the pertinence of using halophytes as the most appropriate “models” to investigate salt tolerance mechanisms in plants. At a more applied level, there are nice and concrete reports on the strong potential of halophytes to desalinate and improve the biological properties of salinized soils. Besides, cultivating halophytes seems to be the only biological approach to decontaminate heavy metal-affected marginal soils. It is known that phytodesalination and phytoremediation are prerequisites in the perspective of rehabilitating marginal lands and their transformation as new productive ecosystems. This book also further highlights the interests of halophytes as precious sources of bioactive substances which have a tremendous potential of utilizations in the agro-food, pharmaceutical, and cosmetic industries. Interestingly, there are also nice reports about the technical and economical feasibility of either cultivating halophytes using seawater or their utilization in integrated farming systems and for biomass production.

To summarize, as plants native to ecosystems with harsh environmental conditions, halophytes are unique and intriguing species. Investigating salt tolerance mechanisms in these plants is of incontestable academic interest, for instance, to generate markers that could be used for breeding programs. But also this generates

information with the practical outcome either for agriculture and industrial users or environmentalists and ecologists. As an ecophysiologicalist who spent more than 35 years working on this topic, I am convinced that the readers of this book will find precious and original data and reviews which will substantially improve their knowledge about salt tolerance traits identified in halophytes and their multiple interests.

Center of Biotechnology of Borj Cedria  
(CBBC), Hammam-Lif, Tunisia

Chedly Abdelly

National Agency for Scientific Research  
Promotion (ANPR), Tunis, Tunisia

# Preface

Halophytes are the plant species that can tolerate high salinity, and their lives are benefitted by salinity up to certain extent and can be adapted in a wide variety of saline habitats such as coastal regions, salt marshes, and inland deserts. There are approximately 1450 halophytic plant species documented (eHALOPH Database), while we consider a total number of plant species which is approximately 350,000 (The Plant List Database), the percentage of halophytes in total plant species is only 0.4%. Since saline habitats and evolutionary history of halophytes are diverse, they have many different adaptation mechanisms, and most of them have more than one adaptation mechanism to able to survive in saline places. Halophytes not only can tolerate salinity but also other stressors like drought, extreme temperature, waterlogging, and toxic metals/metalloids. Halophytes are many times exposed to heavy metal toxicity particularly in the mangroves ecosystem. They act as a major sink of a number of pollutants as they are connected to nearby drainage and rivers.

It is estimated that the world population would be 9.3 billion by 2050, and providing foods for the increasing population will be one of the greatest challenges for agricultural scientists because the arable lands are being decreased at the same time. It is reported that about 7% of the total earth land and 20% of the total arable *area* are *affected* by high *salt* contents. Increasing salinity is one of the negative impacts of climate change, and globally 7% of the total land and 20% of the total arable area are salty. Since halophytes possess special morphological, anatomical, and physiological features, those make them adaptive in saline habitat, learning from the halophytes is one of the viable approaches for understanding salt tolerance mechanisms. The adaptive features of halophytes may be strong genetic traits to be transferred to cultivated plant species toward the improvement and development of salt tolerant cultivars. Apart from the special adaptive features, halophytes have diverse uses such as food, fodder, fuel, and medicines. Therefore, the use of halophytes may be an effective way to enhance food security in the era of climate change. Numerous research works have been carried on different aspects of halophytes in regard to its ecophysiology, special adaptive features, tolerance mechanisms, specialized uses, and so on which made them fascinating plants. This book



presents a collection of 19 chapters written by 65 experts in the field of plant ecophysiology, environmental sciences, biotechnology, botany, and agriculture.

We, the editors, would like to give special thanks to the authors for their outstanding and timely work in producing such fine chapters. We are highly thankful to Ms. Khursheda Parvin for her valuable help in formatting and incorporating editorial changes in the manuscripts. We are highly thankful to Dr. Fumiko Yamaguchi, Senior Editor (Editor, Ecology and Animal Science), Springer, Japan, for her prompt responses during the acquisition. We are also thankful to Sivachandran Ramanan, Production Editor of this book, and all other editorial staffs for the precious help in formatting and incorporating editorial changes in the manuscripts. The editors and contributing authors hope that this book will include a practical update on our knowledge for ecophysiological features of halophytes and mechanisms of abiotic stress tolerance. Moreover, the potential uses of halophytes are also discussed. Graduate students (MS and PhD), university faculty members, and researchers working in the field of plant physiology, ecology, soil science, botany, and agriculture will be highly benefited from this book.

Dhaka, Bangladesh

Izmir, Turkey

Mirza Hasanuzzaman

Kamrun Nahar

Münir Öztürk

# Contents

<b>1</b>	<b>Halophyte Responses and Tolerance to Abiotic Stresses . . . . .</b>	<b>1</b>
	Ganesh Chandrakant Nikalje, Kushi Yadav, and Suprasanna Penna	
<b>2</b>	<b>Behavior of Halophytes and Their Tolerance Mechanism Under Different Abiotic Stresses . . . . .</b>	<b>25</b>
	Asha Sharma, Pooja, Annu Devi, Chanchal Garg, Anita Kumari, Anita Mann, and Ashwani Kumar	
<b>3</b>	<b>An Overview of the Germination Behavior of Halophytes and Their Role in Food Security . . . . .</b>	<b>39</b>
	Gül Nilhan Tuğ and Ahmet Emre Yaprak	
<b>4</b>	<b>Reactive Oxygen Species Production and Scavenging During Seed Germination of Halophytes . . . . .</b>	<b>63</b>
	Aysha Rasheed, Sarwat Ghulam Rasool, Bilquees Gul, M. Ajmal Khan, and Abdul Hameed	
<b>5</b>	<b>Halophyte Growth and Physiology Under Metal Toxicity . . . . .</b>	<b>83</b>
	Anita Kumari, Sunita Sheokand, Pooja, Ashwani Kumar, Anita Mann, Neeraj Kumar, Sarita Devi, Babita Rani, Arvind Kumar, and B. L. Meena	
<b>6</b>	<b>Oxidative Stress and Antioxidant Defence Under Metal Toxicity in Halophytes . . . . .</b>	<b>115</b>
	Anita Kumari, Vinod Goyal, and Sunita Sheokand	
<b>7</b>	<b>Molecular Mechanisms of Osmotic Stress Recovery in Extremophile Plants: What Can We Learn from Proteomics? . . . . .</b>	<b>157</b>
	Nèjia Farhat and Ahmed Debez	

<b>8</b>	<b>Halophytic Microbiome in Ameliorating the Stress</b> . . . . .	171
	Ahmad Mahmood, Ryota Kataoka, Oğuz Can Turgay, and Ahmet Emre Yaprak	
<b>9</b>	<b>Economic Utilization and Potential of Halophytes</b> . . . . .	195
	Sarita Devi, Ashwani Kumar, Sunder Singh Arya, Anita Kumari, Neeraj Kumar, Gurdev Chand, Anita Mann, Vinod Goyal, and Pooja	
<b>10</b>	<b>Halophytes: Prospective Plants for Future</b> . . . . .	221
	Ganesh Chandrakant Nikalje, Shelke Deepak Bhaskar, Kushi Yadav, and Suprasanna Penna	
<b>11</b>	<b>Sustainable Use of Halophytic Taxa as Food and Fodder: An Important Genetic Resource in Southwest Asia</b> . . . . .	235
	Münir Öztürk, Volkan Altay, and Aykut Güvensen	
<b>12</b>	<b>How Could Halophytes Provide a Sustainable Alternative to Achieve Food Security in Marginal Lands?</b> . . . . .	259
	Karim Ben Hamed and Luisa Custódio	
<b>13</b>	<b>Halophytes: The Plants of Therapeutic Medicine</b> . . . . .	271
	Sunder Singh Arya, Sarita Devi, Kirpa Ram, Sunil Kumar, Naveen Kumar, Anita Mann, Ashwani Kumar, and Gurdev Chand	
<b>14</b>	<b>Halophyte Species as a Source of Secondary Metabolites with Antioxidant Activity</b> . . . . .	289
	Milan Stanković, Dragana Jakovljević, Marija Stojadinov, and Zora Dajić Stevanović	
<b>15</b>	<b>Phytoamelioration of the Salt-Affected Soils Through Halophytes</b> . . . . .	313
	Arvind Kumar, Ashwani Kumar, Anita Mann, Geeta Devi, Hariom Sharma, Rachna Singh, and Satish Kumar Sanwal	
<b>16</b>	<b>Multidisciplinary Studies on a Pilot Coastal Desert Modular Farm Growing <i>Salicornia bigelovii</i> in United Arab Emirates</b> . . . . .	327
	Dionysia Angeliki Lyra, R. M. S. Al-Shihi, R. Nuqui, S. M. Robertson, A. Christiansen, S. Ramachandran, S. Ismail, and A. M. Al-Zaabi	
<b>17</b>	<b>Financial Analysis of Halophyte Cultivation in a Desert Environment Using Different Saline Water Resources for Irrigation</b> . . . . .	347
	S. M. Robertson, Dionysia Angeliki Lyra, J. Mateo-Sagasta, S. Ismail, and M. J. U. Akhtar	

**18 *Rhizophora* Biomass of Mangrove Swamp Forests and Its Utilization in Energy and Industrial Production: The Case of Malaysia . . . . . 365**  
I. Faridah-Hanum and A. Latiff

**19 Halophytic Plant Diversity of Duzdag Area in Nakhchivan Autonomous Republic: Azerbaijan . . . . . 383**  
Nuri Vagif Movsumova and Sayyara Jamshid Ibadullayeva

**Correction to: Ecophysiology, Abiotic Stress Responses and Utilization of Halophytes . . . . . C1**

# Contributors

**M. Ajmal Khan** Institute of Sustainable Halophyte Utilization (ISHU), University of Karachi, Karachi, Pakistan

**M. J. U. Akhtar** Pakistan Center for Advanced Studies in Agriculture and Food Security, University of Agriculture, Faisalabad, Pakistan

**R. M. S. Al-Shihi** Marine Environment Research Department (MERD), Ministry of Climate Change and Environment, Dubai, United Arab Emirates

**Volkan Altay** Biology Department, Faculty of Science & Arts, Hatay Mustafa Kemal University, Antakya, Hatay, Turkey

**A. M. Al-Zaabi** Marine Environment Research Department (MERD), Ministry of Climate Change and Environment, Dubai, United Arab Emirates

**Sunder Singh Arya** Department of Botany, Maharishi Dayanand University, Rohtak, Haryana, India

**Karim Ben Hamed** Laboratory of Extremophile Plants, Centre of Biotechnology of Borj Cedria, Hammam-Lif, Tunisia

**Shelke Deepak Bhaskar** Department of Botany, Amruteshwar Art's, Commerce and Science College, Pune, MS, India

**Gurdev Chand** Sher-e Kashmir University, Main Campus SKUAST-Jammu, Chatha, India

**Ganesh Chandrakant Nikalje** Department of Botany, R. K. Talreja College of Arts, Science and Commerce, Affiliated to University of Mumbai, Thane, India

**A. Christiansen** Department of Plant and Environmental Sciences (PLEN), University of Copenhagen, Copenhagen, Denmark

**Luisa Custódio** Faculty of Sciences and Technology, Centre of Marine Sciences, University of Algarve, Faro, Portugal

**Ahmed Debez** Laboratory of Extremophile Plants, Center of Biotechnology of Borj Cedria, (CBBC), Hammam-Lif, Tunisia

**Annu Devi** Department of Botany, Maharishi Dayanand University, Rohtak, Haryana, India

**Geeta Devi** Rashtriya Kisan Post Graduate College, Shamli, Uttar Pradesh, India

**Sarita Devi** Chaudhary Charan Singh, Haryana Agricultural University, Hisar, Haryana, India

**Nèjia Farhat** Laboratory of Extremophile Plants, Center of Biotechnology of Borj Cedria, (CBBC), Hammam-Lif, Tunisia

**I. Faridah-Hanum** Faculty of Forestry, Universiti Putra Malaysia, Serdang, Selangor, Malaysia

**Chanchal Garg** Department of Botany, Baba Masthnath University, Rohtak, Haryana, India

**Vinod Goyal** Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India

**Bilquees Gul** Institute of Sustainable Halophyte Utilization (ISHU), University of Karachi, Karachi, Pakistan

**Aykut Güvensen** Botany Department & Centre for Environmental Studies, Ege University, Izmir, Turkey

**Abdul Hameed** Institute of Sustainable Halophyte Utilization (ISHU), University of Karachi, Karachi, Pakistan

**Sayyara Jamshid Ibadullayeva** The Institute of Botany of ANAS, Baku, Azerbaijan

**S. Ismail** International Center for Biosaline Agriculture (ICBA), Dubai, United Arab Emirates

**Dragana Jakovljević** Department of Biology and Ecology, Faculty of Science, University of Kragujevac, Kragujevac, Serbia

**Ryota Kataoka** Department of Environmental Sciences, Faculty of Life & Environmental Sciences, University of Yamanashi, Yamanashi, Japan

**Arvind Kumar** ICAR – Central Soil Salinity Research Institute, Karnal, Haryana, India

**Ashwani Kumar** ICAR – Central Soil Salinity Research Institute, Karnal, Haryana, India

**Anita Kumari** Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India

**Naveen Kumar** Department of Botany, Maharishi Dayanand University, Rohtak, Haryana, India

**Neeraj Kumar** Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India

**Sunil Kumar** Department of Environmental Sciences, Maharishi Dayanand University, Rohtak, Haryana, India

**Kushi Yadav** Dr. B. Lal Institute of Biotechnology, Rajasthan University, Jaipur, India

**A. Latiff** Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Selangor, Malaysia

**Dionysia Angeliki Lyra** International Center for Biosaline Agriculture (ICBA), Dubai, United Arab Emirates

**Ahmad Mahmood** Department of Environmental Sciences, Faculty of Life & Environmental Sciences, University of Yamanashi, Yamanashi, Japan

**Anita Mann** ICAR – Central Soil Salinity Research Institute, Karnal, Haryana, India

**J. Mateo-Sagasta** International Water Management Institute, Colombo, Sri Lanka

**B. L. Meena** ICAR – Central Soil Salinity Research Institute, Karnal, Haryana, India

**Nuri Vagif Movsumova** The Institute of Botany of ANAS, Baku, Azerbaijan

**R. Nuqui** Marine Environment Research Department (MERD), Ministry of Climate Change and Environment, Dubai, United Arab Emirates

**Münir Öztürk** Botany Department & Centre for Environmental Studies, Ege University, Izmir, Turkey

**Suprasanna Penna** Nuclear Agriculture and Biotechnology Division, Bhabha Atomic Research Centre, Mumbai, India

**Pooja** Regional Center, ICAR – Sugarcane Breeding Institute, Karnal, Haryana, India

**S. Ramachandran** Biotechnology Department, BITS Pilani, Dubai, United Arab Emirates

**Kirpa Ram** Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India

**Babita Rani** Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India

**Aysha Rasheed** Institute of Sustainable Halophyte Utilization (ISHU), University of Karachi, Karachi, Pakistan

**Sarwat Ghulam Rasool** Institute of Sustainable Halophyte Utilization (ISHU), University of Karachi, Karachi, Pakistan

**S. M. Robertson** International Center for Biosaline Agriculture (ICBA), Dubai, United Arab Emirates

**Satish Kumar Sanwal** ICAR – Central Soil Salinity Research Institute, Karnal, Haryana, India

**Hariom Sharma** ICAR-Directorate of Rapeseed-Mustard Research, Bharatpur, Rajasthan, India

**Sunita Sheokand** Department of Botany and Plant Physiology, CCS Haryana Agricultural University, Hisar, India

**Asha Sharma** Department of Botany, Maharishi Dayanand University, Rohtak, Haryana, India

**Rachna Singh** ICAR – Central Soil Salinity Research Institute, Karnal, Haryana, India

**Milan Stanković** Department of Biology and Ecology, Faculty of Science, University of Kragujevac, Kragujevac, Serbia

**Zora Dajić Stevanović** Faculty of Agriculture, University of Belgrade, Belgrade, Serbia

**Marija Stojadinov** Department of Biology and Ecology, Faculty of Science, University of Kragujevac, Kragujevac, Serbia

**Gül Nilhan Tuğ** Department of Biology, Faculty of Science, Ankara University, Ankara, Turkey

**Oğuz Can Turgay** Department of Soil Science and Plant Nutrition, Faculty of Agriculture, Ankara University, Ankara, Turkey

**Ahmet Emre Yaprak** Department of Biology, Faculty of Science, Ankara University, Ankara, Turkey



## About the Editors

**Dr. Mizra Hasanuzzaman** is a Professor of Agronomy at Sher-e-Bangla Agricultural University in Dhaka, Bangladesh. He completed his Ph.D. on ‘Plant Stress Physiology and Antioxidant Metabolism’ at Ehime University, Japan with a scholarship from the Japanese government. Later, he completed his postdoctoral research at the Center of Molecular Biosciences, University of the Ryukyus, Japan, as a recipient of a Japan Society for the Promotion of Science (JSPS) postdoctoral fellowship. He was also awarded the Australian Government’s Endeavour Research Fellowship for postdoctoral research as an Adjunct Senior Researcher at the University of Tasmania, Australia. Dr. Hasanuzzaman’s current work is focused on the physiological and molecular mechanisms of environmental stress tolerance (salinity, drought, flood, and heavy metals/metalloids). He has published over 80 articles in peer-reviewed journals, edited six books and written 30 book chapters on important aspects of plant physiology, plant stress tolerance, and crop production. He is an editor and reviewer for more than 50 peer-reviewed international journals and was a recipient of the ‘Publons Peer Review Award 2017.’ He has been honored by various authorities for to his outstanding contributions to research and education, and received the World Academy of Science Young Scientist Award (2014).

**Dr. Kamrun Nahar** is an Associate Professor, Department of Agricultural Botany at Sher-e-Bangla Agricultural University, Dhaka, Bangladesh. She completed her Ph.D. on ‘Environmental Stress Physiology of Plants’ at Ehime University, Japan on a Japanese Government (MEXT) Scholarship in 2016. Dr. Nahar has been involved in research with field crops emphasizing stress physiology since 2006. She has completed several research projects funded by the Sher-e-Bangla Agricultural University Research System and the Ministry of Science and Technology (Bangladesh). Dr. Nahar has published numerous articles in peer-reviewed journals and books, including 50 articles and chapters related to plant physiology and environmental stresses with Springer, Elsevier, CRC Press, Wiley, etc. She is involved in editorial activities and serves as a reviewer for international journals. Further, she is an active member of over 20 professional societies.

**Dr. Münir Öztürk (Ph.D.)** completed his M.Sc. degree at Jammu & Kashmir University, Kashmir, and his Ph.D. & D.Sc. at Ege University Izmir, Turkey. Over the last 50 years, he has served in different positions at Ege University. Dr. Öztürk was elected as the “Vice President of the Islamic World Academy of Sciences” in 2017, and has received fellowships from the Alexander von Humboldt Society, Japanese Society for Promotion of Science, and National Science Foundation (USA). He is the Founding Director of the Centre for Environmental Studies at Ege University and has worked as a Consultant Fellow, Faculty of Forestry, Universiti Putra Malaysia, Malaysia and as Distinguished Visiting Scientist, ICCBS, Karachi University, Pakistan. His main fields of scientific interest are Plant Eco-Physiology; Medicinal and Aromatic Plants, Conservation of Plant Diversity; Biosaline Agriculture and Crops; Pollution and Biomonitoring. He has published 40 books, 50 book chapters, and more than 200 papers in prominent journals.

# Chapter 1

## Halophyte Responses and Tolerance to Abiotic Stresses



Ganesh Chandrakant Nikalje, Kushi Yadav, and Suprasanna Penna

### Contents

1.1	Introduction .....	2
1.1.1	Flooding .....	3
1.1.2	Heavy Metals .....	5
1.1.3	Salinity .....	7
1.1.4	Combined Stress [Salt + Drought] .....	9
1.1.5	Combined Stress [Salinity + Flood] .....	10
1.1.6	Combined Stress [Salt + Metal] .....	11
1.1.7	Combined Stress [Salt + Nutrient Deficiency] .....	12
1.1.8	Combined Stress [Salinity + Nitrogen Deficiency] .....	12
1.1.9	Combined Stress [Salt + Phosphorous Deficiency] .....	13
1.2	Cross Tolerance .....	14
1.3	Stress Memory .....	14
1.4	Conclusions and Future Prospective .....	15
	References .....	17

**Abstract** Different anthropogenic activities result in the contamination and degradation of the agricultural ecosystem. Improper disposal of industrial waste, use of excess chemical fertilizers, and mining are major sources of soil contamination. These adverse conditions exert a negative effect on crop growth and yield, while a group of plants, known as halophytes, exhibit greater tolerance. These plants are native to such adverse environments and can withstand different abiotic stresses such as salinity, drought, toxic metal stress, and hypoxia. Halophytes grow luxuriantly in saline soils, which make them suitable for saline agriculture. In addition, they are a

---

G. C. Nikalje  
Department of Botany, R. K. Talreja College of Arts, Science and Commerce,  
Affiliated to University of Mumbai, Thane, India

K. Yadav  
Dr. B. Lal Institute of Biotechnology, Rajasthan University, Jaipur, India

S. Penna (✉)  
Nuclear Agriculture and Biotechnology Division, Bhabha Atomic Research Centre, Mumbai,  
India  
e-mail: [prasanna@barc.gov.in](mailto:prasanna@barc.gov.in)

good source of salt-responsive genes and value-added products. Many halophytes show common biochemical and physiochemical responses to salt stress, whereas under multiple stresses, different mechanisms operate. The accumulation of osmolytes such as proline, selectivity in K:N, the exclusion of sodium and vacuolar compartmentalization, the induction of antioxidant molecules (enzymatic and non-enzymatic) are the most common features of halophytic adaptation to stress. The comparative study of halophytes and glycophytes revealed that the former are well equipped with cross-tolerance mechanism and are well prepared before stress imposition. It is also reported that pretreatment/priming with salinity or other stresses in early developmental stage of halophytes improves their salt tolerance at later stage. This observation suggests that halophytes might have stress memory, which helps them to respond better to stress conditions. In this article, we present a current perspective of the general tolerance mechanism and the responses of halophytes to different abiotic stresses such as salt, drought, toxic metal, and combination of these. Understanding the mechanism of such abiotic stresses alone and in combination will help to identify potential halophytes for re-vegetation or possible breeding for redevelopment of salt-affected agricultural lands.

**Keywords** Halophytes · Abiotic stress · Multiple stress · Salinity · Drought · Toxic metals

## 1.1 Introduction

Of the world's total land area, 6% is plagued by excess salinity, mainly in arid and semi-arid areas (Flowers et al. 2010). In agriculture, regional salinity problems are increasing owing to low precipitation, high evaporation, naturally saline soil, poor waste management, and saline water irrigation (Hamed et al. 2013). Usually, plants' response to salinity stress alone differs from their natural biotope behavior in comparison with combined stress (Mittler 2006). It was suggested that plants enhance the negative effect of a first stress in the presence of a second stress (Mittler 2006). Hence, increasing plants' tolerance toward stresses requires multiple stress research (Mittler and Blumwald 2010). The salinity tolerance of halophytes predominantly depends upon uptake control and compartmentalization of  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Cl}^-$  (Flowers and Colmer 2008; Shabala and Mackay 2011). Halophytes survive not only in a saline habitat, but also in drought, heat, and an alkaline habitat (Hamed et al. 2013). Along with these stresses, halophytes are also exposed to various biotic stress such as pathogen attack and abiotic stress, for example, toxic pollutants including pesticides, noxious gases ( $\text{SO}_2$ ,  $\text{NO}_x$ ,  $\text{O}_3$ ), soil acidification, excessive use of fertilizers, heavy metal and intensified UV-B irradiation (Flowers and Colmer 2008). When combined abiotic stress affects halophytes, they usually show tolerance against combined stress (Hamed et al. 2013). Combined stress tolerance is a complex phenomenon to explain because of multitude of events that take place at various stages of plant development.

Salt-tolerant plants with a phenomenal ability of completing their life cycle under saline conditions are known as halophytes (Flowers and Colmer 2008; Grigore et al. 2014). Development of morphological, anatomical, and physiological characteristics during evolution has enabled halophytes capable of growing under high salt conditions (Flowers and Colmer 2008; Grigore et al. 2014). Because of the wide range of stress tolerance environments, halophytes cannot be classified easily, but a few definitions of halophytes are available (Huchzermeyer and Flowers 2013). They are defined as plants capable of accomplishing their life cycle in a living environment with a salt concentration greater than 0.5% NaCl (Xu et al. 2016). Another definition is that they are exposed to salt stress and able to survive for at least a life cycle in an environment that is lethal to a large population of plants (Xu et al. 2016). Halophytes classified on the basis of salt requirement are obligate halophytes, facultative halophytes, and habitat-indifferent halophytes (Braun-Blanquet et al. 1933; Cushman 2001). Obligatory halophytes require salt for their growth, but facultative halophytes can grow in salt-devoid soil in a similar manner to habitat-indifferent halophytes, but differ because they like to grow in salt-devoid soil, but they are also able to survive under saline conditions (Braun-Blanquet et al. 1933). Another classification divides halophytes into reversible extremophytes, irreversible extremophytes, and mesophytes (Grigore et al. 2014). Extremo-halophytes are able to grow massively in a saline environment (Grigore et al. 2014). They are further subclassified on the basis of their habitat, which could be reversible or irreversible (Mishra and Tanna 2017). Modern omics techniques have enabled the understanding of salt tolerance mechanisms in these plants (Xu et al. 2016). With more insights into salt tolerance mechanisms, it has become apparent that there is more to tolerance mechanisms than mere salt tolerance (Xu et al. 2016). In this chapter, we present an account of stress tolerance behavior of halophytes under different abiotic stress conditions and in a combined stress regime. The main attention is given to abiotic stress factors, such as drought, salinity, flood, heavy metal and nutrient deficiency, which are usually lethal to plants other than halophytes.

### ***1.1.1 Flooding***

There are potential conditions of submergence of plants in flood water (Armstrong 1980). Complete submergence of plants results in to reduced O<sub>2</sub> supply to plants because diffusion of gases in water is 10,000 times slower in comparison with air (Armstrong 1980). Under dark conditions, O<sub>2</sub> is deficient, which can inhibit tissue respiration, which finally becomes anoxic (Gibbs and Greenway 2003). Internal O<sub>2</sub> level everyday based on CO<sub>2</sub>, O<sub>2</sub>, and incident light at the site of submergence (Pedersen et al. 2004; Winkel et al. 2013). The amount of CO<sub>2</sub> uptake by shoots for photosynthesis is also reduced because of submergence, mainly under circumstances where there is high resistance across the leaf cuticle (Mommer et al. 2005). In succulent halophytes, internal O<sub>2</sub> production is mainly based on O<sub>2</sub> production during daytime photosynthesis and anoxic when it is dark at night (Pedersen et al. 2006;

Colmer et al. 2013). In association with other problems of exchange of gases under submergence, halophytes also have to maintain an osmotic potential difference between tissues and floodwater, which could lead to swelling of tissues (Flowers and Colmer 2008). In that situation, to constrain such excessive swelling, a strong epidermis may be required, which greatly reduces the rate of gaseous exchange, already diminished because of the gas diffusion pathway and the relatively low surface area to volume (SA/V) ratio of succulent stems (Konnerup et al. 2015). Salicornioideae stem succulents habitually occupy saline, often flooded, habitats around world (Wilson et al. 1993; Shepherd and Wilson 2007). Perennial halophytes, stem succulent species largely endemic to Australia, contain the genus *Tecticornia* (Wilson et al. 1993; Shepherd and Wilson 2007). At ephemeral inland salt lakes, where the environment is distinguished by high soil salinity and episodic flooding punctuated by durations of drought, many *Tecticornia* species are known to grow (Wilson et al. 1993; Shepherd and Wilson 2007). Species zonation was found on the margins of ephemeral salt lakes in Australia in *Tecticornia* communities (English and Colmer 2011, 2013). Usually, *Tecticornia* field distribution has a different tolerance of flood, salinity, and drought, but an elevation in particular plant establishment may depend on the ability of submergence stress tolerance at low marsh conditions (Konnerup et al. 2015).

The SA/V ratio differs among species of succulent halophytes. Plants with thicker stems are more tolerant than others, explained by the internal O<sub>2</sub> dynamics of plants under submergence stress (Colmer et al. 2013). The major barrier is the supply of oxygen to stem tissue under darkness and submergence in the epidermis with the *T. medusa* stem being hypoxic throughout (Konnerup et al. 2013). Oxygen production in photosynthesis builds up owing to the low permeability of the epidermis; for example, *T. medusa* has a less permeable epidermis, but gas films on *T. auriculata* contribute to obtaining a better gas exchange initially after submergence stress, compared with *T. medusa*, in which gas films are absent (Pedersen et al. 2006). Therefore, low epidermis permeability and a low S/A ratio result in different oxygen patterns when there is a manipulation in the presence of light and submergence (Pedersen et al. 2006). Built-up respiratory carbon dioxide does not occur to the same extent in adequate species as they have thin leaves and a more penetrable epidermis without cuticles, which allows them to exchange gases with the surrounding water (Armstrong 1980; Colmer 2003). In waterlogged or submerged plants, the aeration of roots by O<sub>2</sub> diffusion is uncommon (Pedersen et al. 2006). Stem-succulent halophyte species experience problems during the night under submergence stress owing to O<sub>2</sub> depletion (Flowers and Colmer 2008). The O<sub>2</sub> depletion occurs because of the great fluctuation, as during the day time stem-succulent halophytes are voluminous and have low epidermal permeability (Konnerup et al. 2015). The survival of stem-succulent halophytes under submergence stress is associated with carbohydrate conversion, resistant osmotic stress, and the organized production of new tissue after the flood is over (Bailey-Serres and Voesenek 2008). To assist the survival of the plant, either in an environment susceptible to prolonged deep floods or during a short transient flood, the plant resumes better growth after the

flood water has receded, which is considered a quiescence response (Bailey-Serres and Voeselek 2008).

### 1.1.2 Heavy Metals

In the earth's crust, heavy metals are present as a natural component (Lutts and Lefèvre 2015). However, anthropogenic activities have led to the acceleration of heavy metal accumulation in the environment (Han et al. 2013). Various industrial and agricultural activities, for example, mining, smelting, burning of fossil fuel, and the excessive use of fertilizers, pesticides, fungi, and sewage sludge in agricultural practices are responsible for rapidly increasing heavy metal pollutants such as Cu, Cd, Pb, and Zn (Ernst and Nelissen 2000; D'Amore et al. 2005). These pollutants not only pose a serious threat to the environment, but also affect the health of living beings and disrupt the food chain (Sirguey and Ouvrard 2013). In the rhizosphere, through microbial activity, many organic pollutants can be converted into harmless compounds, but heavy metals cannot degrade and hence remain in the soil for long periods of time (Batty and Dolan 2013). These toxic compounds affect various plant species by inhibiting their growth, influencing the water status, photosynthesis, and inducing over-synthesis of reactive oxygen species (ROS), which is responsible for various physiological disorders that finally leads to plant death (Vázquez et al. 1994). There are some exceptional plant species known as hyper-accumulators. These hyper-accumulators display a higher shoot concentration of heavy metals than roots and these have the ability to absorb and also to translocate heavy metal (Vázquez et al. 1994; Yang et al. 2004).

Overproduction of ROS leads to oxidative stress (Demidchik et al. 2010). This oxidative stress is the major cause of oxidative damage to proteins, nucleic acids, and lipid peroxidation, which may also lead to programmed cell death (PCD) (Demidchik et al. 2010; Nikalje et al. 2017a, b, c; Demidchik 2015). Transition metals in response to oxidative stress generate hydroxyl radicals (Zepeda-Jazo et al. 2011; Rodrigo-Moreno et al. 2013), which activate plasma membrane  $\text{Ca}^{2+}$  and  $\text{K}^{+}$  channels, and lead to  $\text{Ca}^{2+}$  influx and  $\text{K}^{+}$  efflux (Demidchik 2015). Halophytes need proper regulation of Na and K concentrations inside their cells, exhibiting a defense response to hydroxyl radicals (Flowers and Muscolo 2015). In *Zygophyllum fabago*, the significance of K relocation in cells of photosynthetically active tissues in response to heavy metal toxicities were studied (Lefèvre et al. 2014). The accumulation of reduced form of transition metals such as Cu, Mn, and Fe is favored by anoxic conditions in some cases (Rodrigo-Moreno et al. 2013). In apoplast and cytosol through the Fenton reaction, transition metals in the presence of  $\text{H}_2\text{O}_2$  catalyze the production of hydroxyl radicals (Rodrigo-Moreno et al. 2013). Essential components of the plant antioxidant defense systems are Cu and Fe, which are part of superoxide dismutase (SOD) (Bose et al. 2014). When Cu and Fe become over-absorbed, they may result in metabolic disturbance such as loss of chloroplast integrity, remodeled plastid membrane, and also inhibits photosynthetic electron transport (Bose et al. 2014). Further ROS overproduction is generated owing to the

disruption of electron transport (Bose et al. 2014). Cu and Zn toxicities are important factors that result in secondary oxidative stress, but Cu and Zn are not involved directly in ROS synthesis (Lutts and Lefèvre 2015). Glutathione, ascorbate, and  $\alpha$ -tocopherol are some of the antioxidants present in plants (Bose et al. 2014; Ozgur et al. 2013). Among antioxidant enzymes, Catalase and peroxidase remove  $H_2O_2$  and, SODs catalyze the disproportion of superoxide radicals to  $O_2$  and  $H_2O_2$  (Bose et al. 2014; Ozgur et al. 2013). The greater constitutive antioxidant defense activities of halophytes play an important role in heavy metal toxicity, salinity and other stress (Wang et al. 2014).

In response to heavy metal and metalloids such as Cd, Ni and As, *Salicornia brachiata* is able to up-regulate catalase (CAT) and SOD activities (Lefèvre et al. 2010). The expression of the *CAT1*, *GST*, and *PrxQ* genes responsible for enzymes activity is increased by cadmium (Cong et al. 2013). In shoots of *Suaeda salsa* in response to Zn, Pb or a combination of both metals, only CAT gene expression level increased significantly in contrast to activities of antioxidant enzymes SOD, glutathione peroxidase (GPxS), and catalase increased in response to Zn and Zn+Pb (Bose et al. 2014; Jithesh et al. 2006). SOD activities play a significant role in the protection of halophytes from extreme environmental changes (Bose et al. 2014; Jithesh et al. 2006).

Glutathione by phytochelatin synthase enzymatically synthesizes phytochelatin and low-molecular-weight polypeptide with general structure  $(\gamma\text{Glu-Cys})_n\text{Gly 'n'}$  varying from 2 to 11 (Lutts and Lefèvre 2015). These polypeptides help to reduce damage under heavy metal stress conditions (Lutts and Lefèvre 2015). In *Avicennia germinans*, on exposure to a polluted environment, shows a high level of tolerance due to the overexpression of the *AvPCS* in response to high concentration of  $Cu^{2+}$  and  $Cd^{2+}$  within few hours (Tennstedt et al. 2008). This over-expression triggers a tolerance mechanism during long-term exposure, and remains transient (Baena-González et al. 2007). Proteins are found in almost all plants, animals, and microbes, that bind to metal and microtubules rich in cysteine (Cobbett 2003). According to the location and distribution of cysteine residue, plant microtubules are divided into four types (Cobbett 2003). Microtubules bind metals efficiently, which leads to metal tolerance. In *Bruguiera gymnorhiza*, Zn, Cu, and Pb up-regulate genes encoding type 2 microtubules (*MT2*) (Lutts and Lefèvre 2015). Such results were also obtained in *Avicennia marina* and transgenic *E. coli* LB4 (Huang and Wang 2009). Expression of gene coding *MT2* has a differing impact according to the various elements (Huang and Wang 2010). As in *Prosopis juliflora*, Cu and Cd stress cannot change the expression of *PjMT2*, but Zn up-regulates the expression of *PjMT2* (Usha et al. 2009). Similarly, in *Salicornia brachiata*, expression of *SbMT2* is unaffected by Pb and is up-regulated in the presence of Zn and Cu (Chaturvedi et al. 2012).

Halophytes lead to osmotic stress owing to the high NaCl concentration in soil, but to maintain internal osmotic pressure and water uptake, various organic compounds accumulate and maintain the water gradient (Flowers and Colmer 2008). These compounds protect cellular structures and enzymes (Lutts and Lefèvre 2015). Halophytes are able to synthesize protective compounds rapidly in response to ion toxicity, which describes the ability of halophytes to tolerate heavy metal stress



(Rastgoo and Alemzadeh 2011). An accumulation of proline in cytosol is frequently associated with vacuolar  $\text{Na}^+$  sequestration (Lutts and Lefèvre 2015). This not only results in osmotic function protecting the protein quaternary structure and stabilizing the cell membrane through direct interaction with phospholipid (Rastgoo and Alemzadeh 2011). In *Aeluropus litoralis*, proline content increased heavy metals such as Cd, Co, Pb, and Ag (Lutts and Lefèvre 2015). In a facultative halophyte, *Mesembryanthemum crystallinum*, Cd exposure caused accumulation of proline and total soluble sugars (Lefèvre et al. 2009). In a similar manner, some amino acids such as histidine, glutamine, alanine or asparagine form complexes in response to metal stress (Sharma and Dietz 2006). In *Suaeda salsa*, in response to Cd and Zn, branch chain amino acids such as leucine, isoleucine, valine, glutamine, glutamate, tyrosine, threonine, arginine, phenylalanine, and tryptophan content increased (Lefèvre et al. 2009). For methionine and GSH/PC synthesis, cysteine is required and acts as a central metabolite in antioxidant defense and metal sequestration (Liu et al. 2011). However, whether these accumulations are related to metal tolerance or protein degradation is unclear (Wu et al. 2013). In different halophyte plant species, heavy metal stress and NaCl in combination may trigger the accumulation and synthesis of quaternary ammonium compounds, such as glycine betaine, resulting in the maintenance of membrane integrity, protecting other cellular structures, stabilizing macromolecule structures, and protecting chloroplasts and Photosystem II (Wu et al. 2013). In leaves of *Atriplex halimus*, over-synthesis of glycine betaine in response to Cd was observed (Bose et al. 2014; Ozgur et al. 2013; Lefèvre et al. 2009). *Sesuvium portulacastrum*, a metallo-protectant halophyte, also accumulates trans-4-hydroxyproline betaine and 3,5,4-trihydroxy-6,7 dimethoxyflavone 3-glucoside (Adrian-Romero et al. 1998; Nikalje et al. 2018). This mechanism differs according to variation in species, such as in *S. salsa*, glycine betaine can decrease on exposure to Cd, while choline increases (Liu et al. 2013). Synthesis of Polyamines (PAs) and small aliphatic amines during plant development is considered to have myriad biological functions in response to ion toxicities. They also act as protective molecules to repair and protect stress-induced injuries by forming a direct interaction with numerous cell structures and biochemical responses in *Atriplex halimus* (Lefèvre et al. 2014; Lutts et al. 2012).

### 1.1.3 Salinity

Almost 72% of the Earth's surface is covered with saline water dominated by  $\text{Na}^+$  and  $\text{Cl}^-$  (Flowers and Colmer 2015). In ocean water, usually 19 g of  $\text{Cl}^-$  and 11 g  $\text{Na}^+\text{Kg}^{-1}$  with substantial concentration of  $\text{Mg}^{2+}$ ,  $\text{SO}_4^{2-}$ ,  $\text{K}^+$  and  $\text{Ca}^{2+}$  are found (Flowers and Muscolo 2015). But the essential nutrients  $\text{H}_2\text{PO}_4^-/\text{HPO}_4^{2-}$  and  $\text{NO}_3^-$  required for plant growth are in very low supply (Flowers and Colmer 2015). Even in such diverse conditions, there are a few exceptional plants such as halophytes that can still survive (Flowers et al. 2010). Hence, halophytes are considered the flora of saline environments (Saslis-Lagoudakis et al. 2015). The salt tolerance capacity of

halophytes develops early in some lineages and in many other origins (Saslis-Lagoudakis et al. 2015). Salt tolerance is observed in a few current species, which indicates that the trait has been gained and lost, or various salt tolerant species are becoming extinct throughout evolution (Saslis-Lagoudakis et al. 2015). Phylogenetic analysis can tell us about the macroevolution of salt tolerance, for example,  $C_4$  photosynthesis metabolism has evolved and increased multiple stress tolerance in plants (Saslis-Lagoudakis et al. 2015). The halophyte species are still evolving and relatively new, but have some common features among them (Flowers and Muscolo 2015). This describes recent advancement in halophytes evolution and systematics (Flowers and Muscolo 2015). Their management of salt tolerance is mainly based on the relationship between salt and water. Halophytes are able to reduce the ion concentration of the solution in the xylem owing to which the quantity of salt delivered to the leaves is assimilated by growth and ion excretion through salt glands (Munns and Tester 2008). The ion regulation through the xylem is usually found in all plants to tolerate salinity stress, but halophytes are capable of  $Na^+$  and  $Cl^-$  exclusion at high salt concentrations (Bennett et al. 2009). Ion retrieval is the removal of  $Na^+$  from xylem sap as it rises to the leaves and leads to a genotypic difference in the “apparent exclusion” of  $Na^+$  from entering leaves (Munns and Tester 2008). Mangroves growing in tidal wetland show a range of 90 to 99.6% of external  $Na^+$  as a degree of “exclusion” (Reef and Lovelock 2015). Such a high range of exclusion in mangroves is achieved by anatomical adaptation that decreases and prevents apoplastic movement of solution from outside the root to the xylem, guards the cell membrane, and their transporters may be determined ions that pass into the xylem. Halophytes can tolerate multiple stresses (Flowers and Colmer 2015).

In a solute compartmentalization model of salt tolerance, it was suggested that osmotic adjustment occurs firstly with the ions ( $Na^+$  and  $Cl^-$  mostly) in their environment, but under high concentrations of toxic salts the potential disturbance of metabolism results in ions becoming concentrated in vacuoles, relative to cytosol (Konnerup et al. 2015). Whether concentrations of  $Na^+$  and  $Cl^-$  are toxic when halophytes are exposed to supra-optimal saline condition is not clear. To decrease growth and eventual death, other factors contribute, such as low cytoplasmic  $K^+$ , cellular dehydration, mineral deficiency, reduced stomatal conductance, which limits the net photosynthesis rate, hormonal imbalance, elevated ROS production, and energetic requirement for either ion transport or the synthesis of compatible solutes (Slama et al. 2015). Diverse compatible solutes (amino acid, quaternary ammonium compound, tertiary sulfonium compound, sugars, and sugar alcohols) also explain the role of osmolytes in the cellular compartmentalization model (Flowers et al. 2015). These osmolytes/organic solutes explain the main halophytes families found, their synthesis pathways, and the importance of osmotic adjustment (Flowers and Colmer 2015). Osmolytes act as chemical chaperones and scavengers of hydroxyl radicals and also describe osmotic regulation and transport, which can participate in salt tolerance by enhancing the use of transgenics or by exogenous application to increase these compounds in plant tissue (Flowers and Colmer 2015). Halophytes reduce stomatal opening by generating ROS, which ultimately maintains water transport to diminish the net salt uptake in shoots (Uzilday et al. 2015). Hence, the

water–water cycle enzyme may prevent excessive ROS accumulation in chloroplasts to protect photosynthesis machinery (Hou and Bartels 2015). The main factor behind salt tolerance is the ability of halophytes to regulate  $\text{Na}^+$  and  $\text{Cl}^-$  uptake while maintaining cytoplasmic  $\text{K}^+$  and  $\text{Mg}^+$  concentrations required for essential enzyme activation (Bose et al. 2015).

### ***1.1.4 Combined Stress [Salt + Drought]***

Plant transpiration, or soil evaporation in plants, is increasing because of a rise in the surface temperature resulting from global climate change (Brown and Pezeshki 2007). Halophytes show positive effect of salinity under drought or water scarcity, such as *Sesuvium portulacastrum*, *Zygophyllum xanthoxylum*, *Atriplex halimus*, *Ipomoea pes-capra*, and *Bruguiera cylindrica* (Slama et al. 2007, 2008; Ma et al. 2012; Atreya et al. 2009). Under the combined stress of salinity and water scarcity, the higher leaf  $\text{Na}^+$  concentration, increased stomatal conductance ( $g_s$ ), and net photosynthesis rate ( $P_n$ ) leads to the inference that  $\text{Na}^+$  can enhance photosynthesis and also perform osmotic adjustment for the improvement of water status (Slama et al. 2007; Martínez et al. 2005). Various species show differential involvement of  $\text{Na}^+$  in total osmotic adjustment, such as under the combined stress condition of salt and drought (Slama et al. 2007). *Atriplex halimus* shows a negligible contribution of  $\text{Na}^+$  in total osmotic adjustment (Zribi et al. 2012). Different species have different approaches to dealing with  $\text{Na}^+$ . For example, in *Atriplex halimus*, absorbed  $\text{Na}^+$  accumulates in trichomes or large bladder cells, but in *Sesuvium portulacastrum*,  $\text{Na}^+$  is sequestered in the vacuoles of parenchyma cells (Slama et al. 2007, 2008). In *Sesuvium portulacastrum*,  $\text{Na}^+$  acts as an osmoticum, because under combined drought and salinity, the stress level of proline accumulation is higher than with salinity alone and with drought alone (Hamed et al. 2013) suggesting that there might be a correlation between proline and water potential, i.e., the level of proline increases when the water potential is low (Slama et al. 2007). Under combined stress, proline improves plant tolerance through mechanisms that provide the multiple functions of proline (ROS), scavenging, protein stabilization, and the regulation of cellular redox potential (Szabados and Saviouré 2010; Glenn et al. 2012). Another mechanism for increasing tolerance toward combined stress is reduced stomatal conductance, leading to increased water use efficiency (WUE), which results in a slow growth rate, but accelerated carbon fixed per unit of water transpired (Glenn and Brown 1998). This mechanism has been reported in *Atriplex canescens*, *Atriplex lentiformis*, and *Atriplex halimus* (Zhu and Meinzer 1999; Nemat Alla et al. 2011).

### 1.1.5 Combined Stress [Salinity + Flood]

Large saline areas are unusually wet land habitats such as coastal zones and salt marshes (Flowers and Colmer 2008). To survive in these habitats, plants have to tolerate both tidal flooding and high salinity, because, owing to these stress conditions, root respiration becomes restricted (Baena-González et al. 2007), which leads to decrease in the activity of  $H^+$ -ATPase and inhibits  $Na^+$  and  $K^+$  intake (Barrett-Lennard 2003; Shabala et al. 2014). Exclusion of toxic compounds, such as  $CO_2$ , ethylene, hydrogen sulfide and carboxylic acid can accumulate in plant tissue in response to reduced  $O_2$  and reduces the uptake of essential nutrients, such as  $NH_4^+$  and  $Ca^{2+}$  (Greenway et al. 2006; Bradley and Morris 1990; Rengel 1992). Despite combined flood and salt stress, there are some halophytes, e.g., *Spartina*, that show high productivity. *Spartina* can increase net primary productivity up to 40 tons dry weight (DW)  $ha^{-1} year^{-1}$  (Mishra and Tanna 2017). Barrett-Lennard listed data on waterlogging and salinity interactions for two species, i.e., *Atriplex amnicola* and *Casuarina* sp. (Barrett-Lennard 2003). Further data on a large number of halophytes (15 species) was added by Colmer and Flowers (Flowers and Colmer 2008). Several species show reduced growth when both waterlogging and salinity were imposed (Hamed et al. 2013). A well-known indicator of a wet land ecosystem is *Phragmites australis* (common reed) because it shows an efficient  $Na^+$  exclusion mechanism from leaves and a high  $K^+ : Na^+$  selective ratio under hypoxic conditions over salinity (Gorai et al. 2010). There are some other similar species, e.g., *Melilotus siculus* and *Puccinellia* shoots. Hence, a high value of intrinsic membrane potential is associated with a high  $H^+$  pumping rate, which describes more about  $K^+$  retention in their roots under combined stress conditions (Jenkins et al. 2010). The importance of  $K^+$  retention by roots and shoots for tolerance to combined salinity and hypoxia is explained by comparative studies between halophytes in contrast to water logging (Jenkins et al. 2010). Flower and co-workers used *Suaeda maritima* as a model halophyte for salinity and flood tolerance studies (Flowers 2004). The plant *S. maritima* showed reduced growth, but it is tolerant and survives under saline flooding conditions (Hamed et al. 2013). Research also revealed that *S. maritima* showed increased antioxidant capacity under both natural and controlled saline flood conditions, which was also compared with well-drained conditions (Hamed et al. 2013). Further insights described the contribution of lactate production by roots to tolerate flooding when *S. maritima* was compared with nonhalophyte species such as *Arabidopsis*, tomato, soybean, and tobacco. The former halophyte showed high production of lactate in roots (Teakle et al. 2013). In *S. maritima*, a large amount of lactate is stored in the root vacuoles (concentration of ~120 mM) along with sodium (concentration of more than 200 mM), leading to sodium lactate solution, which prevents vacuole acidification (Teakle et al. 2013). Unlike *S. maritima*, in some other halophyte species, such as *Limonium*, formation of lactate is continuous, but lactate concentration is much lower because most of the lactate is secreted into medium found by Rivoal and Hanson (Flowers and Colmer 2015). To prevent lowering of cytoplasmic pH, it was important for lactate to transport out of

cytoplasm (Bazihizina et al. 2012). Some plants show reduced growth; others were unaffected, but rarely stimulated. Owing to various tolerance traits, such as salt exclusion,  $K^+ : Na^+$  selectivity and antioxidant capacity are amplified in most of the reported species, whereas pH activity and lactose production are important in *S. maritima* roots (Hamed et al. 2013).

### 1.1.6 Combined Stress [Salt + Metal]

A site for the accumulation of industrial and urban effluents contaminated by heavy metals is natural saline soil (Ghnaya et al. 2005; Zaier et al. 2010). Owing to repeated irrigation with sewage sludge or poor quality water, the use of pesticides and phosphorous fertilizers in agricultural soil leads to metal contamination (Hamed et al. 2013). Some halophytic species are accumulators and some are hyper-accumulators of certain metals such as *Sesuvium portulacastrum*, which can tolerate and accumulate high levels of cadmium and lead in aerial parts (Ghnaya et al. 2005; Zaier et al. 2010). In a similar manner, *Mesembryanthemum crystallinum* is a copper hyper-accumulator because it accumulates  $3500 \mu\text{g g}^{-1}$  DW of copper in its tissues. *Sesuvium portulacastrum* produces a higher biomass than *Mesembryanthemum*; hence, *Mesembryanthemum* is a poor candidate for phytoremediation of contaminated soil (Thomas et al. 1998). *Atriplex halimus* has high biomass, but accumulates low concentrations of lead and cadmium; however, it is a viable candidate for use in phytoextraction in arid and saline soils (Manousaki and Kalogerakis 2009). One more suggested suitable species for phytoremediation because of its high translocation rates of cadmium and lead toward above ground tissue is *Halimione portulacoides* (Lutts et al. 2004). *Salicornia maritima* can accumulate reasonable levels of copper and zinc and tolerates moist or dry saline soil, but it is not a metal hyper-accumulator (Reboreda and Caçador 2007). The high metal tolerance of halophytes is associated with salt tolerance traits that include an antioxidant system, and osmoprotectant synthesis, such as proline, which scavenges free radicals and leaf surface excretion by using specialized salt glands (Milić et al. 2012). The major advantage of halophytes is that they can grow and accumulate metal in saline soil (Jithesh et al. 2006; Thomas et al. 1998). In the presence of optimal salinity, when exposed to cadmium stress *Sesuvium portulacastrum* showed improvement in growth and decreased the cadmium accumulation in roots and shoots (Manousaki and Kalogerakis 2011). In contrast, the quantity of cadmium extraction from plants is higher under combined (salinity and heavy metal stress) than under cadmium alone (Manousaki and Kalogerakis 2011; Ghnaya et al. 2007). It was suggested that  $Na^+$  might indirectly contribute to the tolerance of *Sesuvium portulacastrum* from combined stress by osmotic adjustment (Ghnaya et al. 2007). It was found that the most toxic form of cadmium ( $Cd^{2+}$ ) was removed in favor of another form bound to chloride anion and phytochelatins on NaCl treatment (Ghnaya et al. 2005). A comparison of leaves of Cd+NaCl-treated plants and NaCl-treated alone showed

high proline and polyamine concentrations in leaves under Cd+NaCl conditions, which could have helped *Atriplex halimus* to tolerate cadmium inside leaf tissue (Hamed et al. 2013). Salinity reduces the harmful effect of cadmium in halophytes such as *Kosteletzkya virginica* by reduced oxidative stress and altering the hormonal status through reduced ethylene and abscisic acid accumulation (Han et al. 2013).

### **1.1.7 Combined Stress [Salt + Nutrient Deficiency]**

Usually,  $\text{Na}^+$  and  $\text{Cl}^-$  are dominant in saline soil. Seawater has a dominant ion concentration around 500 mM and sodic soil with  $\text{pH} > 8.5$  has  $\text{CO}_3^{2-}$  and  $\text{HCO}_3^-$  as dominant anions (Reboreda and Caçador 2007; Debez Huchzermeyer et al. 2011). Seawater contains 11 mineral nutrients needed by plants in sufficient concentrations for the growth of crops, but deficient in N and  $\text{PO}_4$  (Hamed et al. 2013; Debez et al. 2011). Halophytes grow on seawater with supplied N and  $\text{PO}_4^{2-}$  and show similar biomass production to that of plants grown on full-nutrient solution such as *Suaeda fruticosa* and *Spartina alterniflora* (Zribi et al. 2012). Hence, in the next section the combined stress of N or  $\text{PO}_4^{2-}$  deficiency with salinity is discussed.

### **1.1.8 Combined Stress [Salinity + Nitrogen Deficiency]**

Nitrogen (N) is an essential plant growth element. Higher plants absorb N in two major forms, i.e.,  $\text{NO}_3^-$  (nitrate) and  $\text{NH}_4^+$  (Bennett et al. 2009). However, in halophytes, the form of N used under high salt conditions varies among species and with both  $\text{NH}_4^+$  and  $\text{NO}_3^-$  (Lu et al. 2005). *Suaeda maritima* shows high nitrate reductase activity in low-marsh (waterlogged) in comparison with high-marsh plants (drained) (Britto and Kronzucker 2002; Lewis et al. 1989; Leidi et al. 1991). As plants response to  $\text{NH}_4^+$  has not been studied, the concept is unclear (Mahmood and Kaiser 2003). It was found that the contribution of nitrate in osmotic adjustment is greater than chloride in *Suaeda physophora* in an intertidal zone, even though the soil has a low nitrate concentration and high salinity (Wetson and Flowers 2010). Under low N and high NaCl conditions, some halophytes, such as *Salicornia bigelovii*, *Phragmites australis*, *Glyceria maxima*, and *Spartina alterniflora*, show that ammonium is beneficial for their growth (Song et al. 2006). *Suaeda salsa* accumulates high  $\text{NO}_3^-$  even in soils with low  $\text{NO}_3^-$  content and regulated Cl at low level at intertidal zone. In addition, the PSII machinery remains intact under high salinity, which is an important characteristic of this plant (Song et al. 2009). When  $\text{NH}_4^+$  was dominant,  $\text{K}^+$  regulated growth in *Salicornia bigelovii*. Owing to comparative inhibition,  $\text{Na}^+$  absorption was suppressed by increased  $\text{NH}_4^+$  and consequently  $\text{K}^+$  was selectively absorbed (Kudo and Fujiyama 2010; Munzarova