

SPRINGER BRIEFS IN PHYSICS

Audrius Dubietis
Arnaud Couairon

Ultrafast Supercontinuum Generation in Transparent Solid-State Media

 Springer

SpringerBriefs in Physics

Series Editors

B. Ananthanarayan, Centre for High Energy Physics, Indian Institute of Science, Bangalore, India

Egor Babaev, Physics Department, University of Massachusetts Amherst, Amherst, MA, USA

Malcolm Bremer, H H Wills Physics Laboratory, University of Bristol, Bristol, UK

Xavier Calmet, Department of Physics and Astronomy, University of Sussex, Brighton, UK

Francesca Di Lodovico, Department of Physics, Queen Mary University of London, London, UK

Pablo Esquinazi, Institute for Experimental Physics II, University of Leipzig, Leipzig, Germany

Maarten Hoogerland, University of Auckland, Auckland, New Zealand

Eric Le Ru, School of Chemical and Physical Sciences, Victoria University of Wellington, Kelburn, Wellington, New Zealand

Hans-Joachim Lewerenz, California Institute of Technology, Pasadena, CA, USA

Dario Narducci, University of Milano-Bicocca, Milano, Italy

James Overduin, Towson University, Towson, MD, USA

Vesselin Petkov, Montreal, QC, Canada

Stefan Theisen, Max-Planck-Institut für Gravitationsphys, Golm, Germany

Charles H.-T. Wang, Department of Physics, The University of Aberdeen, Aberdeen, UK

Andrew Whitaker, Department of Physics and Astronomy, Queen's University Belfast, Belfast, UK

SpringerBriefs in Physics are a series of slim high-quality publications encompassing the entire spectrum of physics. Manuscripts for SpringerBriefs in Physics will be evaluated by Springer and by members of the Editorial Board. Proposals and other communication should be sent to your Publishing Editors at Springer.

Featuring compact volumes of 50 to 125 pages (approximately 20,000–45,000 words), Briefs are shorter than a conventional book but longer than a journal article. Thus Briefs serve as timely, concise tools for students, researchers, and professionals.

Typical texts for publication might include:

- A snapshot review of the current state of a hot or emerging field
- A concise introduction to core concepts that students must understand in order to make independent contributions
- An extended research report giving more details and discussion than is possible in a conventional journal article
- A manual describing underlying principles and best practices for an experimental technique
- An essay exploring new ideas within physics, related philosophical issues, or broader topics such as science and society

Briefs are characterized by fast, global electronic dissemination, straightforward publishing agreements, easy-to-use manuscript preparation and formatting guidelines, and expedited production schedules. We aim for publication 8-12 weeks after acceptance.

More information about this series at <http://www.springer.com/series/8902>

Audrius Dubietis · Arnaud Couairon

Ultrafast Supercontinuum Generation in Transparent Solid-State Media

 Springer

Audrius Dubietis
Laser Research Center
Vilnius University
Vilnius, Lithuania

Arnaud Couairon
Centre de Physique Théorique
Ecole polytechnique, CNRS, Institut
Polytechnique de Paris
Paris, France

ISSN 2191-5423

ISSN 2191-5431 (electronic)

SpringerBriefs in Physics

ISBN 978-3-030-14994-9

ISBN 978-3-030-14995-6 (eBook)

<https://doi.org/10.1007/978-3-030-14995-6>

Library of Congress Control Number: 2019934790

© The Author(s), under exclusive license to Springer Nature Switzerland AG 2019

This work is subject to copyright. All rights are solely and exclusively licensed by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

Preface

Nonlinear propagation of intense femtosecond laser pulses in bulk transparent media produces a wealth of physical effects, whose combined action leads to a specific propagation regime, termed femtosecond filamentation, which in turn produces dramatic broadening of the pulse spectrum, termed supercontinuum generation. Due to low cost, compactness, efficiency, robustness, and virtual alignment insensitivity, femtosecond supercontinuum represents a unique and versatile source of coherent ultrabroadband radiation, whose wavelength range spans a considerable part of the optical spectrum. Altogether, these outstanding properties make femtosecond supercontinuum highly suitable for diverse applications in time-resolved spectroscopy, photonics, femtosecond technology, and lightwave engineering. During recent years, this research field has reached a high level of maturity, both in understanding of the underlying physics and in achievement of exciting practical results. In this book, we present the underlying physical picture and overview the state of the art of femtosecond supercontinuum generation in various transparent solid-state media, ranging from wide bandgap dielectrics to semiconductor materials and in various parts of the optical spectrum, from the ultraviolet to the mid-infrared. A particular emphasis is given to the most recent experimental developments: multi-octave supercontinuum generation with pumping in the mid-infrared spectral range, spectral control, power and energy scaling of broadband radiation, and the development of simple, flexible, and robust pulse compression techniques, which deliver few optical cycle pulses and which could be readily implemented in a variety of modern ultrafast laser systems. The expected audience includes graduate students, professionals, and scientists working in business and academia, in the field of laser-matter interactions and ultrafast nonlinear optics.

Vilnius, Lithuania
Paris, France
January 2019

Audrius Dubietis
Arnaud Couairon

Acknowledgments

This book is the result of many years of collaborative research work. We thank members of our own research teams at Centre de Physique Théorique Ecole Polytechnique, CNRS, Institut Polytechnique de Paris (France), and Vilnius University Laser Research Center (Lithuania). We also cordially thank our national and international colleagues and numerous collaborators worldwide, and we wish to convey our gratitude to them through our acknowledgement to their respective institutions: Laboratoire d'Optique Appliquée, ENSTA ParisTech, Palaiseau, France; Femto-ST, Besançon, France; University Jean-Monnet, ST-Etienne, France; Louisiana State University, Baton Rouge, USA; University of Arizona, USA; INRS, Montréal, Canada; University of Insubria, Como & CNR, Italy; Texas A&M University at Qatar and Foundation for Research and Technology-Hellas, Heraklion, Greece; Moscow State University & Lebedev Institute, Moscow, Russia; Joint Institute for High Temperatures, RAS Moscow, Russia; ICFO Barcelona, Spain; University of Glasgow, UK; Lund University, Sweden; Leibniz University, Hannover, Germany; ETH Zurich, Switzerland; East China Normal University, Shanghai, China; and Tata Institute, Mumbai & ACRHEM, Hyderabad, India.

Preparation of this book was greatly facilitated by Tom Spicer, Cindy Zitter, Zach Evenson, and Femina Joshi from Springer Nature. We thank them for their friendly and professional work.

Contents

1	Introduction	1
	References	3
Part I Physical Picture of Supercontinuum Generation		
2	Governing Physical Effects	9
2.1	Self-focusing of Laser Beams	10
2.2	Self-phase Modulation of Laser Pulses	13
2.3	Nonlinear Absorption and Ionization	16
2.4	Plasma Effects	16
2.4.1	Transition of Electrons from the Valence to the Conduction Band	16
2.4.2	Refractive Index Change	17
2.4.3	Plasma-Induced Phase Modulation	17
2.4.4	The Drude–Lorentz Model	19
2.5	Intensity Clamping	20
2.6	Chromatic Dispersion	22
2.7	Self-steepening and Space-Time Focusing	24
2.8	Four-Wave Mixing and Phase Matching	24
	References	25
3	Femtosecond Filamentation in Solid-State Media	27
3.1	Universal Features of Femtosecond Filamentation	28
3.1.1	Conical Emission	28
3.1.2	Plasma Channel Formation	28
3.1.3	Filament Robustness and Energy Reservoir	29
3.1.4	Conical Waves	30
3.2	Numerical Model	33
3.3	Supercontinuum Generation Under Normal GVD	38
3.4	Supercontinuum Generation in the Region of Anomalous GVD	39

3.5 Supercontinuum Generation Under Zero GVD	40
3.6 Conical Emission	41
References	43

Part II Overview of the Experimental Results

4 General Practical Considerations	49
4.1 Materials	49
4.2 External Focusing	52
4.3 Stability Issues	56
4.4 Effect of Filament Refocusing	58
4.5 Multiple Filamentation	60
References	61
5 Experimental Results	65
5.1 Water as a Prototypical Nonlinear Medium	65
5.2 Glasses	67
5.3 Alkali Metal Fluorides	70
5.4 Laser Hosts	74
5.5 Crystals with Second-Order Nonlinearity	78
5.6 Semiconductors	82
5.7 Other Nonlinear Media	86
References	87
6 New Developments	95
6.1 Power and Energy Scaling	95
6.2 Extracavity Pulse Compression	98
6.2.1 Pulse Compression Exploiting SPM in Normally Dispersive Media	99
6.2.2 Soliton Compression Due to Second-Order Cascading	102
6.2.3 Self-compression Through Filamentation	103
6.2.4 Soliton Compression in Isotropic Nonlinear Media with Anomalous GVD	104
6.2.5 Other Compression Mechanisms	107
6.3 Supercontinuum Generation with Picosecond Laser Pulses	108
6.4 Control of Supercontinuum Generation	112
6.5 Supercontinuum Generation with Non-Gaussian Beams	116
6.6 Other Developments	118
References	119

Chapter 1

Introduction



Supercontinuum (SC) generation is one of the most spectacular and visually perceptible effects produced by the nonlinear propagation of intense ultrashort laser pulse in a transparent medium. The discovery of supercontinuum generation in a bulk solid-state medium dates back to the early years of nonlinear optics, when Alfano and Shapiro reported on white light generation, produced by self-focusing of powerful picosecond pulses in a borosilicate glass sample [1]. The discovery was immediately followed by observations of spectral broadening in various crystals and glasses, confirming the universal nature of the phenomenon [2]. (See also [3] for a complete historical account on the early developments of SC generation in various optical media.)

SC generation in bulk media constitutes a compact, efficient, low cost, highly robust, and virtually alignment-insensitive technique for the generation of coherent ultrabroadband radiation at various parts of the optical spectrum [4]. The physical picture of SC generation in transparent bulk media is unveiled in the framework of femtosecond filamentation, which provides a universal scenario of nonlinear propagation and spectral broadening of intense femtosecond laser pulses in bulk solids, liquids, and gases [5–9]. SC generation in bulk media appears to be a complex process that involves an intricate coupling between spatial and temporal effects: diffraction, group velocity dispersion, self-focusing, self-phase modulation, and multiphoton absorption or ionization. In the space domain, the interplay of these effects leads to the formation of a narrow light channel, termed a “light filament” that is able to propagate over extended distances much larger than the typical diffraction length and which leaves a narrow luminous plasma trail in its wake. In the time domain, the pulse undergoes dramatic transformations: pulse splitting or compression, pulse-front steepening, and generation of optical shocks. These transformations altogether produce a broadband, spatially, and temporally coherent emission with a low angular divergence (supercontinuum), which is accompanied by the generation of colored conical emission that is emitted at different angles with respect to the propagation

axis, forming a beautiful array of concentric colored rings. Therefore, SC generation in bulk is markedly different from SC generation in optical fibers, where the propagation dynamics of the optical pulse is essentially one-dimensional and spectral broadening arises from soliton generation and fission due to the interplay between self-phase modulation and material dispersion [10].

SC generation with femtosecond laser pulses was first reported in 1983 [11], long before the phenomenon of femtosecond filamentation was discovered [12]. In that pioneering experiment, Fork and co-authors observed spectral broadening from the deep ultraviolet to the near infrared by focusing intense 80-fs pulses at 627 nm from the dye laser into an ethylene glycol jet [11]. Apart from large-scale spectral broadening, the authors underlined an improvement of pulse-to-pulse reproducibility and spatial uniformity of the beam, which resulted from the short duration of the input pulse; see also [13] and references therein for an account of SC generation in various solid-state and liquid media using femtosecond dye lasers.

A major breakthrough in femtosecond solid-state laser technology was inspired by the groundbreaking invention of chirped pulse amplification (CPA) technique by D. Strickland and G. Mourou [14], which was awarded the Nobel Prize in Physics in 2018. The CPA concept solved the long-standing problem of safe and efficient amplification of ultrashort optical pulses without the onset of optical damage of the amplifier material and other optical components, enabling a tremendous leap in the peak power and intensity of laser pulses. Shortly after that, the discovery of Kerr lens mode locking has led to the invention of femtosecond Ti:sapphire laser oscillator in 1991 [15]. Demonstration of the CPA technique-based regenerative amplification of the Ti:sapphire oscillator pulses, constituted a significant breakthrough in solid-state laser technology and marked a new era in femtosecond SC generation [16]. The amplified Ti:sapphire lasers outperformed then widely spread femtosecond dye lasers in all essential parameters of operation, setting a new standard for the entire femtosecond solid-state laser technology [17].

The commercial availability of novel femtosecond laser sources as combined with a growing practical knowledge of femtosecond SC generation in transparent condensed media [18–20], boosted the development of femtosecond optical parametric amplifiers (OPAs). In that regard, the SC radiation was recognized as an indispensable seeding source for these devices, which produced femtosecond pulses with unprecedented wavelength tunability well exceeding the tuning range afforded by conventional laser sources [21]. Broad spectral bandwidth and high temporal coherence of the SC radiation allowed compressibility of the pulses down to the Fourier transform limit, contributing to the invention of ultrabroadband noncollinear optical parametric amplifiers, the so-called NOPAs [22], which currently deliver few optical cycle pulses at various parts of the optical spectrum, ranging from the visible to the mid-infrared [23].

The advances in the optical parametric amplification techniques fostered exciting developments in the optical parametric chirped amplification (OPCPA). The general idea of the OPCPA was to replace the laser amplifier by the OPA and was originally proposed as an alternative to existing laser amplifiers in 1992 [24]. At present, OPCPA is deservedly regarded as an important offspring of the CPA technique, since

as compared to the laser amplifier, the OPA offers the advantages of very high gain, broad amplification bandwidth, great wavelength flexibility, low thermal effects and superior intensity contrast of the amplified pulses, offering unique possibilities for the amplification of ultrashort laser pulses [25–27]. Interestingly, the first demonstrations of both, CPA and OPCPA, used an optical fiber to broaden the pulse spectrum before the amplification and compression stages. In that regard, SC generation in condensed bulk media offered a number of advantages due to its robustness and compactness, enabling to elaborate novel and compact architectures of tabletop SC-seeded OPCPA systems [28]. In particular, a considerable effort is currently directed to the development of the SC-seeded OPCPA systems that deliver intense few optical cycle pulses in the mid-infrared spectral region, see, e.g., [29], which is out of the grasp for existing mid-infrared solid-state lasers and laser amplifiers, see, e.g., [30].

These developments in turn facilitated experimental studies of SC generation in the region of anomalous group velocity dispersion (GVD) of dielectric solid-state media, yielding ultrabroadband, multioctave SC with unprecedented wavelength coverage, see, e.g., [31–33]. Moreover, using long-wavelength ultrashort pulses, SC generation was made possible in various highly nonlinear materials, such as narrow bandgap dielectric crystals, soft glasses, and semiconductors to produce octave-spanning SC spectra extending into far infrared. From a future perspective, SC generation represents one of the fundamental building blocks of the emerging third-generation femtosecond technology, which foresees boosting the peak and average powers of few optical cycle pulses simultaneously to the multiterawatt and hundreds of watts range, respectively, thereby paving the way for the generation of powerful sub-cycle pulses with full control over the generated light waves [34]. Finally, within the past decade, the term “supercontinuum generation” has been extended well beyond the optical range, to include high-order and nonperturbative nonlinear optical processes, such as high harmonic generation in the vacuum ultraviolet and X-ray ranges [35].

References

1. Alfano, R.R., Shapiro, L.: Emission in the region 4000 to 7000 Å via four photon coupling in glass. *Phys. Rev. Lett.* **24**, 584–587 (1970)
2. Alfano, R.R., Shapiro, L.: Observation of self-phase modulation and small-scale filaments in crystals and glasses. *Phys. Rev. Lett.* **24**, 592–594 (1970)
3. Alfano, R.R. (ed.): *The Supercontinuum Laser Source*. Springer (2006)
4. Dubietis, A., Tamošauskas, G., Šuminas, R., Jukna, V., Couairon, A.: Ultrafast supercontinuum generation in bulk condensed media (Review). *Lith. J. Phys.* **57**, 113–157 (2017)
5. Chin, S.L., Hosseini, S.A., Liu, W., Luo, Q., Thberge, F., Aközbek, N., Becker, A., Kandidov, V.P., Kosareva, O.G., Schroeder, H.: The propagation of powerful femtosecond laser pulses in optical media: physics, applications, and new challenges. *Can. J. Phys.* **83**, 863–905 (2005)
6. Couairon, A., Mysyrowicz, A.: Femtosecond filamentation in transparent media. *Phys. Rep.* **441**, 47–190 (2007)
7. Kasparian, J., Wolf, J.-P.: Physics and applications of atmospheric nonlinear optics and filamentation. *Opt. Express* **16**, 466–493 (2008)

8. Kandidov, V.P., Shlenov, S.A., Kosareva, O.G.: Filamentation of high-power femtosecond laser radiation. *Quantum Electron.* **39**, 205–228 (2009)
9. Chin, S.L.: *Femtosecond Laser Filamentation*. Springer (2010)
10. Dudley, J.M., Genty, G., Coen, S.: Supercontinuum generation in photonic crystal fiber. *Rev. Mod. Phys.* **78**, 1135–1184 (2006)
11. Fork, R.L., Shank, C.V., Hirlimann, C., Yen, R., Tomlinson, W.J.: Femtosecond white-light continuum pulses. *Opt. Lett.* **8**, 1–3 (1983)
12. Braun, A., Korn, G., Liu, X., Du, D., Squier, J., Mourou, G.: Self-channeling of high-peak-power femtosecond laser pulses in air. *Opt. Lett.* **20**, 73–75 (1995)
13. Wittmann, M., Penzkofer, A.: Spectral superbroadening of femtosecond laser pulses. *Opt. Commun.* **126**, 308–317 (1996)
14. Strickland, D., Mourou, G.: Compression of amplified chirped optical pulses. *Opt. Commun.* **56**, 219–221 (1985)
15. Spence, D.E., Kean, P.N., Sibbett, W.: 60-fsec pulse generation from a self-mode-locked Ti:sapphire laser. *Opt. Lett.* **16**, 42–44 (1991)
16. Norris, T.B.: Femtosecond pulse amplification at 250 kHz with a Ti:sapphire regenerative amplifier and application to continuum generation. *Opt. Lett.* **17**, 1009–1011 (1992)
17. Backus, S., Durfee, C.G., Murnane, M.M., Kapteyn, H.C.: High power ultrafast lasers. *Rev. Sci. Instr.* **69**, 1207–1223 (1998)
18. Brodeur, A., Chin, S.L.: Band-gap dependence of the ultrafast white-light continuum. *Phys. Rev. Lett.* **80**, 4406–4409 (1998)
19. Brodeur, A., Chin, S.L.: Ultrafast white-light continuum generation and self-focusing in transparent condensed media. *J. Opt. Soc. Am. B* **16**, 637–650 (1999)
20. Bradler, M., Baum, P., Riedle, E.: Femtosecond continuum generation in bulk laser host materials with sub- μ J pump pulses. *Appl. Phys. B* **97**, 561–574 (2009)
21. Cerullo, G., De Silvestri, S.: Ultrafast optical parametric amplifiers. *Rev. Sci. Instr.* **74**, 1–18 (2003)
22. Wilhelm, T., Piel, J., Riedle, E.: Sub-20-fs pulses tunable across the visible from a blue-pumped single pass noncollinear parametric converter. *Opt. Lett.* **22**, 1494–1496 (1997)
23. Brida, D., Manzoni, C., Cirmi, G., Marangoni, M., Bonora, S., Villoresi, P., De Silvestri, S., Cerullo, G.: Few-optical-cycle pulses tunable from the visible to the mid-infrared by optical parametric amplifiers. *J. Opt.* **12**, 013001 (2010)
24. Dubietis, A., Jonušauskas, G., Piskarskas, A.: Powerful femtosecond pulse generation by chirped and stretched pulse parametric amplification in BBO crystal. *Opt. Commun.* **88**, 437–440 (1992)
25. Dubietis, A., Butkus, R., Piskarskas, A.P.: Trends in chirped pulse optical parametric amplification. *IEEE J. Sel. Topics Quantum Electron.* **12**, 163–172 (2006)
26. Witte, S., Eikema, K.S.E.: Ultrafast optical parametric chirped-pulse amplification. *IEEE J. Sel. Topics Quantum Electron.* **18**, 296–307 (2012)
27. Vaupel, A., Bodnar, N., Webb, B., Shah, L., Richardson, M.: Concepts, performance review, and prospects of table-top, few-cycle optical parametric chirped-pulse amplification. *Opt. Eng.* **53**, 051507 (2014)
28. Budriūnas, R., Stanislauskas, T., Adamonis, J., Aleknavičius, A., Veitas, G., Gadonas, D., Balickas, S., Michailovas, A., Varanavičius, A.: 53 W average power CEP-stabilized OPCPA system delivering 5.5 TW few cycle pulses at 1 kHz repetition rate. *Opt. Express* **25**, 5797–5806 (2017)
29. Rigaud, P., van de Walle, A., Hanna, M., Forget, N., Guichard, F., Zaouter, Y., Guesmi, K., Druon, F., Georges, P.: Supercontinuum-seeded few-cycle midinfrared OPCPA system. *Opt. Express* **24**, 26494–26502 (2016)
30. Pires, H., Baudisch, M., Sanchez, D., Hemmer, M., Biegert, J.: Ultrashort pulse generation in the mid-IR. *Prog. Quantum Electron.* **43**, 1–30 (2015)
31. Silva, F., Austin, D.R., Thai, A., Baudisch, M., Hemmer, M., Faccio, D., Couairon, A., Biegert, J.: Multi-octave supercontinuum generation from mid-infrared filamentation in a bulk crystal. *Nature Commun.* **3**, 807 (2012)