

SECOND EDITION

INTRODUCTION TO
BIOLOGICAL PHYSICS

FOR THE HEALTH AND LIFE SCIENCES

Kirsten Franklin
Paul Muir
Terry Scott
Paul Yates

The Staff of the Department of Physics,
University of Otago

WILEY

Introduction to Biological Physics for the Health and Life Sciences

Introduction to Biological Physics for the Health and Life Sciences

Second Edition

Kirsten Franklin

Paul Muir

Terry Scott

Paul Yates

And staff of the Department of Physics
University of Otago

WILEY

This edition first published 2019
© 2019 John Wiley & Sons, Ltd

Edition History

John Wiley & Sons Ltd (1e, 2010)

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by law. Advice on how to obtain permission to reuse material from this title is available at <http://www.wiley.com/go/permissions>.

The right of Kirsten Franklin, Paul Muir, Terry Scott and Paul Yates to be identified as the authors of this work has been asserted in accordance with law.

Registered Offices

John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, USA
John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Office

The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

For details of our global editorial offices, customer services, and more information about Wiley products visit us at www.wiley.com.

Wiley also publishes its books in a variety of electronic formats and by print-on-demand. Some content that appears in standard print versions of this book may not be available in other formats.

Limit of Liability/Disclaimer of Warranty

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting scientific method, diagnosis, or treatment by physicians for any particular patient. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. While the publisher and authors have used their best efforts in preparing this work, they make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives, written sales materials or promotional statements for this work. The fact that an organization, website, or product is referred to in this work as a citation and/or potential source of further information does not mean that the publisher and authors endorse the information or services the organization, website, or product may provide or recommendations it may make. This work is sold with the understanding that the publisher is not engaged in rendering professional services. The advice and strategies contained herein may not be suitable for your situation. You should consult with a specialist where appropriate. Further, readers should be aware that websites listed in this work may have changed or disappeared between when this work was written and when it is read. Neither the publisher nor authors shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

Library of Congress Cataloging-in-Publication Data

Names: Franklin, Kirsten, author.

Title: Introduction to biological physics for the health and life sciences / Kirsten Franklin, Paul Muir, Terry Scott, Paul Yates, and staff of the Department of Physics, University of Otago.

Description: Second edition. | Chichester, West Sussex, United Kingdom : John Wiley & Sons Ltd, [2019] |

Includes bibliographical references and index. |

Identifiers: LCCN 2018051884 (print) | LCCN 2018060214 (ebook) |

ISBN 9781118934494 (Adobe PDF) | ISBN 9781118934487 (ePub) |

ISBN 9781118934500 (pbk.) | ISBN 1118934504 (pbk.)

Subjects: LCSH: Biophysics--Textbooks. | Physics--Textbooks. | Medical physics--Textbooks.

Classification: LCC QH505 (ebook) | LCC QH505 .I68 2019 (print) |

DDC 571.4--dc23

LC record available at <https://lccn.loc.gov/2018051884>

Cover image: Human heart circulation in a skeleton © shefkate/Adobe, Red blood cells,
© ktsdesign/Adobe, Woman's eye, © freshidea/Adobe, all other images courtesy of Michael Crawford
Cover design by Michael Crawford

Set in 10/12pt TradeGothicLTStd by Aptara Inc., New Delhi, India

10 9 8 7 6 5 4 3 2 1

CONTENTS

I Mechanics

CHAPTER 1	Kinematics	3	CHAPTER 6	Momentum	61
1.1	Introduction	3	6.1	Introduction	61
1.2	Distance and Displacement	3	6.2	Linear Momentum	61
1.3	Speed and Velocity	3	6.3	Newton's Laws and Momentum	61
1.4	Acceleration	4	6.4	Collisions	62
1.5	Average Velocity or Speed	6	6.5	Elastic Collisions	64
1.6	Change in Displacement Under Constant Acceleration	7	6.6	Summary	66
1.7	The Acceleration Due to Gravity	8	6.7	Problems	66
1.8	Independence of Motion in Two Dimensions	10	CHAPTER 7	Simple Harmonic Motion	69
1.9	Summary	13	7.1	Introduction	69
1.10	Problems	14	7.2	Hooke's Law	69
CHAPTER 2	Force and Newton's Laws of Motion	17	7.3	Simple Harmonic Motion	70
2.1	Introduction	17	7.4	The Simple Pendulum	76
2.2	The Concept of Force	17	7.5	Summary	76
2.3	Kinds of Force	22	7.6	Problems	77
2.4	Newtonian Gravity	25	CHAPTER 8	Waves	79
2.5	Fictitious Forces	26	8.1	Introduction	79
2.6	Summary	27	8.2	SHM and Waves	79
2.7	Problems	27	8.3	Frequency, Wavelength, and Speed	80
CHAPTER 3	Motion in a Circle	31	8.4	The Form of the Wave	81
3.1	Introduction	31	8.5	Types of Wave	82
3.2	Description of Circular Motion	31	8.6	Superposition and Interference	83
3.3	Circular Velocity and Acceleration	32	8.7	Beats	83
3.4	Centripetal Force	33	8.8	Reflection	85
3.5	Sources of Centripetal Force	33	8.9	Standing Waves	85
3.6	Summary	34	8.10	Waves and Energy	85
3.7	Problems	35	8.11	Complex Waveforms	86
CHAPTER 4	Statics	37	8.12	Summary	88
4.1	Introduction	37	8.13	Problems	88
4.2	Equilibrium	37	CHAPTER 9	Sound and Hearing	91
4.3	Torque	38	9.1	Introduction	91
4.4	The Principle of Moments	38	9.2	Sound Waves in Media	91
4.5	Centre of Gravity/Centre of Mass	39	9.3	Pitch and Loudness	94
4.6	Stability	43	9.4	Resonance and Sound Generation	96
4.7	Summary	43	9.5	The Ear	100
4.8	Problems	43	9.6	The Doppler Effect	102
CHAPTER 5	Energy	47	9.7	Summary	104
5.1	Introduction	47	9.8	Problems	105
5.2	What is Energy?	47	II Solids and Fluids	107	
5.3	Work	47	CHAPTER 10	Elasticity: Stress and Strain	109
5.4	Kinetic Energy	49	10.1	Introduction	109
5.5	Potential Energy	51	10.2	Tension and Compression	109
5.6	Conservative Forces	52	10.3	Shear Stress and Strain	112
5.7	Conservation of Total Energy	52	10.4	Bulk Stress and Strain	113
5.8	Power	53	10.5	Elasticity	114
5.9	Summary	58	10.6	Summary	116
5.10	Problems	58	10.7	Problems	116
			CHAPTER 11	Pressure	119
			11.1	Introduction	119
			11.2	Pressure	119
			11.3	Density	120
			11.4	Pascal's Principle	121
			11.5	Measurement of Pressure	124
			11.6	Pressure and the Human Body	129
			11.7	Summary	129
			11.8	Problems	130
			CHAPTER 12	Buoyancy	133
			12.1	Introduction	133
			12.2	The Buoyant Force	133
			12.3	Summary	137
			12.4	Problems	138

CHAPTER 13	Surface Tension and Capillarity	141	CHAPTER 21	Heat Transfer	227
13.1	Introduction	141	21.1	Introduction	227
13.2	Surface Tension	141	21.2	Conduction	227
13.3	Capillarity	144	21.3	Convection	229
13.4	Surfactants and the Lung	146	21.4	Radiation	230
13.5	Summary	147	21.5	Combined Transfer Processes	232
13.6	Problems	147	21.6	Summary	234
			21.7	Problems	235
CHAPTER 14	Fluid Dynamics of Non-viscous Fluids	149	CHAPTER 22	Thermodynamics and the Body	239
14.1	Introduction	149	22.1	Introduction	239
14.2	Definitions of Some Key Terms	149	22.2	The First Law	239
14.3	The Equation of Continuity	149	22.3	Energy and the Body	239
14.4	Bernoulli's Equation	151	22.4	Thermoregulation	242
14.5	Summary	155	22.5	Temperature and Health	244
14.6	Problems	155	22.6	Summary	245
			22.7	Problems	245
CHAPTER 15	Fluid Dynamics of Viscous Fluids	159	CHAPTER 23	Thermodynamic Processes in Ideal Gases	249
15.1	Introduction	159	23.1	Introduction	249
15.2	Viscosity	159	23.2	States, Processes, and Equilibrium	249
15.3	Turbulence	161	23.3	Reversibility	251
15.4	Summary	162	23.4	Work and P - V Diagrams	252
15.5	Problems	162	23.5	Isobaric, Isochoric, Isothermal, and Adiabatic Processes	254
			23.6	Summary	258
CHAPTER 16	Molecular Transport Phenomena	165	23.7	Problems	259
16.1	Introduction	165	CHAPTER 24	Heat Engines And Entropy	263
16.2	Diffusion	165	24.1	Introduction	263
16.3	Osmosis	166	24.2	The Second Law of Thermodynamics	263
16.4	Applications to Biological Systems	167	24.3	Entropy	264
16.5	Summary	168	24.4	Cyclic Processes and Heat Engines	266
16.6	Problems	168	24.5	The First Law for Reversible Processes	270
			24.6	T - S Diagrams for Heat Engines	270
III Thermodynamics		171	24.7	Entropy and Irreversible Processes	273
CHAPTER 17	Temperature and the Zeroth Law	173	24.8	Absolute Entropy And The Third Law of Thermodynamics	274
17.1	Introduction	173	24.9	Summary	275
17.2	Thermal Equilibrium	173	24.10	Problems	276
17.3	Measuring Temperature	175	CHAPTER 25	Energy Availability and Thermodynamic Potentials	279
17.4	Thermal Expansion of Materials	177	25.1	Introduction	279
17.5	Summary	181	25.2	Enthalpy	279
17.6	Problems	182	25.3	Helmholtz Energy	282
			25.4	Gibbs Energy	283
CHAPTER 18	Ideal Gases	185	25.5	Chemical Work and the Chemical Potential	284
18.1	Introduction	185	25.6	Thermodynamic Potentials and Equilibrium	285
18.2	The Gas Laws	185	25.7	Heat Engines and the Efficiency of Metabolism	287
18.3	Biological Applications	190	25.8	Summary	290
18.4	Kinetic Theory of Gases	191	25.9	Problems	291
18.5	Summary	195			
18.6	Problems	195	IV Electricity and DC Circuits	293	
			CHAPTER 26	Static Electricity	295
CHAPTER 19	Phase and Temperature Change	199	26.1	Introduction	295
19.1	Introduction	199	26.2	Charge	295
19.2	Phase Changes	199	26.3	Conductors and Insulators	297
19.3	Temperature Changes	204	26.4	Charging of Objects	297
19.4	Energy Conservation	204	26.5	Polarisation	298
19.5	L and c Values for Water	207	26.6	Summary	299
19.6	Summary	208	26.7	Problems	300
19.7	Problems	209	CHAPTER 27	Electric Force and Electric Field	301
			27.1	Introduction	301
CHAPTER 20	Water Vapour and the Atmosphere	213	27.2	Coulomb's Law	301
20.1	Introduction	213	27.3	Superposition of Electric Forces	302
20.2	Mixtures of Water Vapour and Air	213	27.4	Inverse Square Laws	304
20.3	Partial Pressure and Moisture Content	214	27.5	The Electric Field	305
20.4	Atmospheric Properties	215			
20.5	Psychrometry	216			
20.6	Applications	222			
20.7	Summary	222			
20.8	Problems	223			

27.6	Electric Field Diagrams	305	33.6	Lenses	382
27.7	Superposition of Electric Fields	307	33.7	Summary	388
27.8	Summary	307	33.8	Problems	389
27.9	Problems	308			
CHAPTER 28	Electrical Potential and Energy	311	CHAPTER 34	The Eye and Vision	393
28.1	Introduction	311	34.1	Introduction	393
28.2	Electrical Potential Energy	311	34.2	The Parts of the Eye	393
28.3	Electrical Potential	312	34.3	Emmetropia (Normal Vision)	396
28.4	Electrical Potential and Work	313	34.4	Presbyopia	397
28.5	Equipotential and Field Lines	315	34.5	Myopia	399
28.6	Electrical and External Forces	316	34.6	Hypermetropia (or Hyperopia)	402
28.7	The Heart and ECG	318	34.7	Astigmatism	403
28.8	Summary	320	34.8	Alternative Structure and Placement	403
28.9	Problems	320	34.9	Colour Vision	404
			34.10	Summary	407
			34.11	Problems	407
CHAPTER 29	Capacitance	323	CHAPTER 35	Wave Optics	411
29.1	Introduction	323	35.1	Introduction	411
29.2	The Capacitor	323	35.2	Superposition and Interference	411
29.3	Energy Stored in a Capacitor	325	35.3	Huygens' Principle	412
29.4	Capacitors in Series and Parallel	326	35.4	Diffraction	414
29.5	The Dielectric in a Capacitor	328	35.5	Young's Double-Slit Experiment	414
29.6	Summary	330	35.6	Single-Slit Diffraction	416
29.7	Problems	330	35.7	Diffraction Gratings	419
			35.8	Circular Apertures and Diffraction	420
			35.9	Visual Acuity	420
CHAPTER 30	Direct Currents and DC Circuits	333	35.10	Thin-Film Interference	422
30.1	Introduction	333	35.11	Polarisation	422
30.2	Electric Current	333	35.12	Summary	425
30.3	Current and Drift Velocity	334	35.13	Problems	425
30.4	Direct Versus Alternating Current	335			
30.5	Circuits and Circuit Diagrams	336	CHAPTER 36	Advanced Geometric Optics	429
30.6	Power Sources	336	36.1	Introduction	429
30.7	Resistance and Ohm's Law	336	36.2	Image Formation by Reflection at a Spherical Surface	429
30.8	Resistors and Resistivity	337	36.3	Image Formation by a Refraction at a Single Spherical Surface	432
30.9	Wires	338	36.4	Image Formation by a Thin Lens	437
30.10	Kirchhoff's Laws	339	36.5	Vergences	440
30.11	Resistors in Series and Parallel	341	36.6	Multiple Lenses and Thick Lenses	446
30.12	Power Dissipation	342	36.7	Summary	447
30.13	Alternate Energy Units	343	36.8	Problems	447
30.14	Electric Shock Hazards	343			
30.15	Electricity in Cells	344	CHAPTER 37	Optical Instruments	449
30.16	Summary	346	37.1	Introduction	449
30.17	Problems	347	37.2	Single Converging Lens: The Magnifying Glass	449
			37.3	Microscopes	452
			37.4	Telescopes	455
			37.5	Summary	457
			37.6	Problems	457
CHAPTER 31	Time Behaviour of RC Circuits	351			
31.1	Introduction	351	VI	Radiation and Health	461
31.2	The RC Circuit	351	CHAPTER 38	Atoms and Atomic Physics	463
31.3	Discharging RC Circuit	352	38.1	Introduction	463
31.4	Charging RC Circuit	353	38.2	Parts of the Atom	463
31.5	Summary	356	38.3	Electron Orbitals	464
31.6	Problems	356	38.4	The Böhr Model of the Atom	465
			38.5	Multielectron Atoms	469
			38.6	Quantum Mechanics	470
			38.7	Summary	471
			38.8	Problems	472
V	Optics	359	CHAPTER 39	The Nucleus and Nuclear Physics	475
CHAPTER 32	The Nature of Light	361	39.1	Introduction	475
32.1	Introduction	361	39.2	Nuclei and Isotopes	475
32.2	Electromagnetic Waves	361	39.3	Energy and Mass Units	476
32.3	Reflection	363	39.4	Nuclear Forces	478
32.4	Refraction	364	39.5	Nuclear Decay and Stability	479
32.5	Dispersion	368	39.6	Summary	483
32.6	Summary	370			
32.7	Problems	371			
CHAPTER 33	Geometric Optics	375			
33.1	Introduction	375			
33.2	Ray Diagrams	375			
33.3	Plane Mirrors	377			
33.4	Spherical Mirrors	377			
33.5	Magnification	380			

39.7	Problems	483			
CHAPTER 40	Production of Ionising Radiation	485			
40.1	Introduction	485			
40.2	Nuclear Decay Processes	485			
40.3	Activity and Half-Life	488			
40.4	X-ray Production	491			
40.5	Other Sources of Radiation	494			
40.6	Summary	495			
40.7	Problems	496			
CHAPTER 41	Interactions of Ionising Radiation	499			
41.1	Introduction	499			
41.2	Attenuation and Cross Section	499			
41.3	X-rays and Gamma Radiation	501			
41.4	Particles	505			
41.5	Detection of Ionising Radiation	506			
41.6	Summary	507			
41.7	Problems	508			
CHAPTER 42	Biological Effects of Ionising Radiation	509			
42.1	Introduction	509			
42.2	Mechanisms of Cell Damage	509			
42.3	Dose and Dose Equivalent	510			
42.4	Types of Effect	513			
42.5	Medical Effects and Risk	514			
42.6	Ultraviolet Radiation	514			
42.7	Summary	515			
42.8	Problems	515			
CHAPTER 43	Medical Imaging	519			
43.1	Introduction	519			
43.2	X-ray Imaging	519			
43.3	CT Scan	521			
43.4	PET scan	521			
			43.5	Gamma Camera and SPECT	522
			43.6	Diagnostic Procedures: Dose	522
			43.7	Ultrasound Sonography	522
			43.8	Summary	524
			CHAPTER 44	Magnetism and MRI	525
			44.1	Introduction	525
			44.2	Magnetism	526
			44.3	A Brief Outline of MRI	530
			44.4	Nuclear Magnetic Resonance	531
			44.5	Magnetic Resonance Imaging	537
			44.6	Summary	545
			44.7	Problems	546
			Appendices		550
			APPENDIX A	Physical Constants	551
			A.1	High Precision Mass Values	551
			A.2	Useful Constants	551
			APPENDIX B	Basic Maths and Science Skills	553
			B.1	Measurement and Units	553
			B.2	Basic Algebra	557
			B.3	Exponentials and Logarithms	559
			B.4	Geometry	561
			B.5	Trigonometric Functions	561
			B.6	Vectors	562
			APPENDIX C	Answers To Odd Numbered Problems	565
			SELECTED FURTHER READING		576
			INDEX		579

PREFACE

Second Edition

In the preface to the first edition of this textbook, we stated our primary goals in writing this book. These goals have not changed: we hope to provide the needed basis in physics to health-science students. However, in the second edition of this book we have an additional goal. We recognise that there are a number of physical ideas which appear in other areas in the health sciences which are often introduced in a very cursory way. In this book we hope to provide health-science students with an explanation of these concepts which, while not introductory, may be useful as reference material in other courses and later in their careers as health professionals. As well as broadening the scope of the book, we have taken the opportunity presented by a second edition to plug some gaps in the coverage, correct some slight errors, and to smooth some of the rough edges of the first edition.

Students, like everyone else, learn more effectively by doing and in physics that means working through problems, and they frequently request sets of questions to try. This second edition attempts to provide more support for this perennial request. We have added many new problems across the book, with more variety in the difficulty level to make them accessible to students at all levels, and to provide some problems which will challenge students at all levels.

In addition to a large number of new problems, we have added a number of new sections to the book. The primary additions are to the Optics and Thermodynamics topics. The new chapters in the Optics topic cover optical instrumentation and go deeper into geometric optics by looking at refraction by a single spherical surface. Optical instruments are central to both research and everyday clinical practice in the health sciences, and a basic understanding of magnifiers is useful. The method of vergences is also introduced as an alternative to the standard thin-lens equation approach to image formation, which allows for the more sophisticated reduced eye model to be introduced briefly.

The second large addition to the textbook is a new section in the Thermodynamics topic with chapters covering thermodynamic processes, heat engines, entropy, and thermodynamic potentials. The material presented in these chapters goes further into some thermodynamics concepts than is typical for a first-year physics text. However, students are confronted with concepts like entropy and Gibb's energy in other courses they undertake as part of their training. These students cannot be expected to develop a deep understanding of these ideas in a single term, especially in a course whose main purpose is to teach chemistry or biochemistry. Our primary goal in this section of the book is to provide students with a secondary source of information which they may access later in their course of study, though it may also prove useful in joining physics with chemistry in integrated science programmes.

Finally, we would like to express our gratitude to the many people who have helped in the production of this edition. Firstly, we would like to thank our students whose continuing search for understanding has continued to keep us honest. Secondly, the support we have received from our colleagues in the Department of Physics at the University of Otago has made the completion of both of the editions of this book possible. Finally we would like to thank Sam Belcher and Professor Gerry Carrington. Sam read through the entire book and worked through every single problem. The errors and other difficulties he found have greatly improved this book. The assistance we received from Professor Carrington in the design and writing of the new thermodynamics chapter has been invaluable. His insight and experience in this field have greatly strengthened both the text-

book and our own understanding of the field.

We would also like to acknowledge the contribution of Lara Wilcocks to the development of the first edition of this book.

We have subjected the text of this book to pitiless scrutiny, however it is unavoidable that errors will remain. For these mistakes we apologise in advance. While unavoidable they are still our responsibility alone.

First Edition

Physics is central to an understanding of biomedical science. We are aware that many students studying for a career in biomedicine are not primarily motivated by physics; they are interested in other areas of science. We are also aware that no currently available first-year textbook takes the physics needs of health-science students seriously.

In this textbook we have several goals. Firstly, we are trying to present the necessary base concepts of physics as clearly as possible. Secondly, the textbook is designed to remove any unnecessary conceptual load from students by removing all physics that is not absolutely necessary for health-science students. The decision as to which parts of physics are necessary has been determined in close collaboration with the physicists and teachers of the Department of Physics and the professional clinicians and academics in the Faculty of Health Science at the University of Otago. Thirdly, we are keenly aware that student motivation is always an issue in the study of physics for the health sciences. We have tried to add as many applications to the biomedical sciences as possible to the text in an attempt to aid this motivation. The companion website for this book is available at www.wiley.com/go/biological_physics.

The production of a textbook is an enormous task and this textbook is no exception. In writing this book, we have relied on the expertise and goodwill of a large group of academic colleagues. We would like to express our gratitude to Mr Gordon Sanderson of the Ophthalmology Department and Professor Terence Doyle of the Radiology Department in the University of Otago Medical School. We would like to thank Dai Redshaw for the many hours he has spent reading through the text and working through the problem sets and Dr Phil Sheard from the Department of Physiology for his inspiring review lectures on bioelectricity. We would particularly like to thank Dr Don Warrington for his diligent and careful reading of the entire manuscript, and for his many corrections and suggestions. Finally we would like to thank the staff of the Department of Physics at Otago for the time and support that they have rendered over the past years. While the staff of the Department of Physics are listed as authors of this textbook we would particularly like to thank Gerry Carrington, Pat Langhorne, Craig Rodger, Rob Ballagh, Neil Thomson, and Bob Lloyd.

Finally, the goal of this textbook is to provide for the needs of our students. In order to achieve this goal, we have depended on the feedback provided by our students. There will of course still be errors which have escaped our editing process, and for these we apologise in advance, and we welcome feedback from our readers.

I

Mechanics

Mechanics is the study of motion. It may be divided into two related areas: kinematics and dynamics. Kinematics is the study of the fundamental properties of motion: displacement, acceleration, velocity, distance, and speed. These concepts allow us to quantify motion and this allows for its scientific study. Dynamics is the study of force as described by Newton's three laws. Forces produce accelerations and thus cause changes in the motion of objects.

Mechanics is a fundamental subject in physics: it shows how the forces of nature produce the changes which are observed in nature. The concepts introduced in mechanics will be used throughout the rest of this book.

Mechanics is of central importance in the health sciences. The applications of mechanics in biological systems appear whenever the concepts of force, energy, or momentum appear. There are, however, many more direct applications of the ideas of mechanics. The working of the musculoskeletal system in humans and other vertebrates cannot be understood without an understanding of mechanical concepts such as torque, force, levers, and tension. The energy and forces required for everyday activities in nature – jumping, flying, accelerating to elude capture – can only be evaluated using the techniques introduced in this section.

KINEMATICS

1

1.1	Introduction
1.2	Distance and Displacement
1.3	Speed and Velocity
1.4	Acceleration
1.5	Average Velocity or Speed
1.6	Change in Displacement Under Constant Acceleration
1.7	The Acceleration Due to Gravity
1.8	Independence of Motion in Two Dimensions
1.9	Summary
1.10	Problems

1.1 Introduction

Kinematics is that part of mechanics which is concerned with the description of motion. This is a vital first step in coming to an understanding of motion, since we will not be able to describe its causes, or how it changes, without a clear understanding of the properties of motion. Kinematics is about the definition and clarification of those concepts necessary for the complete description of motion. Only six concepts are needed: time, distance, displacement, speed, velocity, and acceleration.

We will begin by focussing on linear motion in one dimension. Later we will expand this to include motion in two and three dimensions, and we will then look at three particularly important special cases of motion in one and two dimensions: circular motion, simple harmonic motion, and wave motion.

Key Objectives

- To develop an understanding of the concepts used to describe motion: time, distance, displacement, speed, velocity, and acceleration
- To understand the relationships between time, displacement, velocity, and acceleration
- To understand the distinction between average and instantaneous velocity and acceleration
- To understand that the horizontal and vertical components of vector quantities, such as acceleration and velocity, may be treated independently

1.2 Distance and Displacement

Motion is characterised by the direction of movement, as well as the amount of movement involved. It is not surprising that we must use vector quantities in kinematics. The **distance** an object travels is defined as the length of the path that the object took in travelling from one place to another. Distance is a scalar quantity. **Displacement**, on the other hand, is the distance travelled, but with a direction associated. Thus a road trip of 100 km to the north covers the same distance as a road trip of 100 km to the south, but these two trips have quite different displacements. The use of displacement rather than distance to give directions is commonplace.

1.3 Speed and Velocity

We are accustomed to talking about the speed at which an object is moving. We also talk about the velocity with which an object is moving. In normal usage, these two words mean the same thing. We can talk about the speed with which a car is travelling, or we can talk about its velocity. In physics, we redefine these two words, **speed** and **velocity**, so that they have similar, but distinct meanings.




Figure 1.1 A toy car on a race track. How do we characterise its motion?

Key concept:

The **velocity** of an object is the change in its position, divided by the time it took for this change to occur. Velocity is a vector and has both a magnitude and a direction.

Mathematically, the velocity of an object is

$$\mathbf{v} = \frac{\Delta \mathbf{x}}{\Delta t} \tag{1.1}$$

where \mathbf{v} is the velocity vector, $\Delta \mathbf{x}$ is the displacement vector and Δt is the time interval over which the displacement occurs. Note that we will use bold symbols, such as \mathbf{v} , for vectors and normal-weight symbols, such as v , for scalar quantities. Note also that the Greek letter Δ (capital delta) represents the change in a quantity. In the above expression, Eq. (1.1), for example, the change in the position of an object is its final position minus its initial position:

$$\Delta \mathbf{x} = \mathbf{x}_f - \mathbf{x}_i \tag{1.2}$$

Key concept:

Speed is the magnitude of the velocity. Speed is a scalar, and it does not have a direction.

Vectors in Examples

Many of the numerical examples presented will be one-dimensional, so the vector properties of many quantities are reduced to them being simply positive or negative relative to a certain direction, and thus are addressed in the process of constructing the problem.

The speed of an object is the distance travelled, divided by the time it took to travel that distance:

$$v = \frac{\Delta x}{\Delta t} \tag{1.3}$$

Note the differences between Eqns. (1.1) and (1.3). In Eq. (1.1), we use bold symbols for both the \mathbf{v} and the \mathbf{x} , indicating that we are referring to the velocity and the displacement in this equation. In Eq. (1.3), we use normal weight symbols, v and x , indicating that we are referring to the speed and distance in this equation.

Many textbooks use d to represent distances and \mathbf{d} to represent displacements rather than Δx and $\Delta \mathbf{x}$. We will often follow this practice when specific reference to the initial and final positions is not called for.

Consider Figure 1.1. A toy car is travelling in a circle around a toy race track and we wish to characterise its motion. If we are interested only in how fast the car is going, we could say it is travelling at 5 m s^{-1} ($= 18 \text{ km h}^{-1}$). Two cars travelling on the same circle will be perfectly well distinguished by noting the different lengths of the circle they traverse in the same time.

Now consider the situation illustrated in Figure 1.2. In this case, two cars approach the same intersection from different directions. In this situation, we might point out that one of the cars is travelling at 18 km h^{-1} , while the other is travelling at 12 km h^{-1} . However, this will not cover all of the differences between the two cars. Another important fact about them is that they are travelling in different directions. If we wanted to predict where these two cars would be in an hour (for example) it would not be enough to just use the magnitude of their velocity; we would also need to take into account their directions.




Figure 1.2 Linear motion in two directions. The cars are travelling at different speeds and in different directions.

1.4 Acceleration

In kinematics, the **acceleration**, \mathbf{a} , is a vector which quantifies changes in velocity. In everyday conversation we use the word acceleration to mean that the speed of an object is increasing. If an object was slowing down we would say that the object was decelerating. The concept of acceleration in physics is more general and applies to a larger set of situations. In physics, acceleration is defined to be the rate of change (in time) of the velocity:

$$\mathbf{a} = \frac{\Delta \mathbf{v}}{\Delta t} \tag{1.4}$$

This definition implies several characteristics of the acceleration:

1. Acceleration is a vector: it has a direction as well as a magnitude. The acceleration is the rate of change of the velocity, and velocity is a vector, therefore acceleration must also be a vector.
2. The acceleration vector of an object may point in the opposite direction to that object's velocity vector. When this happens, the object's velocity will decrease and may even reverse direction. This means that deceleration (slowing down) is just another acceleration, but in a particular direction.
3. An object may have an acceleration without its speed changing at all. Should the acceleration vector point in a direction perpendicular to the velocity vector, the direction of the velocity vector will change, but its length will not. A good example of this is when an object moves in a circle. In this case, the acceleration is always perpendicular to the velocity, so the speed of the object is constant, but its velocity is constantly changing.

To illustrate these ideas, consider a car which starts from rest ($v_i = 0$) and accelerates along a straight road so that its velocity increases by 2 m s^{-1} every second. The velocity of this car is illustrated at a series of later times in Figure 1.3.




Figure 1.3 A car accelerating at 2 m s^{-2} for 5 s.

Since the velocity changes by the same amount every second (2 m s^{-1}), the acceleration of the car is constant. The velocity is changing at a rate of 2 m s^{-1} per second, or 2 metres per second per second. This acceleration would normally be written as $a = 2 \text{ m s}^{-2}$ (or 2 m/s^2) to the right.

We can calculate the velocity at any time. Since we know how much the velocity increases every second and we also know that the car was initially stationary, we just multiply this rate by the time elapsed since the acceleration began, i.e. we use the equation

$$v = at \quad (1.5)$$

Note that this is a vector equation, so that the velocity is in the same direction as the acceleration. For the car in this example, which is accelerating in a straight line at a constant rate of 2 m s^{-2} from rest, after 4 s the speed is $v = at = 2 \text{ m s}^{-2} \times 4 \text{ s} = 8 \text{ m s}^{-1}$, and so on.

What if the car had not been at rest initially? Suppose that the car in the previous example had been travelling at a constant velocity of 5 m s^{-1} for some unspecified length of time, and then began to accelerate at 2 m s^{-2} . Figure 1.4 shows this car at a sequence of later times. Compare this figure with Figure 1.3.




Figure 1.4 A car accelerates from an initial velocity of 5 m s^{-1} with a constant acceleration of 2 m s^{-2} .

In one most important respect, the situation has not changed. The velocity of the car still increases at the same rate, so that the *change* in velocity after the acceleration begins is given by the equation

$$\Delta v = at \quad (1.6)$$

The difference between Eq. (1.5) and Eq. (1.6) is that we now explicitly recognise that it is the *change* in velocity that we are calculating. In the previous example we calculated the change in velocity, but since the car started at rest, the velocity of the car was the same as how much the velocity had increased. Since we now have a nonzero initial velocity, we must recognise that the change in velocity is the final velocity minus the initial velocity, so

$$v_f - v_i = at$$

Thus after 5 s we would find that $15 \text{ m s}^{-1} - 5 \text{ m s}^{-1} = 2 \text{ m s}^{-2} \times 5 \text{ s}$.

We are normally interested in calculating the final velocity, so we write the above equation in the form

$$v_f = v_i + at \quad (1.7)$$

We can use this equation to find the final velocity at any later time, so long as the acceleration has not changed.

Note that the velocity calculated using this formula is the *instantaneous* velocity of the car at that time. This is the velocity that you would read off the car's speedometer. We will discuss instantaneous and average velocities in more detail next.

1.5 Average Velocity or Speed

The discussion above allows us to calculate the instantaneous velocity of an object moving with constant acceleration. This is the velocity of the object at a particular instant of time. It is also often useful, when we are dealing with motion in a straight line, to use the average velocity of an object to solve problems. The average velocity is

$$v_{\text{av}} = \frac{\text{total displacement}}{\text{total time}} = \frac{d}{t} \quad (1.8)$$

If you drive a car from Dunedin, New Zealand, to Christchurch (370 km away to the north) in 5 h, your average speed is given by

$$v_{\text{av}} = \frac{d}{t} = \frac{370 \text{ km}}{5 \text{ h}} = 74 \text{ km h}^{-1}$$

It is important to realise that this calculation does not require any knowledge of the details of your trip. You may have travelled at a constant 74 km h^{-1} the whole way, or (more likely) you may have varied your speed significantly. You may even have stopped to look at the view and eat lunch for half an hour. These details are not needed for the calculation of the average velocity.

Without explicitly deriving it, an alternative expression for the average velocity vector, \mathbf{v}_{av} , of an object which is undergoing *constant acceleration* is given in Eq. (1.9).

$$\mathbf{v}_{\text{av}} = \frac{1}{2} (\mathbf{v}_i + \mathbf{v}_f) \quad (1.9)$$

where \mathbf{v}_i is the initial velocity and \mathbf{v}_f is the final velocity. Note that neither the magnitude or direction of the acceleration or the time period over which the acceleration acted are required here.

1.6 Change in Displacement Under Constant Acceleration

If an object is undergoing a constant acceleration \mathbf{a} , then the displacement, \mathbf{d} , occurring in some given time, t , is

$$\mathbf{d} = \mathbf{v}_{\text{av}} t = \frac{1}{2} (\mathbf{v}_i + \mathbf{v}_f) t$$

and from Eq. (1.7), we get

$$\mathbf{d} = \frac{1}{2} (\mathbf{v}_i + \mathbf{v}_i + \mathbf{a}t) t$$

and so

$$\mathbf{d} = \mathbf{v}_i t + \frac{1}{2} \mathbf{a}t^2 \quad (1.10)$$

If the object starts at rest (that is $\mathbf{v}_i = 0$) then this can be simplified to

$$\mathbf{d} = \frac{1}{2} \mathbf{a}t^2 \quad (1.11)$$

An Expression Independent of Time

The kinematic equations derived so far all include time. It is useful to have an alternative kinematic equation in which time does not feature. Such an equation can be derived from the kinematic equations already covered starting from of Eq. (1.8) in 1D:

$$d = v_{\text{av}} t$$

The average velocity $v_{\text{av}} = \frac{v_i + v_f}{2}$ as before, and rearranging Equation 1.7 to the form $t = \frac{v_f - v_i}{a}$ gives

$$d = \left(\frac{v_f + v_i}{2} \right) \left(\frac{v_f - v_i}{a} \right)$$

so

$$2ad = v_f^2 - v_i^2$$

with the useful result:

$$v_f^2 = v_i^2 + 2ad \quad (1.12)$$

We will now investigate an acceleration which is particularly important for the motion of objects near the surface of the Earth: the *acceleration due to gravity*.

Example 1.1 *Falling Ball (1D Kinematics)*

Problem: You drop a cricket ball from a 125 m high tower and it accelerates downwards at a rate of 10 m s^{-2} . How far will it fall in 5 s?

Solution: We can solve this problem in two different ways. We can find the average velocity of the ball over the first 5 s and use this average velocity to calculate a displacement, or we can calculate a displacement directly.

- (a) The acceleration of the ball is 10 m s^{-2} downwards and so the velocity increases by 10 m s^{-1} in the downwards direction every second. The initial velocity is 0 m s^{-1} so the final velocity must be

$$v_f = v_i + at = 0 \text{ m s}^{-1} + 10 \text{ m s}^{-2} \times 5 \text{ s} = 50 \text{ m s}^{-1}$$

$v_f = 50 \text{ m s}^{-1}$ in the downwards direction. The average velocity of the ball in the downwards direction is

$$v_{\text{av}} = \frac{0 \text{ m s}^{-1} + 50 \text{ m s}^{-1}}{2} = 25 \text{ m s}^{-1}$$

Using this average velocity, the distance that the cricket ball will fall in 5 s is:

$$d = v_{\text{av}}t = 25 \text{ m s}^{-1} \times 5 \text{ s} = 125 \text{ m}$$

- (b) The second technique uses Eq. (1.11) (since the initial velocity is zero). The change in displacement of the ball is

$$d = \frac{1}{2}at^2 = \frac{1}{2} \times 10 \text{ m s}^{-2} \times (5 \text{ s})^2 = 125 \text{ m}$$

which is the same answer as we found with the previous method.

1.7 The Acceleration Due to Gravity

Air Resistance

Actually, the acceleration is only the same in a vacuum. Objects falling in air are affected by air resistance which reduces the acceleration. Often this is small enough to be ignored.

Galileo found (and countless experiments since have also shown) that all objects falling freely towards the Earth have the same acceleration. (In order to see this effect, we must take into account the effect of air resistance when this is significant.) Thus every object in free fall close to the surface of the Earth has its downward speed increased by approximately 10 m s^{-1} in every second regardless of its mass. Galileo claimed that this was an experimental fact and is reported to have shown it by dropping two balls of unequal mass from the top of the Leaning Tower of Pisa. Later we will discuss the theoretical explanation for this experimental fact when we investigate the relationship between acceleration and the concept of *force*. The value of this constant acceleration is given by

$$g = 9.81 \text{ m s}^{-2} \approx 10 \text{ m s}^{-2} \quad (1.13)$$

This quantity, g , is called the **acceleration due to gravity**. This is the value at sea level on the surface of the Earth; the value will change with altitude.

Consider an object released from rest and accelerating in free fall. Assuming that the air resistance is negligible, we are able to calculate its velocity after 5 s:

$$v = gt = 9.81 \text{ m s}^{-2} \times 5.00 \text{ s} = 49.1 \text{ m s}^{-1}$$

The result is obtained when using the approximation $g = 10 \text{ m s}^{-2}$ is very similar:

$$v = gt = 10 \text{ m s}^{-2} \times 5 \text{ s} = 50 \text{ m s}^{-1}$$

Note that the mass of this object is not even mentioned in the original question, and that we have not considered the vectorial character of either the acceleration due to gravity or of the velocity achieved by this object after 5 seconds. These quantities are of course vectors, but their directions may be assumed to be towards the centre of the Earth and need not be considered in this problem. This is not always the case.

Precision and g

In mathematical examples where we wish to find an accurate result, we will use a value for g of 9.8 or 9.81 m s^{-2} . However, in some cases where we are more interested in the concepts and methods than the exact answer we may approximate the value as 10 m s^{-2} to keep things simpler.

Example 1.2 *A Ball Thrown Straight Up (I) (1D Kinematics)*

Problem: If you throw a cricket ball straight up at 12 m s^{-1} , how high will it go?

Solution: This problem is very similar to the previous one, except now the cricket ball has an initial velocity. This initial velocity is in the opposite direction to the acceleration due to gravity. This means that, at first, gravity will *reduce* the upward velocity of the cricket ball by 10 m s^{-1} every second. At some point the upward velocity of the cricket ball will have been reduced to zero – the cricket ball has stopped travelling up so it has reached its maximum height.

To calculate the maximum height that the ball reaches we need to find the time taken for the upward velocity to decrease to zero as this is also the time for the ball to reach its maximum height. We then calculate the average velocity and use this combined with the elapsed time to calculate the change in displacement of the ball.

For this problem we will define the upwards direction as positive, and define $d = 0 \text{ m}$ to be the height at which the ball was released. The change in velocity of the ball is $v_f - v_i = -12 \text{ m s}^{-1}$ (i.e. the velocity goes from $+12 \text{ m s}^{-1}$ initially to 0 m s^{-1} at its highest point). The time it takes the acceleration due to gravity (-9.81 m s^{-2} , as it points in the downwards direction) to cause this change in velocity is found using

$$\Delta v = gt$$

so

$$t = \frac{\Delta v}{g} = \frac{-12 \text{ m s}^{-1}}{-9.81 \text{ m s}^{-2}} = 1.2 \text{ s}$$

The average velocity is 6.0 m s^{-1} ($v_{\text{av}} = \frac{1}{2}(v_f + v_i) = \frac{1}{2}(12 \text{ m s}^{-1} + 0 \text{ m s}^{-1}) = 6.0 \text{ m s}^{-1}$) and using this we calculate the change in displacement of the cricket ball during the 1.2 seconds it takes to reach the maximum height,

$$d = v_{\text{av}}t = 6.0 \text{ m s}^{-1} \times 1.2 \text{ s} = 7.2 \text{ m}$$

The highest point the ball reaches is 7.2 m above the point at which it was released.

Example 1.3 *A Ball Thrown Straight Up (II) (1D Kinematics)*

Problem: How long does it take the ball in Example 1.2 to fall to its original position from its maximum height?

Solution: The ball reaches a maximum height of 7.2 m above the point at which it is thrown. At its maximum height it has a velocity of 0 m s^{-1} . The time taken to fall a distance of 7.2 m back to its original position can be found using Eq. (1.10):

$$d = v_i t + \frac{1}{2} a t^2$$

Since the initial velocity, v_i , is the velocity of the ball at its maximum height, i.e. $v_i = 0 \text{ m s}^{-1}$ this can be reduced to

$$d = \frac{1}{2} a t^2$$

The time in this equation, t , is the time the ball takes to fall back to its original position.

We rearrange this equation to solve for t where (using the same sign convention as in Example 1.2) $d = -7.2 \text{ m}$ and $a = g = -10 \text{ m s}^{-2}$

$$\begin{aligned} t &= \sqrt{\frac{2d}{g}} \\ &= \sqrt{\frac{2 \times -7.2 \text{ m}}{-9.81 \text{ m s}^{-2}}} \\ &= 1.2 \text{ s} \end{aligned}$$

Note that this is the same time it took the ball to reach its maximum height from the point at which it was initially thrown in Example 1.2. This is a useful general result. Projectile motion is symmetrical about the point of maximum height. It takes the same amount of time to reach the maximum height from a starting height as it does to get back to that height from the maximum height.

Example 1.4 A Ball Thrown Straight Up (III) (1D Kinematics)

Problem: What is the velocity of the ball in Examples 1.2 and 1.3 when it falls back to the height at which it was released?

Solution: As in the previous problem we can think of the ball starting at its maximum height and falling 7.2 m under the influence of gravity to its original height, which we found would take 1.2 s, the same as the time taken to reach its maximum height from the point at which it was released. The change in velocity over the 1.2 s seconds in which it is falling from its maximum height is

$$\Delta v = at = -10 \text{ m s}^{-2} \times 1.2 \text{ s} = -12 \text{ m s}^{-1}$$

The ball is travelling at the same speed at it was when released, but in the opposite direction! Again this is a useful general result we can apply to many kinematics problems without going through an extensive derivation.

Example 1.5 A Ball Thrown Straight Up (IV) (1D Kinematics)

Problem: If the ball in the Example 1.2 was released (travelling upwards) at a height of 1.2 m above the ground, what is the velocity of the ball just before it hits the ground?

Solution: We will first need to find the time it takes for the ball to hit the ground and then use $\Delta v = at$ to find the change in velocity, and hence the final velocity. Initially it is tempting to try to solve this problem directly using Eq. (1.10):

$$d = v_1 t + \frac{1}{2} at^2 \quad (1.10)$$

We know the change in position of the ball $d = -1.2 \text{ m}$ (as, in this case, the ball ends up 1.2 m below its starting point), the initial velocity of the ball ($v_1 = 12 \text{ m s}^{-1}$), and acceleration ($g = -9.81 \text{ m s}^{-2}$). In order to use this equation, however, we would need to solve for t , which would require solving a quadratic equation. Even using the shortcuts highlighted in the previous two examples does not allow us to avoid this quadratic as we still end up with two terms featuring t . If you're confident doing this, that is great. If you are not very confident at solving quadratic equations however, all is not lost.

We can simplify the problem by using the fact that we know a bit more about the situation than is apparent from the question. From Example 1.2, we know that the ball reaches a maximum height of 7.2 m above the point at which it was released. This means that the ball will reach a height of $7.2 + 1.2 = 8.4 \text{ m}$ above the ground. At this maximum height, the velocity of the ball is 0 m s^{-1} , and by ignoring the first part of the ball's motion, we can simplify Eq. (1.10) to $d = \frac{1}{2} at^2$, where d is the change in displacement of the ball as it moves from its maximum height to the ground, $d = -8.4 \text{ m}$, and $a = g = -9.81 \text{ m s}^{-2}$:

$$d = \frac{1}{2} at^2$$

$$t = \sqrt{\frac{2d}{g}} = \sqrt{\frac{2 \times -8.4 \text{ m}}{-9.81 \text{ m s}^{-2}}} = 1.3 \text{ s}$$

This gives a change in velocity of $\Delta v = gt = -9.81 \text{ m s}^{-2} \times 1.3 \text{ s} = -13 \text{ m s}^{-1}$, so the ball will be travelling at 13 m s^{-1} in the downwards direction as it hits the ground.

1.8 Independence of Motion in Two Dimensions

In the previous section, we considered objects which move vertically up and down. This means that in these cases the velocity vector is always parallel to the acceleration vector. These objects would go straight upward and fall straight downward. They will not move horizontally since there is no initial velocity in the horizontal direction, and no acceleration in the horizontal direction to cause a non-zero horizontal velocity to develop. What would happen if the initial velocity was not straight upward? What would have happened

if the initial velocity was at some angle to the vertical? This is the situation shown in Figure 1.5, using the example of a cricket ball launched upward at an angle.




Figure 1.5 The trajectory of a cricket ball initially launched upward at an angle to the vertical. The vertical and horizontal components of the ball's velocity are shown at a number of times in its trajectory.

In Figure 1.5, the cricket ball has a vertical velocity and a horizontal velocity and this results in a net velocity at an angle to the vertical. The acceleration due to gravity, however, acts only in the vertical direction, and changes only the vertical component of the velocity. The horizontal component of the velocity is initially 10 m s^{-1} and is still 10 m s^{-1} when the ball reaches its maximum height at 45 m after 3 s, and is 10 m s^{-1} when the ball reaches the ground again after a total of 6 s. There is no acceleration in the horizontal direction, so the velocity component in this direction *cannot change*.

The vertical component of the velocity is changed by the acceleration due to gravity. This can be seen in Figure 1.5 as well. In point of fact, the vertical component of the velocity behaves in exactly the same way as it did in the examples above. The vertical velocity is initially 30 m s^{-1} upward. After 3 s this has dropped to 0 m s^{-1} when the ball reaches its maximum height. The vertical velocity is again 30 m s^{-1} just before the ball hits the ground after 6 s, but now the velocity is in the downward direction.

This is what we mean when we say that the horizontal and vertical components of the velocity vector are independent. These components are acted on separately by the accelerations in those directions. An acceleration in the horizontal direction would not change the velocity component in the vertical direction. This means that when we are attempting to solve a kinematics problem in three dimensions, we may look at the components of the velocity and acceleration in a given direction independently of their components in the other two directions.

This effect may be seen in the following experiment. Suppose that we have two identi-




Figure 1.6 Two balls are released at the same time from the same height. One ball has a non-zero initial horizontal velocity. Both balls fall the same distance in the same time, and have the same vertical velocity at each point in time.

cal balls which are held on a platform in a darkened room. (The balls do not need to have the same mass for this experiment to work, but we will simplify the discussion by assuming that they do.) Now suppose that we drop one ball directly downward from a platform and at the same instant fire the other ball horizontally out from the same platform. As the balls fall, a strobe light flashes at regular intervals and the trajectory of the two balls is recorded on a camera with a very long exposure time. Figure 1.6 is an example of the sort of image that would be obtained from this experiment.

In this figure, we observe that the balls are at the same height at each interval, i.e. at each flash of the strobe light. This means that their vertical velocity components are the same at each time. Their horizontal velocity components are quite different, however. The ball which is simply dropped has no horizontal velocity component, whereas the horizontal velocity component of the other ball is constant.

Example 1.6 *Projectile Motion (2D Kinematics)*

Problem: A modern artist throws a bottle of paint towards the wall of a nearby building. The bottle leaves the artist's hand at a height of 2.0 m, a speed of 16 m s^{-1} , and at an angle of 30° above the horizontal. If the building is 15 m away:

- At what height H does the bottle hit the wall?
- At what velocity v is the bottle travelling as it hits the wall?

Solution:




Figure 1.7 An artist throws a bottle against a wall.

A good first step for this kind of problem is to draw a diagram like Figure 1.7. Remember that when dealing with 2D kinematics you can always separate out the horizontal and vertical motions. Note also that since we are given numerical quantities to two significant figures, we will use this level of numerical precision throughout the problem, i.e. we will use $g = -9.8 \text{ m s}^{-2}$.

In order to answer both of the questions, we will need to know how long after the bottle leaves the artist's hand it hits the wall. We can calculate this time by looking at the horizontal motion of the bottle. As the acceleration due to gravity is in the vertical direction only, we know that the horizontal velocity is constant and has a magnitude of $v_x = 16.0 \text{ m s}^{-1} \times \cos 30^\circ = 13.9 \text{ m s}^{-1}$.

The time it takes the bottle to travel the 15 m horizontally to the wall is

$$d_x = v_x t \quad \Rightarrow \quad \frac{d_x}{v_x} = \frac{15 \text{ m}}{13.9 \text{ m s}^{-1}} = 1.08 \text{ s}$$

- The difference in height between the bottle's initial height of 2.0 m and its final height of H is h_f . We can calculate this height by using the initial vertical velocity of the ball ($v_{yi} = 16 \text{ m s}^{-1} \times \sin 30^\circ = 8.0 \text{ m s}^{-1}$) and the fact that the ball is accelerating in the vertical direction at a rate of $a = g = -9.8 \text{ m s}^{-2}$.

$$\begin{aligned} h_f &= v_{yi} t + \frac{1}{2} g t^2 \\ &= 8.0 \text{ m s}^{-1} \times 1.08 \text{ s} + \frac{1}{2} \times (-9.8 \text{ m s}^{-2}) \times (1.08)^2 \\ &= 2.93 \text{ m} \end{aligned}$$

So the bottle must hit the wall a total of 4.9 m above the ground (giving solution to two significant figures).

- In order to find the final velocity of the bottle we will have to add together the vertical and horizontal components of the bottle's velocity as it hits the wall. We already know that the horizontal velocity of the bottle is constant as there is no acceleration in the horizontal direction, therefore the horizontal component of the final velocity is $v_x = 14 \text{ m s}^{-1}$. To find the final vertical velocity we can use $v_f = v_i + at$ where $v_i = v_{yi} = 8.0 \text{ m s}^{-1}$ and $a = g = -9.8 \text{ m s}^{-2}$.

$$\begin{aligned}
 v_{yf} &= v_{yi} + gt \\
 &= 8.0 \text{ m s}^{-1} + (-9.8 \text{ m s}^{-2}) \times 1.08 \text{ s} = -2.58 \text{ m s}^{-1}
 \end{aligned}$$

By the time the bottle has hit the wall, it has reached its maximum height (at which $v_y = 0 \text{ m s}^{-1}$) and has started moving back down, hence the negative vertical velocity.




Figure 1.8 The velocity vector components of the bottle as it hits the wall.

We can get the magnitude of the final velocity by vector addition of the two components v_x , and v_{yf} :

$$|v_f| = \sqrt{v_x^2 + v_{yf}^2} = \sqrt{(13.9 \text{ m s}^{-1})^2 + (-2.58 \text{ m s}^{-1})^2} = 14.14 \text{ m s}^{-1}$$

The direction in which the bottle is travelling can be found by using trigonometry.

$$\begin{aligned}
 \tan \theta_f &= \frac{v_{yf}}{v_x} \\
 \theta_f &= \tan^{-1} \left(\frac{v_{yf}}{v_x} \right) = \tan^{-1} \left(\frac{2.58 \text{ m s}^{-1}}{13.9 \text{ m s}^{-1}} \right) = 10.52^\circ
 \end{aligned}$$

So as the bottle hits the wall it is travelling at a speed of 14 m s^{-1} , 11° below the horizontal (to 2 s.f.).

1.9 Summary

Key Concepts

elapsed time (Δt) The time interval between two events.

distance (d or Δx) The length of a path between two spatial positions.

displacement (d or Δx) The vector equivalent of distance, which specifies the distance and direction of one point in space relative to another. It depends only on the initial and final spatial positions, and is independent of the path taken from one position to the other.

speed (v) A scalar measure of the rate of motion. The SI unit of speed is metres per second (m/s or m s^{-1}).

velocity (v) A vector measure of the rate of motion, which specifies both the magnitude and direction of the rate of motion.

acceleration (a) A measure of the rate of change of the velocity. Acceleration is a vector quantity. The SI units of acceleration are m/s^2 or m s^{-2} .

Equations

$$d = v_{av}t$$

$$\Delta v = at$$

$$v_{av} = \frac{1}{2}(v_i + v_f)$$

$$d = v_i t + \frac{1}{2}at^2$$

$$v_f^2 = v_i^2 + 2ad$$

1.10 Problems

When solving the following problems use $g = 9.81 \text{ m s}^{-2}$ unless otherwise specified.

1.1 Figure 1.9 shows five different paths between two end points. The initial position is indicated by i and the final position by f .




Figure 1.9

If each of the paths shown in the figure were travelled in exactly 100 s what are the:

- (a) change in displacement, d .
- (b) distance travelled, D .
- (c) average velocity, v .
- (d) average speed, s

for each journey?

1.2 What is the *average* velocity of the following:

- (a) A snail moves 30 cm north in 2 minutes.
- (b) An elevator is raised 18 m in 15 s.
- (c) An elevator is lowered 18 m in 15 s.
- (d) A lion runs 49 m west in 7 s.
- (e) A jogger runs a complete lap of a 1500 m oval track in 6 minutes.
- (f) A car passes a signpost travelling west at a speed of 10 m s^{-1} and 2 min later passes a second signpost while travelling north at 12 m s^{-1} . The second signpost is 300 m north-west of the first.
- (g) A car decelerates at a constant rate from a velocity of 40 m s^{-1} west to a velocity of 14 m s^{-1} west.

- (h) A train carriage that is initially moving at a velocity of 0.5 m s^{-1} north that accelerates uniformly to a velocity of 1.5 m s^{-1} south.
- (i) A train carriage that is initially moving at a velocity of 0.5 m s^{-1} south that accelerates uniformly to a velocity of 1.5 m s^{-1} south.

1.3 At $t = 0 \text{ s}$ a hovercraft is stationary and accelerating at a uniform rate of 0.8 m s^{-1} east.

What is the velocity of the hovercraft and its displacement with respect to its initial position at:

- (a) $t = 1 \text{ s}$
- (b) $t = 2 \text{ s}$
- (c) $t = 5 \text{ s}$
- (d) $t = 10 \text{ s}$
- (e) $t = 20 \text{ s}$
- (f) $t = 40 \text{ s}$

1.4 The position of a drag car along a 400 m long drag strip is recorded at regular intervals of 2 s. The results are shown in the table below and the finish line is due East from the start line.

	$t \text{ (s)}$	$d \text{ (m East)}$
A	0	0
B	2	16
C	4	64
D	6	144
E	8	256
F	10	400

- (a) What is the average velocity of the car between times A and B?
- (b) What is the average velocity of the car between times A and C?
- (c) What is the average velocity of the car between times A and D?
- (d) What is the average velocity of the car between times A and E?
- (e) What is the average velocity of the car between times A and F?

If we assume that the drag car accelerated at a uniform rate and started off stationary then:

- (f) What is the velocity of the car at time B?
- (g) What is the velocity of the car at time C?
- (h) What is the velocity of the car at time D?
- (i) What is the velocity of the car at time E?
- (j) What is the velocity of the car at time F?
- (k) What is the rate of acceleration of the drag car?

1.5 A car travels along the path shown in Figure 1.10. The car is travelling at a constant speed of 21 m s^{-1} . The radius of the track is 100 m.




Figure 1.10

- (a) What is the velocity of the car as it passes point A?
- (b) What is the velocity of the car as it passes point B?
- (c) What is the velocity of the car as it passes point C?
- (d) What is the velocity of the car as it passes point D?
- (e) What is the average velocity of the car between points A and B?
- (f) What is the average velocity of the car between points A and C?
- (g) What is the average velocity of the car between points A and D?
- (h) What is the average velocity of the car over a complete lap (from A to A)?

1.6 At $t = 0$ s a hovercraft has a velocity of 10 m s^{-1} west and is accelerating at a uniform rate of 0.8 m s^{-2} east.

What is the velocity of the hovercraft and its displacement with respect to its initial position at:

- $t = 1$ s
- $t = 2$ s
- $t = 5$ s
- $t = 10$ s
- $t = 20$ s
- $t = 40$ s

1.7 At $t = 0$ s a hovercraft has a velocity of 10 m s^{-1} west and is accelerating at a uniform rate of 0.8 m s^{-2} north.

What is the velocity of the hovercraft and its displacement with respect to its initial position at:

- $t = 1$ s
- $t = 2$ s
- $t = 5$ s
- $t = 10$ s
- $t = 20$ s
- $t = 40$ s

1.8 You are walking at a speed of 0.80 m s^{-1} . Define the direction that you are walking in as the positive direction. Your friend is walking at a speed of 1.2 m s^{-1} .

- If you and your friend are walking in the same direction, what is the relative velocity of your friend from your point of view?
- If you and your friend are walking in the opposite directions, what is the relative velocity of your friend from your point of view?

1.9 A blue car travelling at 80 km h^{-1} west is passed by a red car travelling in the same direction at 100 km h^{-1} .

- What is the velocity of the red car relative to the blue car?
- What is the velocity of the blue car relative to the red car?

A green car then passes the blue car. The velocity of the green car relative to the blue car is 6 km h^{-1} west.

- What is the velocity of the green car?
- What is the velocity of the green car relative to the red car?

1.10 Three cars are approaching an intersection as shown in Figure 1.11.




Figure 1.11

Car A is travelling at a speed of 45 km h^{-1} , Car B is travelling at a speed of 50 km h^{-1} , and Car C is travelling at a speed of 55 km h^{-1} .

- What is the relative velocity of car B from the point of view of car A at the instant shown in Figure 1.11?
- Does this relative velocity of car B from the point of view of car A change as the positions of the cars change?
- What is the relative velocity of car C from the point of view of car A at the instant shown in Figure 1.11?
- Does this relative velocity of car C from the point of view of car A change as the positions of the cars change?

1.11 A ball is dropped from a height of 10.0 m. The ball was stationary just before you let it go.

- What is the acceleration of the ball just after you let it go?
- What is the acceleration of the ball half-way towards the ground?
- What is the acceleration of the ball just before it hits the ground?

- How long does it take for the ball to hit the ground?
- What is the velocity of the ball just before it hits the ground?

1.12 You throw a ball straight up. The speed of the ball just after it leaves your hand is 14.1 m s^{-1} upwards. The ball rises, reaches its maximum height, then falls down. You catch the ball when it is the same height that you let it go.

- What is the acceleration of the ball just after you let it go?
- What is the acceleration of the ball when it is at its maximum height?
- What is the acceleration of the ball just before you catch it?
- How long does it take the ball to reach its maximum height?
- How long is the ball in the air before you catch it?
- What is the velocity of the ball at the instant when it is at its maximum height?
- What is the velocity of the ball at the instant just before you catch it?

1.13 A bullet is fired horizontally from a gun that is 1.5 m from the ground. The bullet travels at 1000 m s^{-1} and strikes a tree 150 m away. How far up the tree from the ground does the bullet hit? [Neglect air resistance.]

1.14 You are abducted by aliens who transport you to their home world in a galaxy far far away. Oddly, the only thing you can think of doing is measuring the acceleration due to gravity on this strange new world. You drop an alien paperweight from a height of 12 m and use an alien stopwatch to measure the interval of 1.36 s it takes the paperweight to hit the ground below. What is the acceleration due to gravity on the alien home world?

1.15 In a bid to escape from your alien captors you hurl your paperweight straight up towards the door switch on a space ship above you. If the switch is 25 m above you how fast does the paperweight need to leave your hand?

1.16 A cricket ball is hit for six and follows the path shown in Figure 1.12. The initial velocity of the cricket ball is 20 m s^{-1} at an angle of 45° above the horizontal.




Figure 1.12

The cricket ball is shown at 3 positions along its trajectory: Position A just after it is hit, Position B at the balls maximum

height, and Position C just before the ball lands. If the ball is hit at time $t = 0 \text{ s}$ then:

- (a) What is the component of v_A in the x -direction?
- (b) What is the component of v_A in the y -direction?
- (c) What is the acceleration of the ball at Position A?
- (d) What is the component of v_B in the x -direction?
- (e) What is the component of v_B in the y -direction?
- (f) What is the acceleration of the ball at Position B?
- (g) What is the component of v_C in the x -direction?

- (h) What is the component of v_C in the y -direction?
- (i) What is the acceleration of the ball at Position C?

1.17 The following question relates to the trajectory of the cricket ball in Problem 1.16.

- (a) What is the maximum height that the ball reaches?
- (b) How long is the ball in the air?
- (c) How far away from its initial position does the ball land?
- (d) What is the velocity of the ball just before it lands?

2

FORCE AND NEWTON'S LAWS OF MOTION

2.1	Introduction
2.2	The Concept of Force
2.3	Kinds of Force
2.4	Newtonian Gravity
2.5	Fictitious Forces
2.6	Summary
2.7	Problems

2.1 Introduction

We now have a clear and complete description of motion. This description relates the primary properties of motion – displacement, velocity, acceleration and elapsed time – to each other. The question ‘What causes a change in the motion of an object?’ amounts to asking ‘Where does acceleration come from?’. The answer to this question is deceptively simple, accelerations are caused by the application of forces. The purpose of this chapter is to explain clearly what a ‘force’ is in physics and how it can be used to solve problems relating to the motion of objects. When forces are included in the discussion of motion, it is called dynamics.

Key Objectives

- To understand the concept of force.
- To understand the relationship between force and motion.
- To be able to identify action–reaction pairs of forces.
- To understand normal, friction and tension forces.
- To be able to solve straightforward problems in dynamics.

2.2 The Concept of Force

In everyday conversation we use the word ‘force’ quite liberally. The Oxford English Dictionary gives a number of definitions of the noun ‘force’:

1. physical strength or energy accompanying action or movement.
2. (Physics) a measurable influence that causes something to move.
3. pressure to do something backed by the use of threat of violence.
4. influence or power.
5. a person or thing having influence: *a force for peace*.
6. an organised group of soldiers, police or workers.

The word is also used as a verb, as in sentences like: ‘She forced the committee to consider her application seriously.’ In physics, the word ‘force’ has a very precise meaning, and this is given by Newton’s laws. These laws define force by listing the essential properties of a force. If some phenomenon does not have all of these properties, then it is not a force. In this section, we will go through Newton’s laws and explain each of them in turn.

Force Vectors in Figures

In this chapter *force vectors* are represented as acting on the *centre of mass* of an object. It should be noted that other texts use differing conventions for contact force vectors in which a contact force is shown acting at the surface between two objects. The distinction between the two ways of representing contact force vectors is partially style, although when dealing with contact forces that may act to rotate an object the exact position of action of a force is important. Such cases will not be dealt with in this chapter and so the simpler standard in which all force vectors are shown to act at the centre of mass of an object has been used in the interests of clarity.

In later chapters, particularly Chapter 4, we will be considering forces that rotate an object and will draw force vectors for forces resulting from contact, at the point of contact.

Newton's First Law

In the list of Oxford English Dictionary definitions of force given above, the second item is closest to the definition used by physicists. A force causes a *change* in the motion of an object. For example, if an object is at rest, then we must apply a force to it to cause it to accelerate and to develop a non-zero velocity.

Key concept:
Newton's first law
 Any object continues at rest or at constant velocity (constant speed in a straight line) unless an external force acts on it.

Figure 2.1 illustrates the effect of a force on the motion of an object. This object is travelling in a straight line until an external force acts on it in a direction perpendicular to its motion. This causes the object to be deflected from its straight-line motion.




Figure 2.1 A pair of objects initially travel in the same direction. The lower object is subject to a force and its motion then deviates from a straight line while the force acts on it, but continues on in a new straight line after the force ceases to act.

Newton's Second Law

Key concept:
Newton's second law
 An external force gives an object an acceleration. The acceleration produced is proportional to the force applied, and the constant of proportionality is the **mass**.




Figure 2.2 If the force applied to an object does not change, but the mass of the object is doubled, then the acceleration is halved. If the mass does not change and the force is doubled, then the acceleration is doubled also.

Newton's second law can be summarised with the following equation:

$$F = ma \tag{2.1}$$

In this equation, a is the acceleration (in $m\ s^{-2}$) as usual, m is the mass (in kg), and F is the force (in N) (N = newton). The SI unit of force is the newton; one newton (1 N) is the force which would accelerate a 1 kg mass at $1\ m\ s^{-2}$ (i.e. would cause its velocity to increase by $1\ m\ s^{-1}$ in every second).

As is illustrated in Figure 2.2, if the mass of an object on which the force is applied does not change, but the force is doubled, then the acceleration of this object will also be doubled (see diagram to the right in Figure 2.2). If the applied force is not changed, but the mass of the object is doubled, then the acceleration will be halved (see centre diagram of Figure 2.2). For numerical examples of the relationship between mass, force and acceleration, see Table 2.1.