

Multidisciplinary Management of Pediatric Voice and Swallowing Disorders

J. Scott McMurray
Matthew R. Hoffman
Maia N. Braden
Editors

Multidisciplinary Management of Pediatric Voice and Swallowing Disorders

J. Scott McMurray
Matthew R. Hoffman • Maia N. Braden
Editors

Multidisciplinary Management of Pediatric Voice and Swallowing Disorders

 Springer

Editors

J. Scott McMurray
Department of Surgery, Division of
Otolaryngology-Head and Neck Surgery
University of Wisconsin School of
Medicine and Public Health
Madison, WI
USA

Matthew R. Hoffman
Department of Surgery, Division of
Otolaryngology-Head and Neck Surgery
University of Wisconsin School of
Medicine and Public Health
Madison, WI
USA

Maia N. Braden
Department of Surgery, Division of
Otolaryngology-Head and Neck Surgery
University of Wisconsin School of
Medicine and Public Health
Madison, WI
USA

ISBN 978-3-030-26190-0 ISBN 978-3-030-26191-7 (eBook)

<https://doi.org/10.1007/978-3-030-26191-7>

© Springer Nature Switzerland AG 2020

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors, and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

Gratitude with humility does not begin to express what is in my heart for so many people. I give thanks to my co-editors, Matt and Maia, who tirelessly worked this book to completion. I thank the many authors who contributed to this book. I give thanks to my other specialty partners without whom we could not give the care that we do. I give thanks to the patients and their families for allowing me to join their family and help point their children in a direction to achieve their full potential.

I give thanks to Chuck Ford and Diane Bless for taking a chance on me as I started my career. I give thanks to all the cottonoids around the globe who are supportive brothers and sisters, and especially to Mark, Dana, and Mike. I give thanks to Sally Shott, Chuck Myer, and J. Paul Willging. For opening opportunity, I give thanks to Chuck Kashima and Paul Flint. I give thanks to David Hanson and Jack Jiang. To Charles W. Cummings, you started it all for me in otolaryngology as the man I dreamt to try to be. To Robin Cotton, there are no words that can express my gratitude for what you have shared in so many different ways.

Without my mother and father and their unceasing and loving support and confidence, I would never have achieved what I have. To my sister, I give her thanks for not killing me as we were growing up.

I cannot give thanks enough for the blessings of my daughters, Grace and Margot, of whom I am so very proud.

Finally, and most importantly, I give thanks for the love of my wife, Jane. She is the rock in so many ways. She has been understanding and supportive. She has encouraged and

suggested different paths. She has set example for what to try to become. She has always been there for me. I love you and I always will.

I hope this book helps you in your journey, seeking knowledge and honing tools, so that you too may point your children patients in the direction to reach their fullest potential.

–J. Scott McMurray

Thank you to Jack Jiang, for introducing me to the larynx, and Tim McCulloch, for introducing me to the pharynx. Thank you to Scott McMurray, for fostering an interest in pediatric laryngology. Thank you to my daughter, Val, for her laugh, smile, joyful nature, and friendly disposition. Most importantly, thank you to my wife, Marisa, with whom I always have fun and whose support is unwavering, maternal instinct is natural, and patience while I worked on this book was unending; I love you.

–Matthew R. Hoffman

I give boundless thanks to my co-editors for their hard work and dedication to this project, and to the chapter authors for their time and expertise. I am also grateful to our patients and their families, from whom I am constantly learning. I am forever grateful to the amazing mentors I have had throughout my career. Thank you to Diane Bless, who introduced me to voice science and who continues to be a source of inspiration and encouragement; to Mary Sandage and Brian Petty, who taught me how to be a voice clinician; to Edie Hapner, who has always encouraged me to reach outside my comfort zone and try new things; and to Susan Thibeault, for her support in being a clinician researcher and research-based clinician.

Most importantly, I would like to thank my family. I am incredibly thankful to my kids, Finn and Marley, for their patience, encouragement, and willingness to spend time in coffee shops while I wrote and edited. And finally, I would like to thank my husband, Jason, my partner in all adventures, for his support, calm, and especially his sense of humor in this and all things. I love you.

–Maia N. Braden

Foreword

In over 100 years of collective practice, we have seen many changes in the ways voice and swallow problems are managed, largely due to new knowledge of structure and function, developments in technology, and transmission of information through a myriad of journals, e-records, and the Internet. The explosion of information has resulted in the development of subspecialties and evidence-based practices and the underlying knowledge that it is necessary to have a multidisciplinary team to best treat voice and swallowing disorders, a premise we expounded in our 1991 book, *Phonosurgery: Assessment and Surgical Management of Voice Disorders*.

The editors who conceived the current book have succeeded in creating a richly informative yet very readable comprehensive textbook on assessment and management of pediatric voice and swallowing disorders. All three editors are accomplished authors and clinicians practicing at the University of Wisconsin-Madison. J. Scott McMurray, MD, is the UWSMPH chief of Pediatric Otolaryngology; Maia N. Braden, MS, CCC-SLP, manages pediatric voice and swallowing problems at UWAF Children's Hospital; and Matthew R. Hoffman, MD, PhD, is a remarkably accomplished young senior resident with over 40 publications in major journals. They have selected some of the best and brightest clinicians in their respective fields to contribute chapters. Each chapter stands alone as a contribution to our mutual understanding of how to treat this pediatric population, and the combined chapters are likely to enhance the way clinicians practice.

As former teachers and colleagues of these authors, we are delighted that they articulate the importance of collaboration and communication in clinical practice. Both factors are particularly important in dealing with pediatric patients, where sensitivity, flexibility, and creativity promote optimal assessment and successful outcomes. Insofar as we teach by example, we are particularly gratified because these are principles we have embraced in our practice, teaching, lectures, and writings.

This book describes an effective multidisciplinary approach. We are introduced to clinician specialists skilled in care of pediatric patients with disorders of the aerodigestive tract. Along with the parents, otolaryngologists and speech-language pathologists usually play pivotal roles, often in collaboration with gastroenterologists, pulmonologists, and other specialists. Readers will obtain a wide range of these interacting roles so important in treating the pediatric population with voice and swallow disorders. The chapters articulate the importance of recognizing that children are not miniature adults and

that they need pediatric subspecialists who are cognizant of the disorders, issues, assessments, and treatments unique to the pediatric population.

Assessment of pediatric voice and swallowing problems are presented in this book by clinicians and scientists who participated in developing many currently used techniques. The techniques described cover the gamut of visualization; aerodynamic, acoustic, perceptual, quality-of-life, and manometric assessment procedures; as well as surgical and behavioral management practices. Notably, separate chapters are devoted to the problems typically seen in pediatric populations as well as chapters on issues dealing with singers and gender-affirming voice concerns. Relevant measures and concepts are explained with great clarity, making this book a valuable resource for persons who are beginners in the field as well as for veterans who have been practicing for years. The reader will be struck by an overarching strategic principle the authors embrace in assessment and management: *effective communication* with both patient and parent. Communicating with a young child can be challenging, especially with tasks that might cause discomfort or require active patient participation. This can cause the child – and possibly the parent – to become anxious and stressed. Successful examinations often require the examiner to be flexible by adjusting tasks to the patient’s ability. An informed parent will appear less anxious, which can be reassuring to an apprehensive child.

The section on intraoperative evaluation addresses the basics of operating room setup. Topics include essential equipment, solutions to laryngeal exposure problems, and discussion of advantages and limitations of inhalation vs total intravenous anesthesia. Two exciting new technologies are presented that provide alternative approaches to intraoperative assessment and treatment: (1) optical coherence tomography, which is used in diagnosis and depth assessment of vocal fold lesions like cysts, scars, and papilloma, and (2) microendoscopy via minithyrotomy, which is to provide subepithelial access to Reinke’s space.

We expect you will find many familiar things in this book to be reaffirming. You will also discover new things to enhance your practice. Hopefully, you will be struck by unexpected things that will stimulate your imagination and enrich your appreciation of what you do. We are so pleased to have written this foreword because the editors and contributing authors not only effectively articulate our vision of best practices, but most importantly, they advance the challenging field of treating pediatric patients with disorders of the aerodigestive tract.

Diane M. Bless, PhD
Department of Surgery, Division of Otolaryngology,
University of Wisconsin-Madison,
Madison, WI, USA

Charles N. Ford, MD
Department of Surgery, Division of Otolaryngology,
University of Wisconsin-Madison,
Madison, WI, USA

Preface

Kids are amazing. Taking care of kids is a gift and a joy. The triad of child, parent, and provider creates multiple layers of complexity that never cease to stimulate, and upon whom, a solid foundation can be formed for growth and development. Whereas the goal of adult and geriatric medicine is often to help a patient maintain their current but at times fleeting abilities, pediatric medicine focuses on pointing the patient in the right direction to achieve their own ultimate potential. With the correct assessment and the appropriate intervention, a tremendous future can be unleashed.

Disorders of the aerodigestive tract can impair a child's verbal communication and swallow function which can significantly hinder their personal and social development. As pediatric otolaryngologists and speech-language pathologists, we have the opportunity to help these children reach their full potential. Critical to achieving that aim is a multidisciplinary approach. This begins with collaboration between the otolaryngologist and speech-language pathologist. At our institution, this collaborative relationship was started and epitomized by Drs. Charles Ford and Diane Bless. They have served as leaders in our field and personal mentors to us.

Over the last two decades, our understanding of the numerous ways in which a child's voice or swallow can be altered has dramatically grown. The myriad and often complex interaction of acquired and congenital anomalies requires a detailed assessment with thoughtful attention before an accurate lifelong plan can be developed, discussed, and instituted. We know that not all children presenting to the otolaryngology clinic simply have nodules or reflux (though some do, as described in Chaps. 27 and 28) and that we can offer many of them more than observation and reassurance alone that symptoms will resolve with age.

As our understanding of voice and swallowing disorders has increased, the number of clinicians involved in the care of affected children has also increased. We now routinely work with gastroenterologists, pulmonologists, plastic surgeons, geneticists, physical therapists, and occupational therapists. Delivery of healthcare by this number of providers has been significantly enhanced by the creation of multidisciplinary pediatric aerodigestive clinics. At our institution, we call it the UW PACT or University of Wisconsin Pediatric Aerodigestive Care Team. Families can come from distances to see multiple specialists over the course of a single trip, attending clinics and undergoing operative assessments over a brief time span in order to better understand their child's disorder and receive a unified plan to help.

With this book, we sought to create a practical reference that would emphasize the collaborative relationships among clinicians that are critical to effective clinical care. Accordingly, most chapters are written by a physician and a speech-language pathologist. Furthermore, a straightforward framework for approaching, diagnosing, and managing each disorder is presented, including descriptions of relevant operative interventions. It is our hope that this could serve as a useful resource for not only otolaryngologists and speech-language pathologists but all members of the pediatric aerodigestive team and all other providers caring for children affected by voice and swallowing disorders.

We are very grateful to the authors who contributed to this book, without whose time and expertise, this would not have been possible. We are also grateful to Drs. Ford and Bless who provided valuable advice at the onset of the project and have provided valuable advice and mentorship throughout our careers.

Madison, WI, USA

J. Scott McMurray, MD
Matthew R. Hoffman, MD, PhD
Maia N. Braden, CCC-SLP

Contents

1 Pediatric Aerodigestive Programs: Role of the Core Team Members, Speech Language Pathology, Pulmonology, Gastroenterology, Otolaryngology, and Parent/Caregiver	1
J. Scott McMurray, Maia N. Braden, Matthew R. Hoffman, Vivek Balasubramaniam, and Dorota Walkiewicz	
2 Operative Evaluation of the Upper Aerodigestive Tract	13
Matthew R. Hoffman and J. Scott McMurray	
3 Anesthetic Considerations	21
Bridget Muldowney	
4 Perioperative Considerations After Pediatric Laryngeal Surgery	25
Maia N. Braden, Matthew R. Hoffman, and J. Scott McMurray	
5 Imaging Evaluation of the Upper Aerodigestive Tract	31
Tiffany Zens and Kara Gill	
6 Physiology of Voice Production	49
Matthew R. Hoffman, Maia N. Braden, and J. Scott McMurray	
7 Embryonic and Histologic Development of the Vocal Tract	63
Vidisha Mohad, Hailey Hirsch, and Susan L. Thibeault	
8 Nonlinear Acoustic Analysis of Voice Production	73
Hayley H. Raj, Austin J. Scholp, and Jack J. Jiang	
9 Clinical Approach to Acoustic Assessment	83
Elizabeth Heller Murray and GERALYN HARVEY WOODNORTH	
10 Aerodynamic Voice Assessment	89
Hunter Huth, Austin J. Scholp, and Jack J. Jiang	
11 Clinical Approach to Aerodynamic Assessment	99
Elizabeth Heller Murray and GERALYN HARVEY WOODNORTH	
12 Perceptual Evaluation of Voice	103
Maia N. Braden and Sarah D. M. Blakeslee	

13	Health-Related Quality of Life in Pediatric Dysphonia	109
	Shannon M. Theis and Nadine P. Connor	
14	Endoscopic Evaluation of the Pediatric Larynx.	119
	Rita R. Patel, Stephen D. Romeo, Jessica Van Beek-King, and Maia N. Braden	
15	Physiology of Normal Swallow.	135
	Corinne A. Jones	
16	Clinical Evaluation of Swallow	143
	Jesse D. Hoffmeister	
17	Videofluoroscopic Evaluation of the Swallow in Infants and Children	153
	Bryn K. Olson-Greb	
18	Pediatric Flexible Endoscopic Evaluation of Swallowing	167
	Jennifer L. Maybee, Bridget M. Harrington, and Jeremy D. Prager	
19	Manometric Evaluation of Pediatric Swallow	185
	Corinne A. Jones and Jesse D. Hoffmeister	
20	Quality of Life Assessment in Children with Feeding and Swallowing Disorders	195
	Pamela Dodrill and Hayley Henrikson Estrem	
21	Approach to Pediatric Voice Therapy	207
	Maia N. Braden	
22	Benign Mass Lesions.	213
	Matthew R. Hoffman, Maia N. Braden, and J. Scott McMurray	
23	Laryngopharyngeal Reflux	227
	Lauren Sowa, Holly Schmidt, and Mark E. Gerber	
24	Muscle Tension Dysphonia and Puberphonia	239
	Marshall E. Smith and Daniel R. Houtz	
25	Vocal Fold Mobility Impairment.	245
	Karen B. Zur, Kimberly Duffy, and Linda M. Carroll	
26	Laryngomalacia.	259
	Bethany R. Powers, Bryn K. Olson-Greb, and Jessica Van Beek-King	
27	Laryngeal Cleft	269
	Karthik Balakrishnan and Kari A. Krein	
28	Paradoxical Vocal Fold Motion	285
	Mary J. Sandage, Wynne Zhang, and Julina Ongkasuwan	

29	Vascular Anomalies	299
	Austin N. DeHart, Joana M. Mack, P. Spencer Lewis, Shelley E. Crary, and Gresham T. Richter	
30	Vocal Fold Scar	315
	Maria E. Powell and Bernard Rousseau	
31	Recurrent Respiratory Papillomatosis	325
	Brandyn Dunn, Kevin Huoh, and Hema Desai	
32	Glottic Web	339
	Kara D. Meister, April Johnson, and Douglas R. Sidell	
33	Syndromes and Congenital Anomalies	349
	Gregory Rice, Maia N. Braden, and J. Scott McMurray	
34	Esophageal Dysmotility	357
	Matthew R. Hoffman, Maia N. Braden, and J. Scott McMurray	
35	Cricopharyngeal Achalasia	369
	Tony Kille and Laurie Matzdorf	
36	Congenital Tracheal Anomalies: Complete Tracheal Rings, Tracheomalacia, and Vascular Compression	377
	Lyndy J. Wilcox, Claire Miller, and Michael J. Rutter	
37	Eosinophilic Esophagitis	399
	Ravi Sun, Robert Pesek, David Kawatu, Ashley O’Neill, and Gresham T. Richter	
38	Tracheoesophageal Fistula	413
	Scott M. Rickert	
39	Post-intubation Glottic Insufficiency	423
	Michael Shih, Danielle Devore, Sarah E. Hollas, and Julina Ongkasuwan	
40	Dysphonia After Laryngotracheal Reconstruction	431
	Mathieu Bergeron, Lyndy J. Wilcox, and Alessandro de Alarcon	
41	Non-cleft Velopharyngeal Insufficiency	445
	Katherine M. McConville and Catharine B. Garland	
42	The Young Aspiring Singer	465
	Debra Jean Phyland and Neil A. Vallance	
43	Gender-Affirming Voice	477
	Sarah L. Penzell	
44	Weird Wonders of the Larynx	485
	J. Scott McMurray and Matthew R. Hoffman	
	Index	497

Contributors

Karthik Balakrishnan, MD, MPH Division of Pediatric Otolaryngology, Department of Otorhinolaryngology and Mayo Clinic Children's Center, Mayo Clinic, Rochester, MN, USA

Vivek Balasubramaniam, MD Pediatric Pulmonology, Department of Pediatrics, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Mathieu Bergeron, BPharm, MD, FRCSC Department of Pediatric Otolaryngology, CHU Ste-Justine Pediatric Hospital, Montréal, QC, Canada

Sarah D. M. Blakeslee SSM Health, Department of Therapy Services, Speech Language Pathology, Madison, WI, USA

Maia N. Braden, MS, CCC-SLP UW Voice and Swallow Clinics, UW Health, American Family Children's Hospital, Madison, WI, USA

Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Linda M. Carroll, PhD, CCC-SLP, ASHA-F Children's Hospital of Philadelphia, Philadelphia, PA, USA

Nadine P. Connor, PhD Department of Communication Sciences and Disorders, University of Wisconsin-Madison, Madison, WI, USA

Department of Surgery, School of Medicine and Public Health, University of Wisconsin-Madison, Madison, WI, USA

Shelley E. Crary, MD Pediatric Hematology-Oncology, University of Arkansas for Medical Sciences, Arkansas Children's Hospital, Little Rock, AR, USA

Alessandro de Alarcon, MD, MPH Department of Otolaryngology Head and Neck Surgery, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

Austin N. DeHart, MD Department of Otolaryngology-Head and Neck Surgery, Division of Pediatric Otolaryngology, Arkansas Children's Hospital, University of Arkansas for Medical Sciences, Little Rock, AR, USA

Hema Desai, MS CCC-SLP, BCS-S Clinic of Speech Pathology, CHOC Children's Hospital, Orange, CA, USA

Danielle Devore, CCC-SLP Department of Speech, Language and Learning, Texas Children's Hospital, Houston, TX, USA

Pamela Dodrill, PhD, CCC-SLP Feeding and Developmental Therapy Team, Neonatal Intensive Care Unit, Brigham and Women's Hospital, Boston, MA, USA

Kimberly Duffy, MA, CCC-SLP Speech-Language Pathology, Children's Hospital of Philadelphia, The Center for Childhood Communication, Philadelphia, PA, USA

Brandyn Dunn, MD, MPH Department of Otolaryngology, University of California, Irvine, CA, USA

Catharine B. Garland, MD Department of Surgery, Division of Plastic and Reconstructive Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Mark E. Gerber, MD Otolaryngology Head and Neck Surgery, Phoenix Children's Hospital, Phoenix, AZ, USA

Kara Gill, MD Department of Radiology, Division of Pediatric Radiology, American Family Children's Hospital, Madison, WI, USA
University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Bridget M. Harrington, MA, CCC-SLP Department of Audiology, Speech and Learning, Children's Hospital Colorado, Aurora, CO, USA

Hayley Henrikson Estrem, PhD, RN School of Nursing, University of North Carolina Wilmington, Wilmington, NC, USA

Hailey Hirsch, CCC-SLP Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Matthew R. Hoffman, MD, PhD Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Jesse D. Hoffmeister, MS, CCC-SLP Department of Communication Sciences and Disorders, Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Sarah E. Hollas, MA, CCC-SLP Department of Speech, Language and Learning, Texas Children's Hospital, Houston, TX, USA

Daniel R. Houtz, MA, CCC-SLP Intermountain Voice and Swallowing Center, Intermountain Healthcare, Salt Lake City, UT, USA

Kevin Huoh, MD Department of Otolaryngology, CHOC Children's Hospital, Orange, CA, USA

Hunter Huth Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Jack J. Jiang, MD, PhD Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

April Johnson, MA, CCC-SLP Pediatric Voice and Swallowing Center, Department of Rehabilitation, Lucile Packard Children's Hospital Stanford, Palo Alto, CA, USA

Corinne A. Jones, PhD, CCC-SLP Neurology, Dell Medical School at the University of Texas-Austin, Austin, TX, USA

David Kawatu, MD Department of Pediatrics, Division of Pediatric Gastroenterology, Hepatology & Nutrition, Arkansas Children's Hospital, University of Arkansas for Medical Sciences, Little Rock, AR, USA

Tony Kille, MD Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin Hospitals and Clinics, Madison, WI, USA

Kari A. Krein, MS Division of Speech Pathology, Department of Neurology, Mayo Clinic, Rochester, MN, USA

P. Spencer Lewis, MD Pediatric Interventional Radiology, University of Arkansas for Medical Sciences, Arkansas Children's Hospital, Little Rock, AR, USA

Joana M. Mack, MD Pediatric Hematology-Oncology, University of Arkansas for Medical Sciences, Arkansas Children's Hospital, Little Rock, AR, USA

Laurie Matzdorf, MS, CCC-SLP Department of Speech Pathology, University of Wisconsin Hospitals and Clinics, Madison, WI, USA

Jennifer L. Maybee, OTR, MA, CCC-SLP Department of Audiology, Speech and Learning, Children's Hospital Colorado, Aurora, CO, USA

Katherine M. McConville, MA, CCC-SLP Department of Speech-Language Pathology, Michigan Medicine, University of Michigan, Ann Arbor, MI, USA

J. Scott McMurray, MD Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Kara D. Meister, MD Department of Otolaryngology-Head and Neck Surgery, Stanford University School of Medicine, Stanford, CA, USA
Pediatric Aerodigestive and Airway Reconstruction Center, Pediatric Voice and Swallowing Center, Stanford Children's Health, Lucile Packard Children's Hospital, Stanford, CA, USA

Claire Miller, PhD, MHA, CCC-SLP Aerodigestive and Esophageal Center, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

Vidisha Mohad, CCC-SLP Department of Communication Sciences and Disorders, Department of Surgery, Division of Otolaryngology – Head & Neck Surgery, Madison, WI, USA

Bridget Muldowney, MD Department of Anesthesiology, Division of Pediatric Anesthesiology, American Family Children's Hospital, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Elizabeth Heller Murray, PhD, CCC-SLP Department of Otolaryngology and Communication Enhancement, Boston Children's Hospital, Boston, MA, USA

Department of Speech, Language, and Hearing Sciences, Boston University, Boston, MA, USA

Ashley O'Neill, MS, CCC-SLP Department of Audiology and Speech Pathology, Arkansas Children's Hospital University of Arkansas for Medical Sciences, Little Rock, AR, USA

Bryn K. Olson-Greb, MS, CCC-SLP Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Julina Ongkasuwan, MD, FAAP, FACS Adult and Pediatric Laryngology, Department of Otolaryngology Head and Neck Surgery, Baylor College of Medicine, Houston, TX, USA

Rita R. Patel, PhD, CCC-SLP Department of Speech and Hearing Sciences, Indiana University, Bloomington, IN, USA

Sarah L. Penzell, MA, CCC-SLP Center for Audiology, Speech, Language, and Learning, Roxelyn and Richard Pepper Department of Communication Sciences and Disorders, Northwestern University, Evanston, IL, USA

Robert Pesek, MD Department of Pediatrics, Division of Allergy and Immunology, Arkansas Children's Hospital, University of Arkansas for Medical Sciences, Little Rock, AR, USA

Debra Jean Phyland, CPSP, PhD Voice Medicine Australia, East Melbourne, VIC, Australia

Department of Otolaryngology Head and Neck Surgery, School of Clinical Sciences at Monash Health, Monash University, Clayton, VIC, Australia

Maria E. Powell, PhD, CCC-SLP Department of Otolaryngology, Vanderbilt University Medical Center, Nashville, TN, USA

Bethany R. Powers, MD Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Jeremy D. Prager, MBA, MD Pediatric Otolaryngology, Children's Hospital Colorado; Department of Otolaryngology, University of Colorado School of Medicine, Aurora, CO, USA

Hayley H. Raj Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Gregory Rice, MD Department of Pediatrics, Division of Genetics and Metabolism, Medical Genetics Residency Program, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Gresham T. Richter, MD Department of Otolaryngology-Head and Neck Surgery, Division of Pediatric Otolaryngology, Arkansas Children's Hospital, University of Arkansas for Medical Sciences, Little Rock, AR, USA

Scott M. Rickert Department of Otolaryngology, Pediatrics, and Plastic Surgery, New York University Langone Health, New York, NY, USA

Stephen D. Romeo, MD Department of Surgery, Division of Otolaryngology-Head and Neck Surgery University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Bernard Rousseau, PhD, MMHC, CCC-SLP, ASHA Fellow Communication Science and Disorders, School of Health and Rehabilitation Sciences, University of Pittsburgh, Pittsburgh, PA, USA

Michael J. Rutter, MD, FRACS Department of Otolaryngology Head and Neck Surgery, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

Mary J. Sandage, PhD, CCC-SLP Department of Communication Disorders, Auburn University, Auburn, AL, USA

Holly Schmidt, MS CCC-SLP, BCS-S, MBA Department of Pediatric Speech Pathology, Lurie Children's Hospital of Chicago, Chicago, IL, USA

Austin J. Scholp, BS Departments of Surgery and Biomedical Engineering, Division of Otolaryngology-Head and Neck Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Michael Shih Department of Otolaryngology Head and Neck Surgery, Baylor College of Medicine, Houston, TX, USA

Douglas R. Sidell, MD Department of Otolaryngology-Head and Neck Surgery, Stanford University School of Medicine, Stanford, CA, USA

Pediatric Aerodigestive and Airway Reconstruction Center, Pediatric Voice and Swallowing Center, Stanford Children's Health, Lucile Packard Children's Hospital, Stanford, CA, USA

Marshall E. Smith, MD Division of Otolaryngology/Head and Neck Surgery, Primary Children's Hospital, University of Utah School of Medicine, Salt Lake City, UT, USA

Lauren Sowa, MD Department of Surgery, Section of Otolaryngology-Head and Neck Surgery, The University of Chicago Medicine, Chicago, IL, USA

Ravi Sun, BE Department of Otolaryngology-Head and Neck Surgery, University of Arkansas for Medical Sciences, Little Rock, AR, USA

Shannon M. Theis, PhD Department of Communication Disorders and Sciences, Rush University Medical Center, Chicago, IL, USA

Susan L. Thibeault, PhD, CCC-SLP Department of Surgery, Voice and Swallow Clinics, University of Wisconsin School of Medicine and Public Health, Division of Otolaryngology – Head & Neck Surgery, Madison, WI, USA

Neil A. Vallance, FRACS Melbourne Voice Analysis Centre, East Melbourne, VIC, Australia

Jessica Van Beek-King, MD Department of Surgery, Division of Otolaryngology-Head and Neck Surgery, Section of Pediatric Otolaryngology, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Dorota Walkiewicz, MD Pediatric Gastroenterology, Department of Pediatrics, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Lyndy J. Wilcox, MD Division of Pediatric Otolaryngology, Department of Otolaryngology-Head and Neck Surgery, Vanderbilt University, Nashville, TN, USA

Geralyn Harvey Woodnorth, MA, CCC-SLP Department of Otolaryngology and Communication Enhancement, Boston Children's Hospital, Boston, MA, USA

Tiffany Zens, MD Department of Surgery, Division of General Surgery, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

Wynne Zhang, BS BE Baylor College of Medicine, Houston, TX, USA

Karen B. Zur, MD Pediatric Voice Program, Center for Pediatric Airway Disorders, Philadelphia, PA, USA

Otorhinolaryngology – Head and Neck Surgery, Children's Hospital of Philadelphia, Philadelphia, PA, USA

Perelman School of Medicine at the University of Philadelphia, Philadelphia, PA, USA



Pediatric Aerodigestive Programs: Role of the Core Team Members, Speech Language Pathology, Pulmonology, Gastroenterology, Otolaryngology, and Parent/ Caregiver

J. Scott McMurray, Maia N. Braden,
Matthew R. Hoffman, Vivek Balasubramaniam,
and Dorota Walkiewicz

Overview

More and more, children who are medically complicated are surviving and flourishing as medicine advances and treatments improve. These medically complicated children pose specific and often difficult challenges as they present with congenital or acquired disorders of multiple organ systems that can impact breathing, swallowing, growth, and verbal communication. No other group of patients epitomize the need for an interdisciplinary team

approach with a core group of specialists than patients with aerodigestive disorders. The interest and formation of specialized aerodigestive programs have grown globally as their efficacy, efficiency, and economy have been recognized [1–6]. As more teams developed independently, the need for a consensus has been recognized regarding the types of patients and the typical disorders evaluated, the basic and minimum structure and function of the team, and the quality measures that should be followed. Boesch et al. [7] were the first to use the

J. S. McMurray (✉) · M. R. Hoffman
Department of Surgery, Division of Otolaryngology-
Head and Neck Surgery, University of Wisconsin
School of Medicine and Public Health,
Madison, WI, USA
e-mail: mcmurray@surgery.wisc.edu;
mrhoffman@wisc.edu

M. N. Braden
UW Voice and Swallow Clinics, UW Health,
American Family Children's Hospital,
Madison, WI, USA

Department of Surgery, Division of Otolaryngology-
Head and Neck Surgery, University of Wisconsin
School of Medicine and Public Health,
Madison, WI, USA
e-mail: braden@surgery.wisc.edu

V. Balasubramaniam
Pediatric Pulmonology, Department of Pediatrics,
University of Wisconsin School of Medicine and
Public Health, Madison, WI, USA
e-mail: balasubramaniam@surgery.wisc.edu

D. Walkiewicz
Pediatric Gastroenterology, Department of Pediatrics,
University of Wisconsin School of Medicine and
Public Health, Madison, WI, USA

Delphi method to obtain consensus about the structure and function of the aerodigestive program. The future of aerodigestive programs in general and the establishment of the aerodigestive society have been seminally shaped by this work, spawned by a desire to treat these complex patients well.

Based on the consensus developed by Boesch et al. [7], the following definition was developed for an aerodigestive patient. An aerodigestive patient is a child with a combination of multiple and interrelated congenital and/or acquired conditions affecting airway, breathing, feeding, swallowing, or growth that require a coordinated interdisciplinary diagnostic and therapeutic approach to achieve optimal outcomes. This includes (but is not limited to) structural and functional airway and upper gastrointestinal tract disease, lung disease because of congenital or developmental abnormality or injury, swallowing dysfunction, feeding problems, genetic diseases, and neurodevelopmental disability. Common conditions evaluated and treated through aerodigestive programs include structural or physiologic airway disease, congenital or acquired subglottic stenosis, chronic parenchymal lung disease, lung injury from aspiration or infection, gastroesophageal reflux, eosinophilic esophagitis, esophageal dysmotility or stricture, dysphagia, and behavioral feeding problems [8]. Piccone and Boesch [8] polled 50 programs in 31 states in the United States and compiled a list of common presenting conditions based on airway, pulmonary, gastrointestinal, feeding and swallowing, sleep, genetic, and neurologic disorders which are listed in Table 1.1.

There are a significant number of specialists that would be required to cover all of the possible conditions in children with aerodigestive disorders. Through consensus development by Boesch et al. [7], however, the list of essential core members whose input is required for all patients attending an aerodigestive program can be distilled to the following: care coordinator, nursing, speech language pathologist, pulmonologist, gastroenterologist, and otolaryngologist. An aerodigestive program should include these key players at a minimum.

Consensus was also achieved regarding the essential and defining functions and features of

an aerodigestive team evaluation [7]. For maximal efficiency and efficacy, the care cycle for an aerodigestive patient would involve the following work flow: consultation request and care coordination, pre-visit intake, team meeting, prescheduling appointments and procedures, shared clinic visit, combined endoscopy with a single anesthetic encounter, wrap-up visit with the family, summary document, and provision of follow-up care if needed.

Table 1.1 Common aerodigestive presenting conditions

Airway	Choanal atresia
	Laryngomalacia
	Glossoptosis
	Vocal fold paralysis
	Laryngotracheoesophageal cleft
	Stenosis: glottic, subglottic, tracheal, transglottis
	Tracheobronchomalacia
	Tracheoesophageal fistula
Pulmonary	Tracheostomy dependence
	Chronic lung disease of prematurity
	Diffuse lung disease
	Asthma
	Bronchiectasis: aspiration, ciliary dyskinesia, immunodeficiency, post-obstructive
Gastroenterology	Chronic respiratory failure
	Gastroesophageal reflux
	Eosinophilic esophagitis
	Esophageal structure
Feeding and swallowing	Failure to thrive
	Swallow incoordination
	Oral aversion
Sleep	Behavioral feeding problems
	Obstructive sleep apnea
	Central sleep apnea
Genetic	Hypoventilation
	Trisomy 21
	CHARGE association
	Pierre Robin sequence
	22q11 deletion
	VATER/VACTERL
	Craniofacial syndromes
Opitz syndrome	
Neurologic	Cri du chat
	Static encephalopathy
	Chiari malformation
	Hypotonia

Adapted from Piccone and Boesch [8], with permission

The typical aerodigestive program will see patients with a mix of medical and surgical needs. The interdisciplinary approach is important to effectively manage and plan the order of events leading to maximization of medical and surgical interventions and outcomes. Piccione et al. [8] also emphasized that there are several consistent structural elements of an aerodigestive program, namely, a (1) interdisciplinary medical and surgical team, (2) care coordination, (3) team meeting, and (4) combined endoscopy.

The team meeting is essential. This allows for distillation and review of historical events and prior tests. This information may be obtained through a telephone-based intake with caregivers and acquisition of previous medical records. This review will help to formulate a patient visit itinerary based on the team review and available best practice guidelines. It will help to ensure that a complete evaluation will be afforded in a short

and convenient time without needlessly repeating tests with the associated cost and risk. The telephone contact is also a great opportunity to council the family about expectations. The multidisciplinary visit can be overwhelming with the total number of interactions and the length of the overall day. Families are often thankful despite the long day once they realize the extent and expedience of the evaluation they will receive. The itinerary will include essential laboratory tests, radiographs, and swallow studies leading up to the clinic visit with the core provider team. The team visit confirms historical and physical findings and affirms the need and plan for the endoscopies and adjuvant tests requiring anesthesia. Piccione et al. [8] compiled the common aerodigestive diagnostic tests which are adapted in Table 1.2.

Each of the core specialists will bring their perspective and process for evaluating the chief

Table 1.2 Common aerodigestive diagnostic tests

Diagnostic modality	Strengths	Weaknesses
Chest radiograph	Identification of lower respiratory tract disease	Low sensitivity for bronchiectasis
	Low radiation	Limited ability to differentiate causes of lung disease
Chest CT	Distribution and severity of lung findings of various types	Increased radiation
	Differentiation between airway and parenchymal disease	May require sedation for good imaging
Upper GI series	Evaluation of anatomy: peristalsis, stricture, hernia, gastric outlet obstruction, malrotation	Does not evaluate reflux
Radionuclide reflux scan	Physiologic conditions May document aspiration from reflux	Limited sensitivity
Radionuclide salivagram	Assess for aspiration of saliva	Poor sensitivity
Radionuclide parotid scan	Assess function of major salivary glands	
FEES (fiber-optic endoscopic evaluation of swallowing)	Evaluate functional anatomy of swallowing	Blind to moment of pharyngeal swallowing and esophageal phase
	Evaluate airway protective reflexes	Not widely available
	Portable No radiation	
VFSS (videofluoroscopic swallowing study)	Evaluates all phases of swallowing	Radiation exposure
	Evaluates for aspiration	Limited anatomic evaluation
Microlaryngoscopy and rigid bronchoscopy	Superior optical resolution	Difficult access to peripheral airways
	Evaluation of the posterior larynx	Limited assessment of airway dynamics
	Access for instrumentation	Requires anesthesia

(continued)

Table 1.2 (continued)

Diagnostic modality	Strengths	Weaknesses
DISE (drug-induced sleep endoscopy)	Assessment of anatomic site of obstruction during sleep	Only an approximation of sleep state
		May miss REM specific obstruction
Flexible bronchoscopy with lavage	Evaluation of static and dynamic airway lesions, nasal-bronchial	Limited evaluation of posterior larynx
	Access to difficult and peripheral airways	Limited optical resolution
	Evaluation of airway inflammation and infections	Access for instrumentation Requires anesthesia
Esophagogastroduodenoscopy (EGD)	Evaluation of esophageal mucosal disease: acid and eosinophilic	Requires anesthesia
	Evaluation of esophageal, gastric, and duodenal anatomy	
	Obtain intestinal secretions	
	Evaluation of celiac disease	
Esophageal impedance	Identification and characterization of acid and nonacid reflux	Lack of normative data
	May identify dysmotility	Unclear relationship between impedance indices and extra-esophageal disease
Motility studies	Gold standard for dysmotility	Not widely available
Polysomnography (PSG)	Characterization of sleep disordered breathing and sleep architecture	Expensive and cumbersome
	Titration of respiratory support	Availability issues

Adapted from Piccione and Boesch [8], with permission

complaints and symptoms presented by the patient [8]. Although the group encounter, with all present for the clinic interview and the operative endoscopies, has been found to be the most efficient and efficacious, each provider brings unique and individual expertise. Each of the four core disciplines has overlap but also bring a unique role in the evaluation of these complex patients. The role of each core discipline will be outlined in this chapter.

Role of the Speech Language Pathologist

The multidisciplinary voice, swallow, and aerodigestive team can provide comprehensive, patient-centered and evidence-based care for children and adolescents with a variety of disorders impacting voice, swallow, and upper airway. Multidisciplinary team management of aerodi-

gestive disorders in children has been found to be more cost-effective and has better outcomes than stand-alone care [6]. In voice disorders, the model of speech language pathologist and otolaryngologist working together in evaluation and treatment has been well established since the 1980s and became more common in pediatric voice around a decade later. The strength of these teams lies in both the diverse knowledge and skills of the team members and their ability to work collaboratively to evaluate and treat the patient. The speech language pathologist specializing in these areas provides a focused set of knowledge and skills for these patients. We can provide evaluation of structure, function, and behavior of upper airway as they relate to voice, swallow, and breathing. In many cases we can also provide behavioral therapy to change voice, breathing, and swallow function, provide education, and provide compensatory strategies when needed. We have specialized knowledge of laryngeal structure and

function; the mechanics of voice, breathing, and swallowing; and neurologic controls of voice, swallow, and breathing. We provide valuable contributions with our in-depth understanding of behavior change. On any medical team, but especially with complex children, we do not operate in a vacuum and collaborate with surgical and medical personnel in both evaluation and treatment. According to the American Speech-Language-Hearing Association's scope of practice statement, "SLPs share responsibility with other professionals for creating a collaborative culture. Collaboration requires joint communication and shared decision making among all members of the team, including the individual and family, to accomplish improved service delivery and functional outcomes for the individuals served" [9].

Evaluation of Swallow

Dysphagia is relatively common in children. A rate of 0.9% was found in children aged 3–17 [10], and incidence is higher in certain medically complex populations including those with cerebral palsy and craniofacial syndromes [11–13]. There has been a marked increase in diagnoses of dysphagia in the pediatric hospitalized population, from 0.08% in 1997 to 0.41% in 2012 [14]. While exact reasons for this are not clear, it is often attributed both to increased survival rates of extremely preterm infants and improved diagnosis of swallowing disorders. Often the SLP is the first contact a child with dysphagia has with the multidisciplinary team. Children may be referred directly to us for a swallow evaluation or for treatment of feeding or swallowing disorders, or we may care for the child in the NICU from birth. We have the benefit of being able to spend the time to get a comprehensive history and provide ongoing assessment in therapy sessions. The SLP has several methods of evaluating swallowing, including the clinical swallowing evaluation, flexible endoscopic evaluation of swallowing, and video fluoroscopic swallowing study, as well as less frequently used measures including manometry. These may be used in combination depending on the needs of the patient. According to

ASHA, the role of the SLP in evaluation includes participating in determining the appropriateness of instrumental evaluation and follow-up, diagnosing pediatric oral and pharyngeal swallowing disorders, making appropriate referrals to other disciplines, and recommending a safe swallowing and feeding plan [9].

We require the expertise of others when evaluating and planning treatment beyond swallow recommendations for structural and functional deficits impacting swallowing, including (but not limited to) neurologic impairments, cerebral palsy (CP), sensory deficits, tracheoesophageal fistula, laryngeal cleft, acid reflux, esophageal dysmotility, laryngeal mobility impairment, and neurologic disorders.

Treatment of Dysphagia

SLPs on the multidisciplinary team as well as our colleagues working in more general outpatient settings, birth to three, and schools provide feeding and swallowing therapy to habilitate or rehabilitate swallowing and progress feeding skills. Feeding is defined as any aspect of eating or drinking and includes preparing food or liquid for intake, sucking or chewing, and swallowing [15]. Swallowing specifically refers to the complex processes involved in transporting solids, liquids, or saliva from the mouth to the digestive tract while maintaining airway protection [15]. Speech language pathologists are involved in evaluation and treatment of both.

A detailed description of all forms of feeding and swallowing therapy is beyond the scope of this chapter. Approaches to swallowing treatment may include positioning changes, changes in viscosity of bolus, changes in flow rate of bolus, maneuvers, sensory stimulation techniques, oral motor treatments, pacing, and cue-based feeding [15].

Evaluation of Voice

Incidence estimates of pediatric dysphonia are varied, ranging from 1.4% [10] to 26% [16]. Dysphonia rates in children are likely increasing

for some of the same reasons dysphagia rates are increasing, and children are presenting with more complex etiologies of voice disorders, beyond benign lesions. As survival rates of children born extremely preterm, or with complex tracheal or laryngeal anomalies, increase, rates of hoarseness and the complexity of children seen in the voice clinic will increase. For example, 38% of a sample of children born extremely preterm were found to have moderate-severe dysphonia at school age, with only 6% having normal voice [17]. Speech language pathologists often work in collaboration with an ENT in diagnosis and evaluation of voice disorders. According to the ASHA scope of practice, SLPs can perform a comprehensive voice evaluation which includes clinical and instrumental evaluation, assess normal or abnormal vocal function, describe voice quality and function, diagnose a voice disorder, refer to appropriate professionals to provide diagnosis of the underlying cause of the voice disorder (e.g., nodules as a cause of dysphonia), and make referrals to other professionals for other medical, surgical, or behavioral evaluation [9]. We can perform perceptual, acoustic, and aerodynamic evaluation of vocal function. We can also visualize the larynx using rigid or flexible endoscopy with stroboscopy, as well as high-speed digital video imaging of the larynx, and provide skilled interpretation of structure and function based on this. We do not diagnose lesions but can identify and describe the parameters of laryngeal function based on these evaluations and contribute to planning treatment, whether it be behavioral, surgical, or a combination of the two. The voice evaluation is also an important time to assess for stimulability for change based on therapeutic probes.

Treatment of Dysphonia

SLPs in a voice clinic and in other settings plan and deliver skilled treatment to optimize vocal function given the current anatomy, provide pre- and postoperative therapy, and provide therapy to change ingrained vocal functional behaviors. A detailed discussion of the types of voice therapy provided is beyond the scope of this chapter but

can be found in other sections of this book and in these and other resources [18–23].

Evaluation and Treatment of Breathing Disorders

Speech language pathologists are also experts in evaluation and management of laryngeal breathing disorders such as paradoxical vocal fold motion disorder, exercise-induced laryngomalacia, and chronic cough [9]. We can behaviorally and endoscopically evaluate laryngeal, pharyngeal, and respiratory function during breathing and provide interventions related to laryngeal sensitivity and control as well as optimizing respiratory coordination [24–28].

Conclusion

The benefits of working as a part of a multidisciplinary team cannot be overstated, for both clinician and patient. We are able to evaluate based on our areas of expertise and then discuss with other team members based on the findings of their specialized evaluations, providing optimal treatment for patients.

Role of the Otolaryngologist

As a specialist of disorders of the upper aerodigestive tract, the otolaryngologist shares the pathway to both the lungs and the gastrointestinal tract. This unique perspective positions them to be able to relate to both the pulmonologist and gastroenterologist. Working in conjunction with the speech language pathologist, the otolaryngologist can help assess the anatomy and physiologic function of the upper aerodigestive tract. Medical treatments of aerodigestive disorders in children are likely made in conjunction with the gastroenterology and pulmonology regarding reflux, inflammation, or infection. Dynamic surgical interventions of the airway may be suggested after functional assessment in collaboration with the speech pathologist. The typical

aerodigestive problems evaluated by the otolaryngologist can be seen in Table 1.1. The role of the otolaryngologist centers primarily on evaluation of airway surgical issues and aspiration [8].

The otolaryngologist should elicit history specific to obstructive sleep apnea, voice and swallowing disorders, recurrent infection, previous surgical history, or instrumentation of the airway. An assessment of possible congenital or genetic disorders is also essential. Growth and weight gain curves are helpful to assess potential feeding or breathing problems.

The otolaryngologist can offer expertise in office and operative endoscopy to evaluate function and anatomy. Identifying sites of abnormal anatomy, obstruction, or function of the upper aerodigestive tract is the prime modality offered. Expertise in nasopharyngoscopy in the awake patient facilitates anatomical and functional evaluation for airway obstruction, voice disorders, and swallowing dysfunction. Expertise with flexible endoscopy with the patient in a state mimicking sleep is also essential for identifying sites of obstruction causing obstructive sleep apnea. Drug-induced sleep endoscopy (DISE) protocols continue to be developed to bring the patient as close to a state of true sleep as possible [29]. Typical dense general anesthesia for airway endoscopy changes muscular tone and can change the site of obstruction that occurs during this type of sleep and can misdirect the clinician during the evaluation. Accurately identifying the true site of obstruction during normal sleep is required to allow for successful surgical management of obstructive sleep apnea.

Plain radiographs of the airway are often helpful during the assessment of the airway and may be ordered by the otolaryngologist. PA and lateral plain radiographs of the upper aerodigestive tract and chest are often helpful. This affords an assessment of the upper airway and trachea. Obstruction from adenoidal hypertrophy (Fig. 1.1), subglottic narrowing, vascular compression, or complete tracheal rings may first be identified or suspected in these films which are easy to obtain. This will help prepare the team for operative endoscopy and prevent unsuspecting catastrophe in cases such as complete tracheal rings.



Fig. 1.1 Large adenoidal pad filling the nasopharynx in this patient

Other radiographs and studies such as swallow studies, esophagram, upper GI CT and CT chest angiography, or MRI of the head or chest may also be of interest and are discussed by the entire team to determine utility and need.

Rigid airway endoscopy with its superior optics should also be the forte of the otolaryngologist. It offers superior static visualization, sizing, and intervention. Figure 1.2 is an example of severe laryngomalacia seen during direct laryngoscopy. The otolaryngologist member of the aerodigestive program should be strongly versed and capable with endoscopy of the upper aerodigestive tract.

As the surgical representative of the four core members, the otolaryngologist can also offer surgical or procedural correction for certain disorders. Boesch et al. [7] polled aerodigestive programs regarding the types of procedures requiring proficiency by otolaryngologists in aerodigestive programs for open and endoscopic airway reconstruction. Proficiency should be maintained in the following interventional categories: (1) open or endoscopic procedures that directly increase the diameter of the cartilaginous skeleton of the airway, (2) endoscopic treatment of airway obstruction, (3) surgical procedures to treat aspiration, (4) surgical procedures to improve voice, (5) tracheostomy, and (6) foreign body removal.

One of the most rewarding goals encountered in many aerodigestive patients is the relief




Fig. 1.2 (a, b) Operative direct laryngoscopy in a 6-week-old child with signs and symptoms of progressive noisy breathing and retractions since birth. Her symptoms were worse during activity of crying and feeding. She has a diagnosis of laryngomalacia. The infantile larynx is curled and obstructed during inspiration with the supraglottic structures

of tracheotomy dependence. Some patients will require airway reconstruction for the tracheotomy to be successfully removed. As stated previously, successful airway reconstruction requires a multidisciplinary approach [30] with sufficient preoperative evaluation of all airway lesions and non-airway diagnoses, appropriate patient selection, appropriate reconstructive technique, staging and timing, and effective patient optimization [8]. An understanding of the surgical approach for airway reconstruction in the context of static and dynamic lesions is essential and must include an appreciation of the interrelatedness with other comorbidities [8].

Working together, the otolaryngologist can bring expertise in diagnostic airway evaluation and surgical airway interventions to maximize patient outcomes through the aerodigestive team.

Role of the Pulmonologist

The role of the pulmonologist in the aerodigestive team is to provide complementary anatomic airway evaluation. Flexible bronchoscopy allows for a better dynamic assessment of the trachea and bronchus. It also allows for a more distal airway assessment and affords a superior bronchoalveolar lavage to help with culture and identifying any inflammatory markers. The pulmonologist is also integral in diagnosing and optimizing respiratory comorbidities prior to airway reconstruction and assisting in postoperative management. Pulmonologists may evaluate for and manage lung injury due to aspiration, active infectious or inflammatory lung disease, impaired airway clearance, interstitial lung disease, asthma, sleep apnea, dynamic airway lesions, and respiratory muscle weakness [8]. The pulmonologist may also make recommendations regarding radiographs, CT chest and CT angiography, as well as pulmonary function tests that may aid in the evaluation.

A list of recommended procedures that a pediatric pulmonologist should be able to provide at an aerodigestive center has been established by Boesch et al. using the Delphi method among a number of aerodigestive programs [7]. This includes bronchoscopy with bronchoalveolar lavage, balloon dilation, sleep state bronchoscopy, biopsy, foreign body removal, and identification of tracheoesophageal fistula.

Role of the Gastroenterologist

The role of the gastroenterologist is to evaluate growth and nutrition and gastrointestinal barriers to safe and adequate feeding and to diagnose and manage esophageal and other gastrointestinal disorders that may present as aerodigestive symptoms (Table 1.1) [8]. They are also essential in managing gastrointestinal disorders that may complicate airway reconstruction. The array of possible diagnoses includes laryngopharyngeal or gastroesophageal reflux, acidic or eosinophilic esophagitis, reflux aspiration, esophageal dysmotility, esophageal stricture, rumination, gastritis, and malabsorption [8].

The gastroenterologist will rely on history and physical findings, upper GI series, swallow studies, EGD (esophagogastroduodenoscopy) with biopsy, pH probe monitoring, impedance manometry, and specific blood work. Procedures which can be performed by the pediatric gastroenterologist on the aerodigestive team include esophagogastroduodenoscopy with biopsy, dilation, cautery, and placement of percutaneous endoscopic gastrostomy (PEG) or gastrojejunostomy (GJ) tubes [7].

Role of the Parent/Caregiver

When working with children, we cannot exclude the role of the parent(s) or caregiver(s) in evaluation and management. DeCivita and Dobkin [31] describe the therapeutic triad that exists among the medical team, the child, and the parent/caregiver. All have shared decision-making, and the impact of the disorder as well as the treatment burden needs to be considered. In interviewing parents of children with dysphonia, Connor and colleagues found that parents reported concerns with social and emotional issues related to their voice, as well as concerns with being understood and fitting in and concerns about comments from others [32]. Parents of children undergoing voice therapy for dysphonia have discussed concerns about social and emotional outcomes, academic and career success for their children, and the impact that their voice has on peer relationships [33]. At the same time, attendance and adherence to voice therapy depend on the parent, as they are usually the ones scheduling and bringing the child to therapy, as well as helping with and monitoring home practice [33]. With infants, very young children, or older children with cognitive or communication impairments, the parent is the only one able to provide a history and description of problems. Parents bear the financial burden, responsibility for transportation, and responsibility for carrying over medical team recommendations at home. They are responsible for making challenging decisions about their child's care, with varying levels of medical knowledge and experience.

Feeding and swallowing are particularly emotionally loaded areas for families. Parents of chil-

dren born with cleft lip and palate have reported feeling that their ability to feed their baby is linked to their competency as a parent [34]. When feeding modifications are recommended, the parents are primarily responsible for thickening liquids, providing positioning and pacing, and ensuring adequate oral intake. Parents have described feeding difficulties with children as a journey lasting from birth and discussed the impact it has on daily life, from the ability to leave the house to schedule activities, and the need to plan ahead extensively [35]. Parents of children with failure to thrive (FTT) [36] described not feeling heard by medical professionals, conversely feeling nurtured by others; feeling comparisons; being afraid; doing what needed to be done.

When partnering with parents, the medical team needs to take into account their values, their understanding of the medical issues facing their child, and their resources and abilities to cope. In many cases, it is easy for a medical team to see themselves as providing a service for the child, rather than partnering with the child and family, especially when treating conditions that require multiple interventions, whether they are medical, surgical, or behavioral.

Conclusion

The multidisciplinary approach to children with aerodigestive disorders is rewarding. Through this approach, efficient, cost-effective, patient-centered, family-focused, and consistent care can be delivered. The number of new diagnoses and the speed to diagnosis increase. The overall cost to diagnosis decreases, and the efficiency in the OR increases, freeing up more OR time for other unique procedures [6]. These complex and medically fragile patients deserve our best and concerted care to help them reach their maximal potential.

References

1. Galligan MM, Bamat TW, Hogan AK, Piccione J. The pediatric aerodigestive center as a tertiary care-based medical home: a proposed model. *Curr Probl Pediatr Adolesc Health Care*. 2018;48(4):104–10. <https://doi.org/10.1016/j.cppeds.2018.03.006>.

2. Mudd PA, Silva AL, Callicott SS, Bauman NM. Cost analysis of a multidisciplinary aerodigestive clinic: are such clinics financially feasible? *Ann Otol Rhinol Laryngol.* 2017;126(5):401–6. <https://doi.org/10.1177/0003489417699420>.
3. Appachi S, Banas A, Feinberg L, et al. Association of enrollment in an aerodigestive clinic with reduced hospital stay for children with special health care needs. *JAMA Otolaryngol Head Neck Surg.* 2017;143(11):1117–21. <https://doi.org/10.1001/jamaoto.2017.1743>.
4. Garcia JA, Mistry B, Hardy S, et al. Time-driven activity-based costing to estimate cost of care at multidisciplinary aerodigestive centers. *Laryngoscope.* 2017;127(9):2152–8. <https://doi.org/10.1002/lary.26354>.
5. Rotsides JM, Krakovsky GM, Pillai DK, et al. Is a multidisciplinary aerodigestive clinic more effective at treating recalcitrant aerodigestive complaints than a single specialist? *Ann Otol Rhinol Laryngol.* 2017;126(7):537–43. <https://doi.org/10.1177/0003489417708579>.
6. Skinner ML, Lee SK, Collaco JM, Lefton-Greif MA, Hoch J, Au Yeung KJ. Financial and health impacts of multidisciplinary aerodigestive care. *Otolaryngol Neck Surg.* 2016;154(6):1064–7. <https://doi.org/10.1177/0194599816637830>.
7. Miller CK, Wooten CT, DeBoer EM, et al. Structure and functions of pediatric aerodigestive programs: a consensus statement. *Pediatrics.* 2018;141(3):e20171701. <https://doi.org/10.1542/peds.2017-1701>.
8. Piccione J, Boesch RP. The multidisciplinary approach to pediatric aerodigestive disorders. *Curr Probl Pediatr Adolesc Health Care.* 2018;48(3):66–70. <https://doi.org/10.1016/j.cppeds.2018.01.002>.
9. American Speech Language and Hearing Association. Scope of practice in speech-language pathology. 2016. www.asha.org/policy.
10. Bhattacharyya N. The prevalence of pediatric voice and swallowing problems in the United States. *Laryngoscope.* 2015;125(3):746–50. <https://doi.org/10.1002/lary.24931>.
11. Benfer KA, Weir KA, Bell KL, Ware RS, Davies PSW, Boyd RN. Oropharyngeal dysphagia in preschool children with cerebral palsy: oral phase impairments. *Res Dev Disabil.* 2014;35(12):3469–81. <https://doi.org/10.1016/j.ridd.2014.08.029>.
12. Calis EAC, Veugelers R, Sheppard JJ, Tibboel D, Evenhuis HM, Penning C. Dysphagia in children with severe generalized cerebral palsy and intellectual disability. *Dev Med Child Neurol.* 2008;50(8):625–30. <https://doi.org/10.1111/j.1469-8749.2008.03047.x>.
13. de Vries IAC, Breugem CC, van der Heul AMB, Eijkemans MJC, Kon M, van der Molen ABM. Prevalence of feeding disorders in children with cleft palate only: a retrospective study. *Clin Oral Investig.* 2013; <https://doi.org/10.1007/s00784-013-1117-x>.
14. Horton J, Atwood C, Gnagi S, Teufel R, Clemmens C. Temporal trends of pediatric dysphagia in hospitalized patients. *Dysphagia.* 2018;33(5):655–61. <https://doi.org/10.1007/s00455-018-9884-9>.
15. Arvedson J, Brodsky L, editors. Pediatric swallowing and feeding: assessment and management. 2nd ed. Albany: Singular Publishing; 2002.
16. Carding PN, Roulstone S, Northstone K, ALSPAC Study Team. The prevalence of childhood dysphonia: a cross-sectional study. *J Voice.* 2006;20(4):623–30. <https://doi.org/10.1016/j.jvoice.2005.07.004>.
17. French N, Kelly R, Vijayasekaran S, et al. Voice abnormalities at school age in children born extremely preterm. *Pediatrics.* 2013;131(3):e733–9. <https://doi.org/10.1542/peds.2012-0817>.
18. Ramig LO, Verdolini K. Treatment efficacy: voice disorders. *J Speech Lang Hear Res.* 1998;41(1):S101–16. <http://www.ncbi.nlm.nih.gov/pubmed/9493749>
19. Verdolini K, Druker DG, Palmer PM, Samawi H. Laryngeal adduction in resonant voice. *J Voice.* 1998;12(3):315–27. [https://doi.org/10.1016/S0892-1997\(98\)80021-0](https://doi.org/10.1016/S0892-1997(98)80021-0).
20. Verdolini Abbott K. Some guiding principles in emerging models of voice therapy for children. *Semin Speech Lang.* 2013;34(02):080–93. <https://doi.org/10.1055/s-0033-1342979>.
21. Stemple JC, Lee L, D’Amico B, Pickup B. Efficacy of vocal function exercises as a method of improving voice production. *J Voice.* 1994;8(3):271–8. [https://doi.org/10.1016/S0892-1997\(05\)80299-1](https://doi.org/10.1016/S0892-1997(05)80299-1).
22. Stemple JC, Hapner ER, editors. Voice therapy: clinical case studies. 4th ed. San Diego: Plural Publishing; 2014.
23. Hartnick C, Ballif C, De Guzman V, et al. Indirect vs direct voice therapy for children with vocal nodules: a randomized clinical trial. *JAMA Otolaryngol Head Neck Surg.* 2018;144(2):156–62. <https://doi.org/10.1001/jamaoto.2017.2618>.
24. Sandage MJ, Zelazny SK. Paradoxical vocal fold motion in children and adolescents. *Lang Speech Hear Serv Sch.* 2004;35(4):353–62. [https://doi.org/10.1044/0161-1461\(2004\)034](https://doi.org/10.1044/0161-1461(2004)034).
25. Sandage M. Sniffs, gasps, and coughs: irritable larynx syndrome across the lifespan. *ASHA Lead.* 2006;11(9):16–21.
26. Gallena SJK, Solomon NP, Johnson AT, Vossoughi J, Tian W. The effect of exercise on respiratory resistance in athletes with and without paradoxical vocal fold motion disorder. *Am J Speech Lang Pathol.* 2015;24(3):470–9. https://doi.org/10.1044/2015_AJSLP-14-0110.
27. Mathers-Schmidt BA. Paradoxical vocal fold motion: a tutorial on a complex disorder and the speech-language pathologist’s role. *Am J Speech Lang Pathol.* 2001;10(2):111–25. [https://doi.org/10.1044/1058-0360\(2001\)012](https://doi.org/10.1044/1058-0360(2001)012).
28. Murry T, Sapienza C. The role of voice therapy in the management of paradoxical vocal fold motion,

- chronic cough, and laryngospasm. *Otolaryngol Clin N Am.* 2010;43(1):73–83. <https://doi.org/10.1016/j.otc.2009.11.004>.
29. Friedman NR, Parikh SR, Ishman SL, et al. The current state of pediatric drug-induced sleep endoscopy. *Laryngoscope.* 2017;127(1):266–72. <https://doi.org/10.1002/lary.26091>.
30. Meier JD, White DR. Multisystem disease and pediatric laryngotracheal reconstruction. *Otolaryngol Clin N Am.* 2012;45(3):643–51. <https://doi.org/10.1016/j.otc.2012.03.004>.
31. De Civita M, Dobkin PL. Pediatric adherence as a multidimensional and dynamic construct, involving a triadic partnership. *J Pediatr Psychol.* 2004;29(3):157–69.
32. Connor NP, Cohen SB, Theis SM, Thibeault SL, Heatley DG, Bless DM. Attitudes of children with dysphonia. *J Voice Off J Voice Found.* 2008;22(2):197–209.
33. Braden MN, Leer EV, McConville K, Blakeslee SDM. Patient, parent, and speech-language pathologists' perceptions of pediatric voice therapy through interviews. *Am J Speech Lang Pathol.* 2018;27(4):1385–404.
34. Lindberg N, Berglund AL. Mothers' experiences of feeding babies born with cleft lip and palate. *Scand J Caring Sci.* 2014;28(1):66–73.
35. Estrem HH, Pados BF, Thoyre S, Knafk K, McComish C, Park J. Concept of pediatric feeding problems from the parent perspective. *MCN Am J Matern Child Nurs.* 2016;41(4):212–20.
36. Thomlinson EH. The lived experience of families of children who are failing to thrive. *J Adv Nurs.* 2002;39(6):537–45.