

NEPHROLOGY

Lecture Notes



Edited by
Surjit Tarafdar

LN

WILEY Blackwell

Lecture Notes: Nephrology

Welcome to the series home page for the *Lecture Notes* books for medical students and junior doctors.

- The series covers over 35 subjects providing the concise core knowledge required by medical students and junior doctors
- Each book written by an expert teacher in note form, with bullet lists, illustrations, and summary boxes
- A panel of medical students reviews each text to ensure that the coverage is exactly right for its audience
- Ideal as course textbooks or revision aids

To view the full range of other titles in the Lecture Notes series, scan this QR code:



Lecture Notes: Nephrology

A Comprehensive Guide to Renal Medicine

Edited by

Dr Surjit Tarafdar

General Physician and Nephrologist
Blacktown and Mt Druitt Hospital
Blacktown
New South Wales, Sydney, Australia

Conjoint Senior Lecturer
Department of Medicine
Western Sydney University
Sydney, Australia

WILEY Blackwell

This edition first published 2020

© 2020 by John Wiley & Sons Ltd

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by law. Advice on how to obtain permission to reuse material from this title is available at <http://www.wiley.com/go/permissions>.

The right of Surjit Tarafdar to be identified as the author of editorial material in this work has been asserted in accordance with law.

Registered Office(s)

John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, USA

John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Office

9600 Garsington Road, Oxford, OX4 2DQ, UK

For details of our global editorial offices, customer services, and more information about Wiley products visit us at www.wiley.com.

Wiley also publishes its books in a variety of electronic formats and by print-on-demand. Some content that appears in standard print versions of this book may not be available in other formats.

Limit of Liability/Disclaimer of Warranty

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting scientific method, diagnosis, or treatment by physicians for any particular patient. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. While the publisher and authors have used their best efforts in preparing this work, they make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives, written sales materials or promotional statements for this work. The fact that an organization, website, or product is referred to in this work as a citation and/or potential source of further information does not mean that the publisher and authors endorse the information or services the organization, website, or product may provide or recommendations it may make. This work is sold with the understanding that the publisher is not engaged in rendering professional services. The advice and strategies contained herein may not be suitable for your situation. You should consult with a specialist where appropriate. Further, readers should be aware that websites listed in this work may have changed or disappeared between when this work was written and when it is read. Neither the publisher nor authors shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

Library of Congress Cataloging-in-Publication Data

Names: Tarafdar, Surjit, editor.

Title: Lecture notes: Nephrology : a comprehensive guide to renal medicine /
edited by Dr. Surjit Tarafdar.

Other titles: Nephrology

Description: Hoboken, NJ : Wiley Blackwell, 2020. | Includes
bibliographical references and index.

Identifiers: LCCN 2019035286 (print) | LCCN 2019035287 (ebook) | ISBN
9781119058045 (paperback) | ISBN 9781119058083 (adobe pdf) | ISBN
9781119058113 (epub)

Subjects: MESH: Kidney Diseases

Classification: LCC RC918.N43 (print) | LCC RC918.N43 (ebook) | NLM WJ
300 | DDC 616.6/1–dc23

LC record available at <https://lccn.loc.gov/2019035286>

LC ebook record available at <https://lccn.loc.gov/2019035287>

Cover Design: Wiley

Cover Image: © PIXOLOGICSTUDIO/Science Photo Library/Getty Images

Set in 8.5/11pt Utopia by SPi Global, Pondicherry, India

To my family for their patience and support. Their belief in my ability helped me to go through those days when I was overwhelmed and exhausted.

Contents

List of Contributors, ix

Foreword, xi

Acknowledgements, xiii

- 1 Clinical Implications of Renal Physiology, 1
- 2 Investigation of Renal Diseases, 21
- 3 Disorders of the Renal Tubules Leading to Disturbances of Acid–Base and Potassium, 41
- 4 Renal Stones, 53
- 5 Kidney Cancer, 67
- 6 Acute Kidney Injury, 77
- 7 Chronic Kidney Disease, 89
- 8 Nephrotic Syndrome, 111
- 9 Glomerulonephritis, 127
- 10 Renal Vasculitis and Lupus Nephritis, 143

- 11 Renovascular Hypertension, Pregnancy-Related Hypertension, and Thrombotic Microangiopathies, 163
- 12 Hereditary and Familial Renal Diseases, 181
- 13 Hepatorenal Syndrome, 195
- 14 Infections of the Kidney, 203
- 15 Diabetic Nephropathy, 215
- 16 Kidney Disease in Myeloma and Other Monoclonal Gammopathies, 231
- 17 Tubulointerstitial Diseases, 245
- 18 Maintenance Haemodialysis, 257
- 19 Peritoneal Dialysis: Principles, Indications, and Common Complications, 277
- 20 Renal Transplantation, 295
- Index, 321

List of Contributors

Edwin Anand

Assistant Professor
Jacobs School of Medicine & Biomedical Sciences
The State University of New York at Buffalo
Buffalo, NY, USA

Anthea Anantharajah

Clinical Immunologist / Immunopathologist
The Canberra Hospital, Garran, ACT;
Clinical Lecturer, Australian National University
Acton, Australia

Dipankar Bhattacharjee

MD(Calcutta), FRCP (Edin), FRCP (Lon), FCPS (B'Desh)
Consultant physician and Nephrologist, NHS, UK
Honorary Senior Lecturer (Ex) United Kingdom

Robert Carroll

Consultant Nephrologist, Central Northern Adelaide
Renal and Transplantation Service, Central Adelaide
LHN, SA Health, Adelaide, Australia

Philip Clayton

Consultant Nephrologist, Central Northern Adelaide
Renal and Transplantation Service, Central Adelaide
LHN, SA Health
Adelaide, Australia

Alexander Gilbert

Advances Trainee in Nephrology, Royal Prince Alfred
Hospital, Sydney, Australia

Pankaj Hari

Professor, Department of Paediatrics, All India
Institute of Medical Sciences, New Delhi, India

Yena Hye

Department of Geriatric Medicine
Nepean-Blue Mountains Hospital
Penrith, Australia

Rajini Jayaballa

MBChB, FRACP, Staff Specialist in Diabetes and
Endocrinology, Western Sydney Diabetes, Blacktown &
Mount Druitt Hospitals, Sydney, Australia

R. Jayasurya

Consultant Nephrologist, Apollo Gleneagles Hospital
Kolkata, India

Lukas Kairaitis

Head of Renal Services, Blacktown & Mount Druitt
Hospitals, Sydney; Sub-Dean, Blacktown & Mount
Druitt Clinical School, School of Medicine
Western Sydney University
Sydney, Australia

Amanda Mather

Staff Specialist Renal Physician, Royal North Shore
Hospital
Sydney, Australia

Stephen McDonald

Consultant Nephrologist, Central Northern Adelaide
Renal and Transplantation service, Central Adelaide
LHN, SA Health, Adelaide;
South Australia Executive Officer, ANZDATA Registry,
SA Health and Medical Research Institute;
Clinical Professor, Adelaide Medical School,
University of Adelaide, Adelaide, Australia

Mark McLean

Staff Specialist in Diabetes and Endocrinology,
Blacktown & Mount Druitt Hospitals, Westmead
Hospital, Sydney; Conjoint Professor of Medicine –
Western Sydney University
Sydney, Australia

Karumathil Murali

FRACP, MD Senior Staff specialist in Renal Medicine
Wollongong Hospital; Clinical Associate Professor
Graduate School of Medicine, University of
Wollongong
Wollongong, Australia

M.K. Phanish

Consultant Nephrologist, Renal Services; Lead
consultant,
Living Donor Kidney Transplantation, Epsom;
St.Helier University Hospitals NHS Trust; Lead
consultant, Diabetes-Renal Services, CUH, Epsom, UK

Priyamvada P.S.

Additional Professor, Department of Nephrology
JIPMER, Puducherry, India

Richard J. Quigg

Arthur M. Morris Professor of Medicine and
Biomedical Informatics Chief,
Division of Nephrology Jacobs School of Medicine &
Biomedical Sciences
The State University of New York, The State University
of New York, Buffalo, NY, USA

Gowri Raman

Current Palliative Care Advanced Trainee, University
of Newcastle
Australia

Wayne Rankin

Chemical Pathologist, SA Pathology and Clinical
Senior Lecturer, Discipline of Medicine, University of
Adelaide
Adelaide, Australia

Vinay Sakhuja

Emeritus Professor, Department of Nephrology, Post
Graduate Institute of Medical Education & Research
Chandigarh, India

Sarah So

Renal Advanced Trainee
Westmead and Blacktown Hospitals
Sydney, Australia

Kamal Sud

Department Head of Renal Medicine, Clinical
Associate Professor
University of Sydney - Nepean Clinical School
Nepean Hospital
Kingswood, Australia

Sanjay Swaminathan

Professor, Clinical Dean, Blacktown & Mount Druitt
Clinical School, Western Sydney University; Clinical
Immunologist, Westmead and Blacktown Hospital
Sydney, Australia

Surjit Tarafdar

Consultant Nephrologist and General Physician,
Blacktown and Mount Druitt Hospital
Blacktown, NSW, Sydney; Conjoint Senior Lecturer
Department of Medicine, Western Sydney University
Sydney, Australia

Muh Geot Wong MBBS, PhD, FRACP

Staff Specialist, Department of Renal Medicine, Royal
North Shore Hospital; Senior Clinical Lecturer
University of Sydney
Sydney, Australia

Foreword

The title 'Lecture Notes' undersells the comprehensive nature of this textbook, which fills a largely vacant niche in nephrology bookshelves, between lecture notes for medical students and junior trainees usually covering specific aspects of nephrology, and comprehensive speciality tomes targeted at those requiring specialized knowledge about any or all aspects of nephrology. *Lecture Notes: Nephrology* is aimed specifically at early-stage physician trainees who in the Australian system are preparing to undertake the basic physician trainee examination, or more senior trainees who have completed that exam, but are seeking expert knowledge as a trainee in general medicine, or are just starting specialized training in nephrology. However, its reach is much broader than that and it will appeal to anyone wanting to understand nephrology at a level greater than that found in general medical textbooks, including more advanced nephrology trainees, subspecialty nephrologists seeking knowledge beyond their subspecialty, general physicians and nephrologists wishing to brush up on all aspects of nephrology, and teachers of nephrology at any level. Whereas its reach is broad, it does not pretend to cater to those seeking highly specialized knowledge of particular aspects of nephrology. It is full of practical, contemporary information useful for day-to-day application on any aspect and at any level of renal diagnosis, investigation and management. Even the nephrologist who has been practising his or her craft for many years will find exciting new gems. In reviewing the book I found myself looking forward to reading each of the chapters at a more leisurely pace, and to applying its new knowledge for teaching and clinical practise.

The authors of *Lecture Notes: Nephrology* range from international, national and local experts on

particular aspects of renal science and its clinical application, to advanced trainees in nephrology who under the supervision of senior colleagues need to have this level of expertise and knowledge for their day-to-day supervision of nephrology patients. Each chapter has been reviewed thoroughly by other authors and content experts, and achieves high standards of readability, immediacy and clinical applicability. The chapters cover all aspects of nephrology, including physiology appropriately linked to clinical disease, general nephrology, dialysis and transplantation. Of particular relevance to those preparing for nephrology examinations, and others wishing to test how well they have assimilated the book's new knowledge, is a comprehensive set of multiple choice questions, which have been written by those well placed to do so; that is, trainees preparing for or having recently completed general physician and nephrology examinations.

Surjit Tarafdar and his colleagues have produced a renal textbook of high utility, which will occupy a place of honour on nephrology bookshelves.



David CH Harris
AM, MD (USyd), BS, FRACP
Professor of Medicine,
University of Sydney
Director of Nephrology & Dialysis,
Western Sydney Renal Service
Past-President, International Society of Nephrology

Acknowledgements

I am indebted to the following groups and individuals for their contributions:

- The Histopathology Department at ICPMR, Westmead Hospital, Sydney, Australia.
- Dr Pei Dai, Advanced Trainee in Immunology at Westmead Hospital, Sydney, for the Immunofluorescence slides.
- Dr Simon Gruenewald, Consultant Nuclear Physician at Westmead Public Hospital, Sydney, and Dr Basim Alqutawneh, Consultant Radiologist at Blacktown Hospital, Sydney, for the radiographs and imaging material.
- Professor Jeremy Chapman, Associate Professor Gopala Rangan and Dr Brian Nankivell from the Department of Renal Medicine of Westmead Hospital, Sydney, and Professor Ravindra Prabhu A, Department of Nephrology at Kasturba Medical College, Manipal, India, for reviewing portions of the book.
- Dr Alexander Gilbert, renal Advanced Trainee at Prince Alfred Hospital, Sydney, for proofreading almost the entire book in addition to co-writing Chapter 14.
- Lashnika Bandaranayake and Thilan Pasanjith Subasinghe, the wonderful team of fifth-year medical students from Western Sydney University, who helped with most of the diagrams and illustrations.
- To the lovely librarians at Blacktown Hospital, Sydney, for letting me stay back in the Peter Zelas Library many a day past closing time.
- To the following junior doctors and registrars who reviewed the book and provided valuable feedback on the language and content:
 - Dr Shrikar Tummala
 - Dr Tu Hao Tran
 - Dr Hannah Jsu
 - Dr Shaun Khanna
 - Dr Sumreen Nawaz
 - Dr Ye Min Kuang
 - Dr Marlies Pinzon
 - Dr Serge Geara
- To Yogalakshmi Mohanakrishnan, the project editor from the publisher's office, for her patience, guidance and prompt responses.
- Finally, I am extremely thankful to Professor David Harris from Westmead Hospital, Sydney. Despite being the President of the International Society of Nephrology at a time when the World Congress of Nephrology was being held in Australia, Professor Harris found time to review the book and provide pearls of wisdom which added immensely to the quality of the finished work.

Clinical Implications of Renal Physiology

Surjit Tarafdar

Summary

- Besides maintaining a stable acid base, electrolyte, and fluid status of the body, kidneys also have an important endocrine role in producing and secreting 1,25-dihydroxycholecalciferol (calcitriol), renin, and erythropoietin
- More than 98% of water in the filtered urine is reabsorbed in the tubules; 90–95% of water is reabsorbed as it follows sodium (Na^+), which is avidly reabsorbed by the Na^+ -deficient epithelial cells, except in the collecting duct (CD), where 5–10% of water is reabsorbed (independent of Na^+) under the direct influence of vasopressin or anti-diuretic hormone (ADH)
- The tubular epithelial cells constantly lose three Na^+ and gain two potassium (K^+) ions from the basolateral membrane (due to the Na^+ - K^+ -ATPase), which keeps these cells deficient in Na^+
- The countercurrent mechanism, which is dependent on the impermeability of the thick ascending limb (TAL) to water, leads to the creation of an increasing osmotic gradient from the cortex to the deeper medulla, which in turn enables ADH to reabsorb water in the CD
- Aldosterone helps in Na^+ reabsorption in the CD and also leads (directly) to K^+ and (indirectly) to hydrogen (H^+) secretion into urine
- All diuretics act by inhibiting tubular reabsorption of Na^+
- Formation of urine begins in the glomerular capillaries where the filtrate has to cross the three filtration layers: endothelium, glomerular basement membrane (GBM) and the foot processes of the podocytes; all these three layers are negatively charged and hence repel anionic proteins like albumin
- Nephrotic syndrome, which is marked by abnormally increased filtration of plasma proteins in the urine, may be due to widening of the pores in the three filtration layers, but is almost always associated with loss of negative charges in these layers
- Nephritis, which is due to glomerular inflammation, is characterized by haematuria with red blood cell (RBC) casts and dysmorphic RBCs, some degree of oliguria, hypertension, and reduction in glomerular filtration rate
- Goodpasture's disease, which is characterized by antibodies against subtype of type IV collagen, can lead to nephritis and haemoptysis, as this particular collagen is found predominantly in the GBM and alveolar membranes of the lungs
- Familial hypocalciuric hypercalcaemia, which manifests with hypercalcaemia, characteristically low urinary calcium and normal to high serum parathyroid hormone (PTH) level, is due to mutation in the calcium-sensing receptor (found in the kidney and parathyroid gland) leading to abnormally increased renal

reabsorption of calcium and inappropriate secretion of PTH

- Distal renal tubular acidosis (type 1 RTA) is due to an inability of the distal tubules to excrete H^+ , whilst proximal renal tubular acidosis (type 2 RTA)

is due to an inability of the proximal tubules to reabsorb bicarbonate (HCO_3^-)

- Metabolic acidosis leads to hyperkalaemia and vice versa

The kidneys are paired retroperitoneal structures that are normally located between the transverse processes of the T12–L3 vertebrae, with the left kidney typically somewhat more superior in position than the right. Each kidney has an outer cortex and an inner medulla which protrudes into the pelvis. The pelvis is practically the funnel-shaped dilated upper end of the ureter.

The kidney maintains a stable acid base, electrolyte, and fluid status inside the body by selective elimination or retention of water, electrolytes, and other solutes (Table 1.1). It does so by three mechanisms:

1. Filtration of blood in the glomerulus to form an ultrafiltrate (water with low molecular weight solutes) which then enters the tubule.
2. Selective reabsorption of water, electrolytes, and solutes from the tubules into the interstitium and peritubular capillaries.
3. Selective secretion from the peritubular capillaries across the tubular epithelium into the tubular fluid.

Besides these mechanisms, the kidneys also play an active endocrine role by the production and secretion of:

- *1,25-dihydroxycholecalciferol*: cholecalciferol is derived from 7-dehydrocholesterol in the skin on exposure to the ultraviolet rays in sunlight. Cholecalciferol then undergoes two subsequent hydroxylations by 25-hydroxylase in the liver and 1-hydroxylase in the proximal tubules of the kidney to yield 1,25-dihydroxycholecalciferol

(calcitriol). Calcitriol is the active form of vitamin D, without which calcium cannot be absorbed from the intestine.

- *Renin*: discussed later under juxtaglomerular apparatus.
- *Erythropoietin (EPO)*: specialized interstitial cells in the inner cortex and outer medulla of the kidney produce and secrete EPO, which stimulates red blood cell (RBC) production in the bone marrow.

The kidney consists of nephrons with the supporting interstitium, collecting ducts (CDs), and the renal microvasculature. The nephron consists of a glomerulus and a twisted tubule which drains into the CD. The tubule consists of a proximal and a distal tubule connected by Henle's loop [1]. Each kidney has approximately one million nephrons and we cannot develop new nephrons after birth.

Cortex

This is the outer layer of the kidney and all the glomeruli are located here (Figure 1.1). The tubules of the superficial and midcortical nephrons are situated entirely within the cortex. The juxtamedullary nephrons (in the deeper regions of the cortex and nearer the medulla) have longer tubules and their loop of Henle goes down into the medulla and helps in the countercurrent mechanism, as discussed later.

Medulla

The renal medulla contains the loops of Henle of the juxtamedullary nephrons, vasa recta (peritubular capillaries surrounding the long loops of the juxtamedullary nephrons), and the CDs. The medulla consists of 7–10 conical subdivisions called pyramids, whose broad base faces the cortex, and the apical papilla points into the minor calyx. After traversing through the pyramid, the CDs open at the papilla and drain the urine into the minor calyx. Two or three minor calyces converge to form a major calyx, through

Table 1.1 Urine to plasma ratios of some physiologically important body substances

Substance	Urine to plasma ratio
Glucose	0
Sodium	0.6
Urea	60
Creatinine	150




Figure 1.1 The renal cortex, medulla (with the pyramids), and minor and major calyces with their relation to the ureter and renal vasculature.

which urine continues into the renal pelvis, which is the funnel-shaped dilated proximal end of the ureter. There are usually two or three major calyces in each kidney.

Renal Vasculature

The kidneys receive 1.2–1.3 l of blood per minute (about 25% of the cardiac output), making them highly vascular organs. After originating from the aorta, the renal artery enters the renal sinus and divides into the interlobar arteries, which extend towards the cortex in the spaces between the medullary pyramids. At the junction between the cortex and medulla, the interlobar arteries divide and pass over into the arcuate arteries. The arcuate arteries give rise to the interlobular arteries, which rise radially through the cortex. It is interesting to note that none of these arteries penetrates the medulla.

Afferent arterioles arise from the interlobular arteries and supply the glomerular tufts. The glomerular tufts are drained by the efferent arterioles, which then form the peritubular plexus and are of two types: cortical and juxtamedullary. The shorter cortical efferent arterioles arise from the superficial and midcortical nephrons and supply the cortex. The longer juxtamedullary efferent arterioles, which arise from the deeper nephrons, represent the sole blood supply to the medulla and are termed vasa recta. Whilst the descending vasa recta supply blood to the medulla, the ascending vasa recta drain it.

Glomerulus and Filtration across it

The glomerulus is the invagination of a tuft of capillaries into the dilated proximal end of the nephron called the Bowman's capsule (Figure 1.2). Supplied by the afferent and drained by the efferent arteriole, the glomerular capillary bunch is attached to the mesangium on the inner side and covered by the glomerular basement membrane (GBM) on the outer side. In a way, the GBM forms the skeleton of the glomeruli, with the podocytes (visceral epithelial cells with long foot processes) on the outer side and the capillaries and mesangium on the inner side. Thus, blood within the glomerular capillary is separated from the urinary space by endothelium, GBM, and podocyte.

Formation of urine begins with the filtration of blood from the glomerular capillaries. For this to happen blood must cross three layers:

- **Endothelium:** these highly fenestrated cells have 50–100 nm pores with a highly electronegative luminal surface.
- **GBM:** with podocytes externally and endothelium/mesangium internally, its main constituents are collagen IV, laminin, and heparin, all of which are negatively charged.
- **Visceral epithelium (podocytes):** a big cell body floating in the urinary space with long primary processes which affix on the glomerular capillaries by foot processes, with filtration slits sized 30–40 nm




Figure 1.2 Parts of glomerulus and the juxtaglomerular apparatus (JGA). PCT, proximal convoluted tubule.

in between. These slits are bridged by the slit diaphragms, which are themselves penetrated by small pores sized about 8 nm. The luminal surfaces of both the podocyte foot processes and the slit diaphragms are rich in negatively charged proteins such as podocalyxin in the former and nephrin in the latter.

Whilst water can freely cross these three layers, the filtration of macromolecules depends on both size and charge. Molecules bigger than 8 nm are normally completely restricted by the GBM. Albumin with a diameter of 7.2 nm would have been effectively filtered were it not for its negative charge.

Clinical Notes

Common Glomerular Pathologies

- Nephrotic syndrome is characterized by proteinuria $>3.5\text{ g}/24\text{ hours}$, hypoalbuminaemia, generalized oedema, and hyperlipidaemia. Nephrotic syndrome is due to abnormally increased filtration of plasma proteins across the glomerular capillary. This may be due to an increase in the size of the pores mentioned earlier, but there is always an associated loss of negative charges across the filtration barrier. Proteinuria leads to hypoalbuminaemia, with a resultant decrease in plasma oncotic pressure in turn leading to oedema. In an effort to compensate for the low oncotic pressure, the liver starts to make excessive lipids, leading to hyperlipidaemia.

- Nephritis is characterized by haematuria with RBC casts and dysmorphic RBCs, some degree of oliguria, hypertension, and reduction in glomerular filtration rate (GFR). Nephritis is due to glomerular inflammation leading to leakage of RBCs across the filtration barrier. As the RBCs traverse the inflamed filtration barrier and travel down the tubules, they develop cytoplasmic blebs on their surface and are seen as dysmorphic RBCs under the microscope. Whilst travelling through the thick ascending loop of Henle, the RBCs get struck in the physiologically secreted tubular Tamm–Horsfall protein, leading to the formation of RBC casts. Thus, the presence of dysmorphic RBCs and RBC casts in a patient with haematuria helps to differentiate nephritis as the pathology rather than ureteric or bladder pathologies.
- Type IV collagen molecules are composed of three alpha chains that form triple-helical structures through specific interactions of C-terminal non-collagenous domains. GBM and alveolar capillary basement membrane collagen consist of alpha3, -4, and -5 chains, unlike other basement membranes with alpha1 (two of them) and alpha2 chains. Two disease processes pathogenetically caused by defects primarily in the GBM are:
 - a. Goodpasture's disease, characterized by anti-GBM antibody against the alpha3 chain of the type IV collagen present in the GBM and alveolar membranes in the lungs [2]. Patients present with nephritis and haemoptysis.
 - b. Alport syndrome, which results from mutations in genes encoding the alpha3, -4, and -5 chains of type IV collagen (80% are due to mutations in the alpha5 chain) [3]. It is the commonest inherited form of nephritis and is often associated with sensorineural deafness and ocular abnormalities due to the presence of similar alpha3, -4, and -5 chains in the type IV collagen in the basement membranes of the lens and cochlea.

Tubules

With a GFR of 125 ml/min, about 180 l of urine is formed at the glomerular level each day. Thankfully, under physiological conditions almost 98–99% of this filtered urine is reabsorbed by the tubules. More than 90% of this tubular water reabsorption is a passive process and water just follows the sodium (Na^+) being reabsorbed. Only in the CD is water absorbed actively and independently of Na^+ under the action of vasopressin or anti-diuretic hormone (ADH).

It is thus obvious that most of the water reabsorption in the tubules is dependent on the reabsorption of Na^+ . So, what makes the tubules take up Na^+ so avidly?

Why do the Tubules Reabsorb Na^+ and Water?

The tubular epithelial cells, irrespective of which segment of the tubule they belong to, have a basolateral Na^+/K^+ -ATPase that continuously pumps out three Na^+ ions and brings in two potassium (K^+) ions, thus making the cells permanently Na^+ deficient. Thus, the

fact that the tubular epithelial cells are always deficient in Na^+ helps in the Na^+ reabsorption from the tubules. In the same vein, the fact that the tubular cells are rich in K^+ helps them to secrete K^+ into the tubular lumen.

Water is absorbed osmotically along with solutes, with Na^+ being the predominant solute (Table 1.2; Figure 1.3). The two exceptions are in the thick ascending limb (TAL) of the loop of Henle, which is impermeable to water, and the CD, where water is reabsorbed by the action of vasopressin, as discussed later.

Proximal Convoluted Tubule

The bulk of tubular reabsorption occurs in the proximal convoluted tubule (PCT), with 65–70% of the total Na^+ being reabsorbed here and about the same amount of water being reabsorbed passively. The PCT is also responsible for reabsorption of the bulk of phosphate (PO_4^{3-}), bicarbonate (HCO_3^-), potassium (K^+), calcium (Ca^{2+}) and almost all glucose, amino acids, uric acid, and low molecular weight proteins such as microglobulins and the light chain fraction of immunoglobulins.

NaHCO_3 is filtered in the glomeruli and dissociates in the lumen of the PCT into Na^+ and HCO_3^- . Na^+ is

Table 1.2 Absorption of various ions and water across different segments of the renal tubules

Ion or substance	PCT	Loop of Henle	DCT	CD
Na ⁺	65–70%	30–35%	5%	1–2%
H ₂ O	65–70%	25%	5%	5–10%
K ⁺	45–50%	40–45%		Secreted by principal cells
Ca ⁺	70%	20%	10–15%	
Mg ⁺	15–25%	60–70%	5–10%	
PO ₄ ³⁻	70–80%	Insignificant	Insignificant	Insignificant
HCO ₃ ⁻	85–90%			Secreted by beta- intercalated cells

Ca⁺⁺: calcium; CD: collecting duct; DCT: distal convoluted tubule; HCO₃⁻: bicarbonate; H₂O: water; K⁺: potassium; Mg⁺: magnesium; Na⁺: sodium; PCT: proximal convoluted tubule; PO₄³⁻: phosphate.





Figure 1.3 Diagram showing the movement of sodium (Na^+), chloride (Cl^-), potassium (K^+), and water (H_2O) across various portions of the renal tubule, with the site of action of different diuretics highlighted. ADH: anti-diuretic hormone; NaHCO_3^- : sodium bicarbonate; TAL: thick ascending limb.

reabsorbed by the Na^+ -hungry tubular epithelial, as already discussed. The HCO_3^- remaining (from the original NaHCO_3) in the lumen combines with hydrogen (H^+) by the action of the tubular carbonic anhy-

dase (CA) to form carbonic acid (H_2CO_3), which then splits into carbon dioxide (CO_2) and water (H_2O). CO_2 enters the tubular cell and now the reaction goes in the reverse direction, with CO_2 combining with H_2O



Reabsorption of Na^+ in the PCT facilitates the reabsorption of glucose, amino acids, phosphate, and uric acid.

Clinical Notes

Disorders of the Proximal Renal Tubules

- Proximal renal tubular acidosis (type 2 RTA) is due to a defect in the reabsorption of HCO_3^- leading to urinary loss of HCO_3^- and the resultant metabolic acidosis. As already explained, the reabsorption of HCO_3^- in this segment is linked to the coupled exchange of H^+ and Na^+ (Figure 1.4). As Na^+ reabsorption here is linked to the reabsorption of glucose, amino acids, phosphate, and uric acid, the patient may present with glycosuria, hypophosphataemia, and hypouricaemia (due to urinary losses), a condition that is termed Fanconi syndrome. Urine pH is usually less than 5.5 despite bicarbonaturia, as the intact distal tubules excrete excess H^+ as a compensatory mechanism for the acidosis.
- Type 2 RTA is discussed further in the clinical notes following the section on the distal convoluted tubule and collecting duct.
- Acetazolamide, which is a CA inhibitor often used by ophthalmologists for the treatment of glaucoma, can also lead to a type 2 RTA-like picture, with full-blown Fanconi syndrome at times. The anti-epileptic drug topiramate and anti-HIV medication tenofovir, by their CA inhibitor activity, can also lead to proximal RTA and Fanconi syndrome [4].
- PCT is responsible for the reabsorption of the easily filtered light chain fragments of immunoglobulins. In myeloma, the excess of light chains is toxic to the PCT and may lead to proximal RTA with or without Fanconi syndrome [5].
- In prerenal AKI, serum urea increases disproportionately to creatinine due to its enhanced PCT reabsorption that follows the enhanced transport of Na^+ and water. Elevated serum urea/creatinine may also be seen in upper gastrointestinal haemorrhage, since the upper intestinal proteolytic enzymes break down haemoglobin to amino acids, which are then reabsorbed by the body and lead to the generation of urea.

Clinical Notes

Disorders of the Loop of Henle

- *Bartter syndrome*: an autosomal recessive syndrome that characteristically presents with hypokalaemia and metabolic alkalosis is described later in the chapter with Gitelman syndrome (following the section on the distal convoluted tubule and collecting duct).
- *Familial hypocalciuric hypercalcaemia (FHH)*: this benign autosomal dominant condition is caused by inactivating mutations in the gene for the CaSR.

CaSR is predominantly found in the parathyroid gland and the kidneys. In the parathyroid gland, it decreases the secretion of PTH in response to serum Ca^{2+} . Under normal conditions, the TAL CaSR, on being activated by Ca^{2+} , downregulates the production of Ca^{2+} -carrying transcellular proteins Claudrin 16 and 19 by forming Claudrin 14, which has an inhibitory effect on the two [7]. The mutated CaSR in FHH is unable to inhibit Claudrin 16 and 19, leading to uninhibited Ca^{2+} reabsorption and hence hypocalciuria and hypercalcaemia, and at the same time a non-suppressible serum PTH level. It is easy to confuse this condition with primary hyperparathyroidism unless one looks at the urinary Ca^{2+} , which is characteristically low. A spot urine Ca^{2+} /creatinine ratio of $<1/100$ is helpful in differentiating the hypercalcaemia with inappropriately high serum PTH levels seen in FHH from primary hyperparathyroidism (where the urinary Ca^{2+} would be high).

About 40–50% of the filtered urea is reabsorbed in the PCT. As urea is reabsorbed in this section along with Na^+ , in conditions of volume contraction when the PCT reabsorbs more Na^+ , the net reabsorption of urea is also significantly increased. Thus, a disproportionate rise in blood urea concentration in comparison to creatinine might be a marker of a prerenal cause of acute kidney injury (AKI).

Loop of Henle

The loop of Henle consists of the thick and thin descending limbs and the thick ascending limb (TAL), with a thin ascending limb seen only in long looped nephrons.




Figure 1.5 Transport mechanisms in the thick ascending limb of the loop of Henle: The basolateral (peritubular) membrane due to the action of $\text{Na}^+-\text{K}^+-\text{ATPase}$ pumps three sodium (Na^+) out of, and two potassium (K^+) into, the cell. This creates a low intracellular Na^+ concentration, which allows the cell to absorb Na^+ from the urinary lumen by the electroneutral $\text{Na}^+-\text{K}^+-2\text{Cl}^-$ (NKCC2). The cell is rich in K^+ and so much of the reabsorbed K^+ recycles back into the lumen through the renal outer medullary potassium channel (ROMK) on the apical/luminal membrane. This movement of one cationic K^+ into the lumen plus the flux of one reabsorbed Na^+ (by $\text{Na}^+-\text{K}^+-\text{ATPase}$ pump) and two chloride (Cl^-) out of the cell and into the peritubular capillary (via Cl^- channels) causes the lumen to become more positively charged compared with the cell and peritubular capillaries/interstitium. The resultant electrical gradient promotes passive reabsorption of cations calcium (Ca^{2+}) and magnesium (Mg^{2+}) via the paracellular pathway. The Cl^- channels require interaction with a small protein called barttin to function normally. ADP, adenosine diphosphate; ATP, adenosine triphosphate; P_i , inorganic phosphate.

The loop of Henle is responsible for the reabsorption of 30–35% of the luminal Na^+ but only 25% of the water, as the TAL is resistant to the movement of water across it.

The frusemide-sensitive and electroneutral $\text{Na}^+-\text{K}^+-2\text{Cl}^-$ cotransporter reabsorbs one Na^+ and K^+ each along with two Cl^- ions from the TAL lumen to the tubular epithelial cell (Figure 1.5). High intracellular K^+ concentration (due to the basolateral $\text{Na}^+-\text{K}^+-\text{ATPase}$ pumping out three Na^+ and bringing in two K^+ ions constantly) causes the reabsorbed K^+ to be secreted back into the lumen via the renal outer medullary potassium (ROMK) channel. The Cl^- is absorbed into blood by the basolateral Cl^- channel (CIC-Kb), which has barttin as a crucial accessory protein. Movement of one positive charge (K^+) into the lumen and net of one negative charge into the peritubular capillary (1 Na^+ and 2 Cl^-) causes the lumen to be more positive compared to the tubular peritubular capillary and interstitium. This electrical gradient promotes the passive reabsorption of cations Ca^{2+} and Mg^{2+} via the paracellular cleft between

the cells. Membrane proteins Claudrin 16 and 19 help in this paracellular reabsorption of Ca^{2+} and Mg^{2+} and are themselves tightly regulated by the calcium-sensing receptor (CaSR) (Figure 1.6) [6].

Countercurrent Mechanism

A major function of the loop of Henle is to create an interstitial osmotic gradient that increases from the renal cortex (290 mOsm/kg) to the tip of the medullary pyramid (1200 mOsm/kg) (Figure 1.7). This countercurrent mechanism is fundamentally dependent on the impermeability of the TAL to water. The reabsorption of Na^+ (and its passage to the interstitium by the basolateral $\text{Na}^+-\text{K}^+-\text{ATPase}$ pump) in the absence of water in the ascending limb has two consequences:




Figure 1.6 Absorption of calcium (Ca^{2+}) and magnesium (Mg^{2+}) through the paracellular pathway in the thick ascending limb of the loop of Henle. Sodium (Na^+) is reabsorbed via the apical $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ cotransporter and pumped out basolaterally by the $\text{Na}^+/\text{K}^+/\text{ATPase}$; chloride (Cl^-) exits via the Cl^- channel (ClC-Kb); and potassium (K^+) is secreted into the tubular lumen by the renal outer medullary potassium channel (ROMK), thereby generating a lumen-positive transtubular electric potential. Ca^{2+} and Mg^{2+} are reabsorbed passively across the paracellular pathway because of this lumen-positive electrical potential. Claudrin 16 and 19 are the two chief transport proteins which help to carry the Ca^{2+} and Mg^{2+} across the paracellular cleft. Blood Ca^{2+} levels regulate this process by stimulating the Ca^{2+} -sensing receptor (CaSR), which then upregulates Claudin-14 (CLDN14), a protein that normally binds to and inhibits Claudin 16 and -19. CaSR controls the production of Claudin 14 by the calcineurin-NFATc1-microRNA pathway. NFAT: nuclear factor of activated T cells.




Figure 1.7 Countercurrent mechanism: reabsorption of sodium (Na^+) with relative resistance to the reabsorption of water by the thick ascending limb of the loop of Henle leads to generation of the interstitial osmotic gradient; the gradient is higher for the longer loops of the juxtamedullary nephrons, which travel deeper into the medulla. H_2O , water; $NaCl$, sodium chloride.

1. A higher concentration of Na^+ in the interstitium leads to higher interstitial osmotic pressure, with the highest osmotic pressure developed in tips of medullary pyramids where the longer loops of Henle reach.
2. The tubular fluid leaving the TAL is hypotonic because of excess water in comparison to Na^+ .

The creation of this gradient helps in the reabsorption of water in the medullary CD under the influence of vasopressin, with water moving from the hypo-osmolar tubular environment to the hyper-osmolar medullary interstitium.

Clinical Notes

Can Frusemide Lead to Hyponatraemia?

Frusemide acts by interrupting the Na^+ uptake in the TAL by the $Na^+-K^+-2Cl^-$ transporter. This interferes with the countercurrent mechanism and the generation of the medullary interstitial osmotic gradient. As discussed, without this gradient, ADH cannot reabsorb water in the medullary CD. Thus, frusemide leads to the excretion of a dilute urine which would prevent the development of hyponatraemia.

Distal Convoluted Tubule and Collecting Duct

The distal convoluted tubule (DCT) is responsible for the reabsorption of 5% of Na^+ by the thiazide-sensitive sodium chloride ($NaCl$) cotransporter. Between 10% and 15% of the Ca^{2+} is also reabsorbed in this segment in an active process, unlike the more passive process in the proximal parts (70% of Ca^{2+} is co-transported passively with Na^+ in the PCT). Ca^{2+} enters tubular cells via the transient receptor potential vanilloid 5 (TRPV5) channel, and 5–10% of the total Mg^{2+} reabsorption also occurs in this segment by the transient receptor potential melastatin 6 (TRPM6) and TRPM7 channels.

The CD is responsible for reabsorption of up to 2% of the Na^+ and water, but this can vary depending on the needs of the body. The two chief types of cells found in the CD are:

- Principal cells, which are responsible for Na^+ reabsorption and K^+ secretion (under the influence of aldosterone) and water reabsorption (under the influence of ADH).

- Intercalated cells, which are responsible for the secretion of H^+ (by alpha-intercalated cell) or HCO_3^- (by beta-intercalated cell).

ADH

ADH is synthesized in the supraoptic and paraventricular nuclei of the hypothalamus and travels down the pituitary stalk to be stored in the posterior pituitary. Secreted in response to decreased intravascular volume or increased plasma osmotic pressure as in dehydration, the main physiological effects of ADH are retention of water by the kidneys and vasoconstriction.

There are three types of ADH receptors: V1A, V1B, and V2. V1A receptors cause vasoconstriction by increasing vascular smooth muscle intracellular Ca^{2+} concentration, whilst V1B receptors lead to adrenocorticotrophic hormone (ACTH) release by the pituitary gland. V2 receptors are found on the principal cells in the CD and via the cyclic adenosine monophosphate (cAMP) pathway lead to movement and fusion of the aquaporin 2 (AQP2) water channels to the luminal membrane (Figure 1.8). The medullary osmotic gradient established due to the selective reabsorption of Na^+ in the TAL helps in the water being reabsorbed in the CD.




Figure 1.8 Anti-diuretic hormone (vasopressin), acting through cyclic adenosine monophosphate (cAMP), increases the total number of aquaporin-2 molecules in the principal cells of the collecting duct; aquaporin-2 aids in water reabsorption in this part of the renal tubule. H_2O : water.

Aldosterone

Aldosterone is secreted by the zona glomerulosa of adrenal cortex and helps in Na^+ reabsorption and K^+ and H^+ excretion by the distal portions of the tubules. On binding to the mineralocorticoid receptor, aldosterone upregulates and activates the basolateral Na^+/K^+ -ATPase in the principal cells, causing lower intracellular Na^+ , and at the same time upregulates epithelial sodium channels (ENaC), increasing apical membrane permeability for Na^+ . The resultant

intracellular movement of Na^+ causes luminal negativity, leading in turn to secretion of the positively charged K^+ . The Na^+ and K^+ movement is not a strict one-to-one exchange and more Na^+ is reabsorbed than K^+ is lost. So, the same luminal negativity leads to H^+ secretion by the neighbouring alpha-intercalated cells. The potassium-sparing diuretics (amiloride and triamterene) act by directly inhibiting ENaC channels; spironolactone acts by competing with aldosterone for binding to the mineralocorticoid receptor (Figure 1.9).




Figure 1.9 Ion transport in collecting tubule principal cells. Aldosterone, after combining with the cytosolic mineralocorticoid receptor (aldo-receptor), leads to enhanced sodium (Na^+) reabsorption and potassium (K^+) secretion by increasing both the number of open Na^+ channels and the number of Na^+/K^+ -ATPase pumps. Whilst spironolactone competes with aldosterone for the aldosterone receptor, amiloride and triamterene act by directly inhibiting the epithelial Na^+ channel. ENaC: epithelial sodium channel.

Clinical Notes

Diuretics, the Distal Tubules, Vasopressin and Aldosterone

- The unifying mechanistic principle of action of all diuretics is to prevent tubular Na^+ reabsorption. Both thiazide and loop diuretics can lead to hypokalaemia and metabolic alkalosis by the following mechanisms:
 - With more Na^+ flowing to the distal parts of the nephron, the tubular cells try to reclaim the excess Na^+ , resulting in loss of positive charges from the lumen. The tubular cells try to compensate by secretion of the positively charged K^+ (by principal cells) and H^+ (by alpha-intercalated cell) into the lumen.
 - Intravascular volume contraction because of diuretics causes activation of the renin-angiotensin-aldosterone (RAS), which contributes to the urinary K^+ and H^+ losses due to the effect of aldosterone.
- Potassium-sparing diuretics can lead to hyperkalaemia and metabolic acidosis. Spironolactone competes with aldosterone for binding to the mineralocorticoid receptor, whilst amiloride and triamterene act by directly inhibiting the ENaC. The resultant effect is inhibition of K^+ and H^+ secretion.
- Effects of diuretics on Ca^{2+} : Let us recapitulate that the paracellular reabsorption of Ca^{2+} in the loop of Henle is dependent on the positive electrical gradient in the lumen (Figures 1.6). Loop diuretics by inhibiting the $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ transporter inhibit generation of this gradient and hence prevent reabsorption of Ca^{2+} in this segment of the tubule. On the other hand, thiazide diuretics increase proximal Na^+ and water reabsorption due to volume depletion, leading in turn to increased passive proximal calcium reabsorption. Thus, whilst furosemide helps to excrete Ca^{2+} in the urine, thiazide diuretics decrease urinary Ca^{2+} excretion and lead to hypercalcaemia.

- Lithium causes nephrogenic diabetes insipidus by causing decreased expression of AQP 2 genes.
- Primary hyperaldosteronism should always be suspected in a patient with hypokalaemia, metabolic alkalosis, and hypertension (more than half of patients are normokalaemic). This is discussed in more detail in Chapter 3.
- Bartter and Gitelman syndromes, both autosomal recessive disorders, are characterized by hypokalaemia, metabolic alkalosis, and low to normal blood pressure, with hypomagnesaemia being an additional distinctive feature of Gitelman syndrome, which tends to present in adults and is generally the milder of the two. The tubular defects in sodium chloride reabsorption in Bartter and Gitelman syndromes are almost identical to those seen with chronic ingestion of loop and thiazide diuretics, respectively [8, 9, 10]. Additional down-regulation of the Mg^{2+} -reabsorbing TRPM6 channel in Gitelman syndrome causes Mg^{2+} wasting and the resultant characteristic hypomagnesaemia. These disorders are discussed in more detail in Chapter 3.
- Distal renal tubular acidosis (type 1 RTA) is due to an inability of the distal tubules to excrete H^{+} , leading to metabolic acidosis. The kidney characteristically cannot lower the urinary pH to less than 5.5 due to a deficiency of H in the urine, in contrast to type 2 RTA where the urinary pH is usually greater than 5.5 (Table 1.3) [11].

Table 1.3 Differences between types 1 and 2 renal tubular acidosis (RTA)

RTA1 (distal RTA)	RTA2 (proximal RTA)
Inability to secrete H^{+}	Inability to reabsorb HCO_3^{-}
Urine pH >5.5 (absence of H^{+} in urine)	Urine pH <5.5 (compensatory rise in DCT secretion of excess H^{+})
Associations: Sjögren's syndrome, systemic lupus erythematosus, primary biliary cirrhosis, autoimmune hepatitis	Associations: myeloma, Wilson's disease, tenofovir, acetazolamide, gentamicin
No Fanconi syndrome	Fanconi syndrome may be seen: glycosuria, phosphaturia, uric aciduria, and aminoaciduria (low serum phosphate and urate concentrations)
Proximal tubules reabsorb all alkali including citrate, which normally keeps Ca^{2+} in urine soluble, leading to increased predisposition to nephrolithiasis and nephrocalcinosis	No renal stones
Treat with alkali and K^{+} replacement	Same treatment, but needs bigger doses of alkali as glomeruli lose the administered alkali and tubules cannot reabsorb it due to the nature of the defect

DCT: distal convoluted tubule; H^{+} : hydrogen; HCO_3^{-} : bicarbonate; K^{+} : potassium.

Glomeruli and Tubule Balance each other

An increase or decrease in glomerular filtration is balanced by increased or decreased tubular reabsorption. Similarly, an increase in tubular flow leads to a decrease in glomerular filtration and vice versa.

Glomerulotubular Balance

Increased filtration of protein-free fluid across the glomeruli leads to the blood flowing to efferent and peritubular arterioles being more concentrated. The resultant increased oncotic pressure in the peritubular capillaries aids in enhanced reabsorption of water across the tubular epithelium.