

MATHEMATICS AND STATISTICS



Statistical Topics and Stochastic Models for Dependent Data with Applications

**Edited by
Vlad Stefan Barbu and Nicolas Vergne**

ISTE

WILEY

Statistical Topics and Stochastic Models
for Dependent Data with Applications

Series Editor
Nikolaos Limnios

**Statistical Topics and Stochastic
Models for Dependent Data
with Applications**

Edited by

Vlad Stefan Barbu
Nicolas Vergne

ISTE

WILEY

First published 2020 in Great Britain and the United States by ISTE Ltd and John Wiley & Sons, Inc.

Apart from any fair dealing for the purposes of research or private study, or criticism or review, as permitted under the Copyright, Designs and Patents Act 1988, this publication may only be reproduced, stored or transmitted, in any form or by any means, with the prior permission in writing of the publishers, or in the case of reprographic reproduction in accordance with the terms and licenses issued by the CLA. Enquiries concerning reproduction outside these terms should be sent to the publishers at the undermentioned address:

ISTE Ltd
27-37 St George's Road
London SW19 4EU
UK

www.iste.co.uk

John Wiley & Sons, Inc.
111 River Street
Hoboken, NJ 07030
USA

www.wiley.com

© ISTE Ltd 2020

The rights of Vlad Stefan Barbu and Nicolas Vergne to be identified as the authors of this work have been asserted by them in accordance with the Copyright, Designs and Patents Act 1988.

Library of Congress Control Number: 2020938718

British Library Cataloguing-in-Publication Data
A CIP record for this book is available from the British Library
ISBN 978-1-78630-603-6

Contents

Preface	xi
Vlad Stefan BARBU and Nicolas VERGNE	
Part 1. Markov and Semi-Markov Processes	1
Chapter 1. Variable Length Markov Chains, Persistent Random Walks: A Close Encounter	3
Peggy CÉNAC, Brigitte CHAUVIN, Frédéric PACCAUT and Nicolas POUYANNE	
1.1. Introduction	3
1.2. VLMCs: definition of the model	6
1.3. Definition and behavior of PRWs	9
1.3.1. PRWs in dimension one	9
1.3.2. PRWs in dimension two	13
1.4. VLMC: existence of stationary probability measures	15
1.5. Where VLMC and PRW meet	19
1.5.1. Semi-Markov chains and Markov additive processes	19
1.5.2. PRWs induce semi-Markov chains	20
1.5.3. Semi-Markov chain of the α -LIS in a stable VLMC	22
1.5.4. The meeting point	23
1.6. References	27
Chapter 2. Bootstraps of Martingale-difference Arrays Under the Uniformly Integrable Entropy	29
Salim BOUZEBDA and Nikolaos LIMNIOΣ	
2.1. Introduction and motivation	29
2.2. Some preliminaries and notation	30
2.3. Main results	35
2.4. Application for the semi-Markov kernel estimators	36

2.5. Proofs	41
2.6. References	45

Chapter 3. A Review of the Dividend Discount Model: From Deterministic to Stochastic Models

Guglielmo D'AMICO and Riccardo DE BLASIS

3.1. Introduction	47
3.2. General model	48
3.3. Gordon growth model and extensions	50
3.3.1. Gordon model	50
3.3.2. Two-stage model	51
3.3.3. H model	52
3.3.4. Three-stage model	52
3.3.5. N-stage model	53
3.3.6. Other extensions	53
3.4. Markov chain stock models	54
3.4.1. Hurley and Johnson model	54
3.4.2. Yao model	56
3.4.3. Markov stock model	57
3.4.4. Multivariate Markov chain stock model	61
3.5. Conclusion	64
3.6. References	65

Chapter 4. Estimation of Piecewise-deterministic Trajectories in a Quantum Optics Scenario

Romain AZAÏS and Bruno LEGGIO

4.1. Introduction	69
4.1.1. The postulates of quantum mechanics	69
4.1.2. Dynamics of open quantum Markovian systems	71
4.1.3. Stochastic wave function: quantum dynamics as PDPs	74
4.1.4. Estimation for PDPs	76
4.2. Problem formulation	77
4.2.1. Atom-field interaction	77
4.2.2. Piecewise-deterministic trajectories	78
4.2.3. Measures	80
4.3. Estimation procedure	80
4.3.1. Strategy	80
4.3.2. Least-square estimators	82
4.3.3. Numerical experiments	83
4.4. Physical interpretation	86
4.5. Concluding remarks	87
4.6. References	88

Chapter 5. Identification of Patterns in a Semi-Markov Chain 91

Brenda Ivette GARCIA-MAYA and Nikolaos LIMNIOS

5.1. Introduction	91
5.2. The prefix chain	93
5.3. The semi-Markov setting	94
5.4. The hitting time of the pattern	100
5.5. A genomic application	102
5.6. Concluding remarks	106
5.7. References	106

Part 2. Autoregressive Processes 109

Chapter 6. Time Changes and Stationarity Issues for Continuous Time Autoregressive Processes of Order p 111

Valérie GIRARDIN and Rachid SENOUSI

6.1. Introduction	111
6.2. Basics	112
6.3. Stationary AR processes	114
6.3.1. Formulas for the two first-order moments	114
6.3.2. Examples	116
6.3.3. Conditions for stationarity of $CAR_1(p)$ processes	118
6.4. Time transforms	125
6.4.1. Properties of time transforms	125
6.4.2. MS processes	131
6.5. Conclusion	132
6.6. Appendix	133
6.7. References	136

Chapter 7. Sequential Estimation for Non-parametric Autoregressive Models 139

Ouerdia ARKOUN, Jean-Yves BRUA and Serguei PERGAMENCHTCHIKOV

7.1. Introduction	139
7.2. Main conditions	141
7.3. Pointwise estimation with absolute error risk	142
7.3.1. Minimax approach	142
7.3.2. Adaptive minimax approach	144
7.3.3. Non-adaptive procedure	145
7.3.4. Sequential kernel estimator	148
7.3.5. Adaptive sequential procedure	151
7.4. Estimation with quadratic integral risk	153
7.4.1. Passage to a discrete time regression model	155
7.4.2. Model selection	159

7.4.3. Main results	161
7.5. References	164

Part 3. Divergence Measures and Entropies 167

**Chapter 8. Inference in Parametric and Semi-parametric Models:
The Divergence-based Approach 169**

Michel BRONIATOWSKI

8.1. Introduction	169
8.1.1. Csiszár divergences, variational form	170
8.1.2. Dual form of the divergence and dual estimators in parametric models	172
8.1.3. Decomposable discrepancies	178
8.2. Models and selection of statistical criteria	183
8.3. Non-regular cases: the interplay between the model and the criterion	184
8.3.1. Test statistics	185
8.4. References	187

**Chapter 9. Dynamics of the Group Entropy Maximization
Processes and of the Relative Entropy Group Minimization
Processes Based on the Speed-gradient Principle 189**

Vasile PREDA and Irina BĂNCESCU

9.1. Introduction	190
9.1.1. The SG principle	191
9.1.2. Entropy groups	193
9.2. Group entropies and the SG principle	196
9.2.1. Total energy constraint	199
9.3. Relative entropy group and the SG principle	202
9.3.1. Equilibrium stability	205
9.3.2. Total energy constraint	205
9.4. A new (G, a) power relative entropy group and the SG principle	206
9.5. Conclusion	210
9.6. References	210

**Chapter 10. Inferential Statistics Based on Measures of
Information and Divergence 215**

Alex KARAGRIGORIOU and Christos MESELIDIS

10.1. Introduction	215
10.2. Divergence measures	216
10.2.1. φ -Divergences	216
10.2.2. α -Divergences	217
10.2.3. Bregman divergences	218

10.3. Properties of divergence measures	219
10.4. Model selection criteria	220
10.5. Goodness of fit tests	222
10.5.1. Simple null hypothesis	222
10.5.2. Composite null hypothesis	223
10.6. Simulation study	227
10.7. References	231
Chapter 11. Goodness-of-Fit Tests Based on Divergence Measures for Frailty Models	235
Filia VONTA	
11.1. Introduction	235
11.2. The proposed goodness-of-fit test	236
11.3. Main results	240
11.4. Frailty models	243
11.5. Simulations	244
11.5.1. Linear models for the estimation of critical values	247
11.5.2. Size of the test	248
11.6. References	250
List of Authors	253
Index	257

Preface

This collective book stems from a workshop that took place in Rouen, October 3–5, 2018, within the framework of the project *Random Models and Statistical Tools, Informatics and Combinatorics (MOUSTIC – Modèles aléatoires et Outils Statistiques, Informatiques et Combinatoires)*, financed by the Region of Normandy and the European Regional Development Fund. The main idea was to bring together leading scientists working on probabilistic and statistical topics for dependent data, as well as on associated applications.

This is the way this book was written, with the intention to offer to the scientific community a part of the latest advances in the field of stochastic modeling for dependent data, authored by leading experts in the field. This is a crucial aspect in a time when we face an increasing need for more and more complex models capable of capturing the main features of increasingly complex applications. From a technical point of view, our book is important for gathering theoretical developments and applications related to Markov type models (semi-Markov processes, autoregressive processes, piecewise deterministic Markov processes, and variable length Markov chains) as well as probabilistic/statistical techniques issued from information theory, based on divergence measures and entropies.

This volume is divided into three parts: the first one examines Markov and semi-Markov processes, the second one deals with autoregressive processes and the last one presents divergence measures and entropies. Particular attention is given to applications of these methods in various fields such as finance, DNA analysis, quantum physics and survival analysis.

The workshop in Rouen and, consequently, the present book, would not have been possible without the support of the Laboratory of Mathematics Raphaël Salem, the Laboratory of Mathematics Nicolas Oresme, the University of Rouen Normandy, the Region of Normandy, the French Statistical Society-SFds, the MAS group of the

French Society of Applied and Industrial Mathematics-SMAI, the Normandie-Mathématiques Research Federation and the Normastic Research Federation.

We would like to thank all the speakers of the workshop who contributed, although some of them indirectly, to the quality of the present volume. Our thanks go also to the anonymous reviewers for their valuable work: without their support, this volume could not have been successfully completed.

We would also like to thank Professor Nikolaos Limnios for proposing and encouraging us to elaborate this collective volume as well as the editorial staff of ISTE Ltd for their technical support.

Vlad Stefan BARBU
Nicolas VERGNE
Rouen, May 2020

PART 1

Markov and Semi-Markov Processes

Variable Length Markov Chains, Persistent Random Walks: A Close Encounter

We consider a walker on the line that at each step keeps the same direction with a probability that depends on the time already spent in the direction the walker is currently moving. These walks with memories of variable length can be seen as generalizations of directionally reinforced random walks (DRRWs) introduced in Mauldin *et al.* (1996). We give a complete and usable characterization of the recurrence or transience in terms of the probabilities to switch the direction. These conditions are related to some characterizations of existence and uniqueness of a stationary probability measure for a particular Markov chain: in this chapter, we define the general model for words produced by a variable length Markov chain (VLMC) and we introduce a key combinatorial structure on words. For a subclass of these VLMC, this provides necessary and sufficient conditions for existence of a stationary probability measure.

1.1. Introduction

This is the story of the encounter between two worlds: the world of random walks and the world of VLMCs. The meeting point turns around the semi-Markov property of underlying processes.

In a VLMC, unlike fixed-order Markov chains, the probability to predict the next symbol depends on a possibly unbounded part of the past, the length of which depends on the past itself. These relevant parts of pasts are called *contexts*. They are stored in a *context tree*. With each context, a probability distribution is associated, prescribing the conditional probability of the next symbol, given this context.

VLMCs are now widely used as random models for character strings. They were introduced in Rissanen (1983) to perform data compression. When they have a finite memory, they provide a parsimonious alternative to fixed-order Markov chain models, in which the number of parameters to estimate grows exponentially fast with the order; they are also able to capture finer properties of character sequences. When they have infinite memory – this will be our case of study in this chapter – they are a tractable way to build non-Markov models and they may be considered as a subclass of “*chaînes à liaisons complètes*” (Doebelin and Fortet 1937) or “chains with infinite order” (Harris 1955).

VLMCs are used in bioinformatics, linguistics and coding theory to model how words grow or to classify words. In bioinformatics, both for protein families and DNA sequences, identifying patterns that have a biological meaning is a crucial issue. Using VLMC as a model enables one to quantify the influence of a meaning pattern by giving a transition probability on the following letter of the sequence. In this way, these patterns appear as contexts of a context tree. Note that their length may be unbounded (Bejerano and Yona 2001).

In addition, if the context tree is recognized to be a signature of a family (say, of proteins), this gives an efficient statistical method to test whether or not two samples belong to the same family (Busch *et al.* 2009).

Therefore, estimating a context tree is an issue of interest and many authors (statisticians or not, applied or not) stress the fact that the height of the context tree should not be supposed to be bounded. This is the case in Galves and Leonardi (2008) where the algorithm `CONTEXT` is used to estimate an unbounded context tree, or in Garivier and Leonardi (2011). Furthermore, as explained in Csiszár and Talata (2006), the height of the estimated context tree grows with the sample size, so that estimating a context tree by assuming *a priori* that its height is bounded is not realistic.

There is extensive literature on the construction of efficient estimators of context trees, as well for finite or infinite context trees. This chapter is not a review of statistics issues, which would already be relevant for finite memory VLMC. This is a study of the probabilistic properties of infinite memory VLMC as random processes, and more specifically of the main property of interest for such processes: existence and uniqueness of a stationary measure.

As has already been said, VLMC are a natural generalization to infinite memory of Markov chains. It is usual to index a sequence of random variables forming a Markov chain with positive integers and to make the process grow to the right. The main drawback of this habit for an infinite memory process is that the sequence of the process is read from left to right, whereas the (possibly infinite) sequence giving the past needed to predict the next symbol is read in the context tree from right to left,

thus giving rise to confusion and lack of readability. For this reason, in this chapter, the VLMC grows to the left. In this way, both the process sequence and the memory in the context tree are read from left to right.

Classical random walks have *independent* and identically distributed increments. In the literature, *persistent* random walks (PRMs), also called *Goldstein-Kac random walks* or *correlated random walks*, refer to random walks having a Markov chain of finite order as an increment process. For such walks, the dynamics of trajectories has a short memory of given length and the random walk itself is not Markovian anymore. What happens whenever the increments depend on a *non-bounded* past memory?

Consider a walker on $\mathbb{Z}$, allowed to increment its trajectory by -1 or 1 at each step of time. Assume that the probability to keep the current direction ± 1 depends on the time already spent in the said direction – the distribution of increments thus acts as a reinforcement of the dependency from the past. More precisely, the process of increments of such a one-dimensional random walk is a Markov chain on the set of (right-)infinite words, with variable – and unbounded – length memory: a VLMC. The concerned VLMC is defined in section 1.3.1. It is based on a context tree called a *double comb*. Later, section 1.3.2 deals with a two-dimensional persistent random walk defined in an analogous manner on $\mathbb{Z}^2$ by a VLMC based on a context tree called a *quadruple comb*.

These random walks that have an unbounded past memory can be seen as a generalization of “directionally reinforced random walks (DRRW)” introduced by Mauldin *et al.* (1996), in the sense that the persistence times are anisotropic ones. For a one-dimensional random walk associated with a double comb, a complete characterization of recurrence and transience, in terms of changing (or not) direction probabilities, is given in section 1.3.1. More precisely, when one of the random times spent in a given direction (the so-called *persistence times*) is an integrable random variable, the recurrence property is equivalent to a classical drift-vanishing. In all other cases, the walk is transient unless the weight of the tail distributions of both persistent times are equal. In two-dimensional random walk, sufficient conditions of transience of recurrence are given in section 1.3.2.

Actually, because of the very specific form of the underlying driving VLMC, these PRWs turn out to be in one-to-one correspondence with so-called *Markov additive processes*. Section 1.5 examines the close links between PRWs, Markov additive processes, semi-Markov chains and VLMC.

In section 1.2, the definition of a general VLMC and a couple of examples are given. In section 1.3, the PRWs are defined and known results on their recurrence properties are collected. In view of section 1.5 where we show how PRW and VLMC meet through the world of semi-Markov chains, section 1.4 is devoted to results – together with a heuristic approach – on the existence and unicity of stationary measures for a VLMC.

1.2. VLMCs: definition of the model

Let $\mathcal{A}$ be a finite set, called the *alphabet*. Here $\mathcal{A}$ will most often be the standard alphabet $\mathcal{A} = \{0, 1\}$, but also $\mathcal{A} = \{d, u\}$ (for *down* and *up*) or $\mathcal{A} = \{n, e, w, s\}$ (for the cardinal directions). Let

$$\mathcal{R} = \{\alpha\beta\gamma\cdots : \alpha, \beta, \gamma, \cdots \in \mathcal{A}\}$$

be the set of *right-infinite* words over $\mathcal{A}$, written by simple concatenation. A VLMC on $\mathcal{A}$, defined below and most often denoted by $(U_n)_{n \in \mathbb{N}}$, is a particular type of $\mathcal{R}$ -valued discrete time Markov chain where:

- the process evolves between time n and time $n + 1$ by adding one letter *on the left* of U_n ;
- the transition probabilities between time n and time $n + 1$ depend on a finite – but not bounded – prefix¹ of the current word U_n .

Giving a formal frame of such a process leads to the following definitions. For a complete presentation of VLMC, one can also refer to Cénac *et al.* (2012).

As usual, a *tree* on $\mathcal{A}$ is a set $\mathcal{T}$ of finite words – namely a subset of $\bigcup_{n \in \mathbb{N}} \mathcal{A}^n$ – which contains the empty word $\emptyset$ (the *root* of $\mathcal{T}$) and which is prefix-stable: for all finite words $u, v, uv \in \mathcal{T} \implies u \in \mathcal{T}$. A tree is made of *internal nodes* ($u \in \mathcal{T}$ is internal when $\exists \alpha \in \mathcal{A}, u\alpha \in \mathcal{T}$) and of *leaves* ($u \in \mathcal{T}$ is a leaf when it has no child: $\forall \alpha \in \mathcal{A}, u\alpha \notin \mathcal{T}$).

DEFINITION 1.1 (Context tree).– A context tree on $\mathcal{A}$ is a saturated tree on $\mathcal{A}$ having an at most countable set of infinite branches.

The tree $\mathcal{T}$ is *saturated* whenever any internal node has $\#(\mathcal{A})$ children: for any finite word u and for any $\alpha \in \mathcal{A}, u\alpha \in \mathcal{T} \implies (\forall \beta \in \mathcal{A}, u\beta \in \mathcal{T})$. A right-infinite word on $\mathcal{A}$ is an *infinite branch* of $\mathcal{T}$ when all its finite prefixes belong to $\mathcal{T}$.

Following the vocabulary introduced by Rissanen, a *context* of the tree is a leaf or an infinite branch. A finite or right-infinite word on $\mathcal{A}$ is an *external node* when it is neither internal nor a context. See Figure 1.1 which illustrates these definitions, as well as the *pref* function defined hereunder.

DEFINITION 1.2 (pref function).– Let $\mathcal{T}$ be a context tree. If w is any external node or any context, the symbol *pref* w denotes the longest (finite or infinite) prefix of w that belongs to $\mathcal{T}$.

¹ In fact, an infinite prefix might be needed in a denumerable number of cases.

In other words, $\text{pref } w$ is the only context c for which $w = c \cdots$. For a more visual presentation, hang w by its head (its left-most letter) and insert it into the tree; the only context through which the word goes out of the tree is its pref.




Figure 1.1. A context tree on the alphabet $\mathcal{A} = \{0, 1\}$. The dotted lines are possibly the beginning of infinite branches. Any word that writes $1000 \cdots$, like the one represented by the dashed line, admits 1000 as a pref. For a color version of this figure, see www.iste.co.uk/barbu/data.zip

With these definitions, it is now possible to define a VLMC.

DEFINITION 1.3 (VLMC).— Let $\mathcal{T}$ be a context tree. For every context c of $\mathcal{T}$, let q_c be a probability measure on $\mathcal{A}$. The VLMC defined by $\mathcal{T}$ and by the $(q_c)_c$ is the $\mathcal{R}$ -valued discrete-time Markov chain $(U_n)_{n \in \mathbb{N}}$ defined by the following transition probabilities: $\forall n \in \mathbb{N}, \forall \alpha \in \mathcal{A}$,

$$\mathbb{P}(U_{n+1} = \alpha | U_n) = q_{\text{pref}(U_n)}(\alpha). \quad [1.1]$$

To get a realization of a VLMC as a process on $\mathcal{R}$, take a (random) right infinite word

$$U_0 = X_0 X_{-1} X_{-2} X_{-3} \cdots$$

At each step of time $n \geq 0$, one gets U_{n+1} by adding a random letter X_{n+1} on the left of U_n :

$$\begin{aligned} U_{n+1} &= X_{n+1} U_n \\ &= X_{n+1} X_n \cdots X_1 X_0 X_{-1} X_{-2} \cdots \end{aligned}$$

under the conditional distribution [1.1].

REMARK 1.1.— Probabilizing a context tree consists, as in definition 1.3, of endowing it with a family of probability measures on the alphabet, indexed by the set of contexts. This vocabulary is used below.

REMARK 1.2.– Assume that the context tree is finite and denote its height by h ; in this condition, the VLMC is just a Markov chain of order h on $\mathcal{A}$. On the contrary, when the context tree is infinite, and this is mainly our case of interest, the VLMC is generally *not* a Markov process on $\mathcal{A}$.

EXAMPLE 1.1.– Take $\mathcal{A} = \{\mathbf{n}, \mathbf{e}, \mathbf{w}, \mathbf{s}\}$ as an (ordered) alphabet, so that the daughters of an internal node are represented, as shown on the left side of Figure 1.2. Making the transition probabilities $\mathbb{P}(U_{n+1} = \alpha U_n | U_n)$ depend only on the length of the largest prefix of the form $\mathbf{n}^k$ ($k \geq 0$) of U_n amounts to taking a comb as a context tree, as shown on the right side of Figure 1.2. Its finite contexts are the $\mathbf{n}^k \alpha$ where $k \geq 0$ and $\alpha \in \mathcal{A} \setminus \{\mathbf{n}\}$.




Figure 1.2. *On the left: how one can represent trees on $\mathcal{A} = \{\mathbf{n}, \mathbf{e}, \mathbf{w}, \mathbf{s}\}$. On the right, the so-called left comb on $\mathcal{A} = \{\mathbf{n}, \mathbf{e}, \mathbf{w}, \mathbf{s}\}$*

EXAMPLE 1.2.– Take again $\mathcal{A} = \{\mathbf{n}, \mathbf{e}, \mathbf{w}, \mathbf{s}\}$ as an alphabet. Making the transition probabilities $\mathbb{P}(U_{n+1} = \alpha U_n | U_n)$ depend only on the length of the largest prefix of the form α^k ($k \geq 1$) of U_n , where α is *any* letter, amounts to taking a *quadruple comb* as a context tree, as shown on the right side of Figure 1.3. In the same vein, if one takes $\mathcal{A} = \{u, d\}$, the *double comb* is the context tree, as shown on the left side of Figure 1.3. In the corresponding VLMC, the transitions depend only on the length of the last current run u^k or d^k , $k \geq 1$. The double comb and the quadruple comb are used below to define PRWs.




Figure 1.3. *The double comb and the quadruple comb*

EXAMPLE 1.3.– Take $\mathcal{A} = \{0, 1\}$ (naturally ordered for the drawings). The left comb of right combs, shown on the left side of Figure 1.4, is the context tree of a VLMC that makes its transition probabilities depend on the largest prefix of U_n of the form $0^p 1^q$.

If one has to take into consideration the largest prefix of the form $0^p 1^q$ or $1^p 0^q$, one has to use the double comb of opposite combs, as shown on the right side of Figure 1.4.




Figure 1.4. Context trees on $\mathcal{A} = \{0, 1\}$: the left comb of right combs (on the left) and a double comb of opposite combs (on the right)

DEFINITION 1.4 (Non-nullness).—A VLMC is called non-null when no transition probability vanishes, i.e. when $q_c(\alpha) > 0$ for every context c and for every $\alpha \in \mathcal{A}$.

Non-nullness appears below as an irreducibility-like assumption made on the driving VLMC of PRWs and for existence and unicity of an invariant probability measure for a general VLMC as well.

1.3. Definition and behavior of PRWs

In this section, the so-called PRWs are defined. A PRW is a random walk driven by some VLMC. In dimensions one and two, results on transience and the recurrence of PRW are given. These results are detailed and proven in Cénac *et al.* (2018b, 2013) in dimension one and in Cénac *et al.* (2020) in dimension two.

1.3.1. PRWs in dimension one

In this section, we deal with one-dimensional PRWs. Note that, contrary to the classical random walk, a PRW is generally not Markovian. Let $\mathcal{A} := \{d, u\} = \{-1, 1\}$ (d for down and u for up) and consider the *double comb* on this alphabet as a context tree, probabilize it and denote by $(U_n)_n$ a realization of the associated VLMC. The n^{th} increment X_n of the PRW is given as the first letter of U_n : define the persistent random walk $S = (S_n)_{n \geq 0}$ by $S_0 = 0$ and, for $n \geq 1$,

$$S_n := \sum_{\ell=1}^n X_\ell, \quad [1.2]$$

so that for any $n \geq 1, m \geq 0$,

$$\mathbb{P}(S_{m+1} = S_m + 1 | U_m = d^n u \dots) = q_{d^n u}(u)$$

$$\mathbb{P}(S_{m+1} = S_m - 1 | U_m = u^n d \dots) = q_{u^n d}(d).$$

Furthermore, for the sake of simplicity and without loss of generality, we condition the walk to start almost surely (a.s.) from $\{X_{-1} = u, X_0 = d\}$ – this amounts to changing the origin of time. In this model, a walker on a line keeps the same direction with a probability that depends on the discrete time already spent in the direction the walker is currently moving (see Figure 1.5). This model can be seen as a generalization of DRRWs introduced in Mauldin *et al.* (1996).




Figure 1.5. A one-dimensional PRW. For a color version of this figure, see www.iste.co.uk/barbu/data.zip

Taking different probabilized context trees would lead to different probabilistic impacts on the asymptotic behavior of resulting PRWs. Moreover, the characterization of the recurrent versus transient behavior is difficult in general. We state here exhaustive recurrence criteria for PRWs defined from a double comb.

In order to avoid trivial cases, we assume that S cannot be frozen in one of the two directions with a positive probability. Therefore, we make the following assumption.

ASSUMPTION 1.1 (Finiteness of the length of runs).— For any $\alpha, \beta \in \{u, d\}$, $\alpha \neq \beta$,

$$\lim_{n \rightarrow +\infty} \left(\prod_{k=1}^n q_{\alpha^k \beta}(\alpha) \right) = 0. \quad [1.3]$$

Let τ_n^u and τ_n^d be, respectively, the length of the n^{th} rise and of the n^{th} descent.

Then, by a renewal-type property (see Cénac *et al.* 2013, proposition 2.3), $(\tau_n^d)_{n \geq 1}$ and $(\tau_n^u)_{n \geq 1}$ are independent sequences of i.d.d. random variables. Their distribution tails are straightforwardly given by: for any $\alpha, \beta \in \{u, d\}$, $\alpha \neq \beta$ and $n \geq 1$,

$$\mathbb{P}(\tau_1^\alpha \geq n) = \prod_{k=1}^{n-1} q_{\alpha^k \beta}(\alpha). \quad [1.4]$$

Note that assumption 1.1 amounts to supposing that the *persistence times* τ_n^d and τ_n^u are a.s. finite. The *jump times* (or *breaking times*) are: $B_0 = 0$ and, for $n \geq 1$,

$$B_{2n} := \sum_{k=1}^n (\tau_k^d + \tau_k^u) \text{ and } B_{2n+1} := B_{2n} + \tau_{n+1}^d. \quad [1.5]$$

In order to deal with a more tractable random walk built with the possibly unbounded but i.d.d. increments $Y_n := \tau_n^u - \tau_n^d$, we introduce the underlying *skeleton* random walk $(M_n)_{n \geq 1}$, which is the original walk observed at the random times of up-to-down turns:

$$M_n := \sum_{k=1}^n Y_k = S_{B_{2n}}. \quad [1.6]$$

Two main quantities play a key role in the asymptotic behavior, namely the expectations of the lengths of runs: with formula [1.4], let

$$\Theta_d := \mathbf{E}[\tau_1^d] = \sum_{n \geq 1} \prod_{k=1}^{n-1} q_{d^k u}(d) \text{ and } \Theta_u := \mathbf{E}[\tau_1^u] = \sum_{n \geq 1} \prod_{k=1}^{n-1} q_{u^k d}(u). \quad [1.7]$$

Actually, Θ_d and Θ_u are already discussed in Cénac *et al.* (2013, proposition B1), where it is shown that the driving VLMC of a one-dimensional PRW admits a unique invariant probability measure if and only if $\Theta_d < \infty$ and $\Theta_u < \infty$.

Note that the expectation of Y_1 is well defined in $[-\infty, +\infty]$ whenever at least one of the persistence times τ_1^u or τ_1^d is integrable. Thus, as soon as $\Theta_d < \infty$ or $\Theta_u < \infty$, let

$$\mathbf{d}_M := \mathbf{E}[Y_1] = \underbrace{\Theta_u - \Theta_d}_{\in [-\infty, +\infty]} \quad [1.8]$$

and

$$\mathbf{d}_S := \frac{\mathbf{E}[\tau_1^u] - \mathbf{E}[\tau_1^d]}{\mathbf{E}[\tau_1^u] + \mathbf{E}[\tau_1^d]} = \frac{\Theta_u - \Theta_d}{\Theta_u + \Theta_d} \in [-1, 1]. \quad [1.9]$$

An elementary computation shows that $\mathbb{E}(M_n) = n\mathbf{d}_M$ and $\mathbb{E}(S_n) \sim n\mathbf{d}_S$ when n tends to infinity. Thus, $\mathbf{d}_M$ and $\mathbf{d}_S$ appear as asymptotic drifts when the walks $(M_n)_n$ and $(S_n)_n$ respectively, turn out to be transient (see Table 1.1). The behavior of the walk also depends on quantities $J_{\alpha|\beta}$, defined for α and $\beta \in \mathcal{A}$, $\alpha \neq \beta$ by:

$$J_{\alpha|\beta} := \sum_{n=1}^{\infty} \frac{n\mathbb{P}(\tau_1^\alpha = n)}{\sum_{k=1}^n \mathbb{P}(\tau_1^\beta \geq k)}.$$

A complete and usable characterization of the recurrence and the transience of the PRW in terms of the probabilities to persist in the same direction or to switch is given in proposition 1.1. Its proof relies on a criterion of Erickson (1973), applied to the skeleton walk $(M_n)_n$, which is simpler to deal with because its increments are independent.

PROPOSITION 1.1.– Under the non-nullness assumption and assumption 1.1, the random walk $(S_n)_n$ is recurrent or transient as described in Table 1.1.

	$\Theta_u < \infty$		$\Theta_u = \infty$	
$\Theta_d < \infty$	Recurrent $\mathbf{d}_S = 0$	Drifting $+\infty$ $\mathbf{d}_S > 0$	Drifting $+\infty$	
		Drifting $-\infty$ $\mathbf{d}_S < 0$		
$\Theta_d = \infty$	Drifting $-\infty$		Recurrent $J_{u d} = J_{d u} = \infty$	Drifting $+\infty$ $\infty = J_{u d} > J_{d u}$ Drifting $-\infty$ $\infty = J_{d u} > J_{u d}$

Table 1.1. *Recurrence versus transience (drifting) for $(S_n)_n$ in dimension one*

The most fruitful situation emerges when both running times τ_1^u and τ_1^d have infinite means. In that case, the recurrence properties of $(S_n)_n$ are related to the behavior of the skeleton random walk $(M_n)_n$ defined in [1.6], the drift of which, $\mathbf{d}_M$, is not defined. Thus, the behavior of $(S_n)_n$ depends on the comparison between the distribution tails of τ_1^u and τ_1^d defined in [1.4], expressed by the quantities $J_{\alpha|\beta}$. Note that the case when both $J_{u|d}$ and $J_{d|u}$ are finite does not appear in the table since it would imply that $\Theta_u < \infty$ and $\Theta_d < \infty$ (see Erickson 1973).

In all three other cases, the drift $\mathbf{d}_S$ is well defined and the PRW is recurrent if and only if $\mathbf{d}_S = 0$. In that case, $\lim_{n \rightarrow \infty} \frac{S_n}{n} = \mathbf{d}_S = 0$. Note that modifying one transition q_c transforms a recurrent PRW into a transient one, since $\mathbf{d}_S$ becomes non-zero.

GONZÁLEZ VELASCO Miguel, del PUERTO GARCÍA Inés, YANEV George P.
Controlled Branching Processes
(*Branching Processes, Branching Random Walks and Branching Particle Fields Set – Volume 2*)

HARLAMOV Boris
Stochastic Analysis of Risk and Management
(*Stochastic Models in Survival Analysis and Reliability Set – Volume 2*)

KERSTING Götz, VATUTIN Vladimir
Discrete Time Branching Processes in Random Environment
(*Branching Processes, Branching Random Walks and Branching Particle Fields Set – Volume 1*)

MISHURA YULIYA, SHEVCHENKO Georgiy
Theory and Statistical Applications of Stochastic Processes

NIKULIN Mikhail, CHIMITOVA Ekaterina
Chi-squared Goodness-of-fit Tests for Censored Data
(*Stochastic Models in Survival Analysis and Reliability Set – Volume 3*)

SIMON Jacques
Banach, Fréchet, Hilbert and Neumann Spaces
(*Analysis for PDEs Set – Volume 1*)

2016

CELANT Giorgio, BRONIATOWSKI Michel
Interpolation and Extrapolation Optimal Designs I: Polynomial Regression and Approximation Theory

CHIASSEIRINI Carla Fabiana, GRIBAUDO Marco, MANINI Daniele
Analytical Modeling of Wireless Communication Systems
(*Stochastic Models in Computer Science and Telecommunication Networks Set – Volume 1*)

GOUDON Thierry
Mathematics for Modeling and Scientific Computing

KAHLE Waltraud, MERCIER Sophie, PAROISSIN Christian
Degradation Processes in Reliability
(*Mathematical Models and Methods in Reliability Set – Volume 3*)

KERN Michel
Numerical Methods for Inverse Problems

RYKOV Vladimir
Reliability of Engineering Systems and Technological Risks
(*Stochastic Models in Survival Analysis and Reliability Set – Volume 1*)

2015

DE SAPORTA Benoîte, DUFOUR François, ZHANG Huilong
Numerical Methods for Simulation and Optimization of Piecewise Deterministic Markov Processes

DEVOLDER Pierre, JANSSEN Jacques, MANCA Raimondo
Basic Stochastic Processes

LE GAT Yves
Recurrent Event Modeling Based on the Yule Process
(*Mathematical Models and Methods in Reliability Set – Volume 2*)

2014

COOKE Roger M., NIEBOER Daan, MISIEWICZ Jolanta
Fat-tailed Distributions: Data, Diagnostics and Dependence
(*Mathematical Models and Methods in Reliability Set – Volume 1*)

MACKEVIČIUS Vigirdas
Integral and Measure: From Rather Simple to Rather Complex

PASCHOS Vangelis Th
Combinatorial Optimization – 3-volume series – 2nd edition
Concepts of Combinatorial Optimization / Concepts and Fundamentals – volume 1
Paradigms of Combinatorial Optimization – volume 2
Applications of Combinatorial Optimization – volume 3