


Edited by
Alfredo Ricci and Luca Bernardi

Methodologies in Amine Synthesis

Challenges and Applications



Methodologies in Amine Synthesis

Methodologies in Amine Synthesis

Challenges and Applications

Edited by Alfredo Ricci and Luca Bernardi

WILEY-VCH

Editors

Prof. Alfredo Ricci

University of Bologna
Department of Industrial Chemistry
“Toso Montanari”
Viale del Risorgimento 4
40136 Bologna
Italy

Prof. Luca Bernardi

University of Bologna
Department of Industrial Chemistry
“Toso Montanari” & INSTM
Research Unit Bologna
Viale del Risorgimento 4
40136 Bologna
Italy

■ All books published by **Wiley-VCH** are carefully produced. Nevertheless, authors, editors, and publisher do not warrant the information contained in these books, including this book, to be free of errors. Readers are advised to keep in mind that statements, data, illustrations, procedural details or other items may inadvertently be inaccurate.

Library of Congress Card No.:
applied for

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library.

Bibliographic information published by the Deutsche Nationalbibliothek

The Deutsche Nationalbibliothek lists this publication in the Deutsche Nationalbibliografie; detailed bibliographic data are available on the Internet at <<http://dnb.d-nb.de>>.

© 2021 WILEY-VCH GmbH, Boschstr. 12, 69469
Weinheim, Germany

All rights reserved (including those of translation into other languages). No part of this book may be reproduced in any form – by photoprinting, microfilm, or any other means – nor transmitted or translated into a machine language without written permission from the publishers. Registered names, trademarks, etc. used in this book, even when not specifically marked as such, are not to be considered unprotected by law.

Print ISBN: 978-3-527-34739-1

ePDF ISBN: 978-3-527-82617-9

ePub ISBN: 978-3-527-82619-3

oBook ISBN: 978-3-527-82618-6

Typesetting SPi Global, Chennai, India

Printed on acid-free paper

10 9 8 7 6 5 4 3 2 1

Contents

Preface *xi*

1	Substitution-type Electrophilic Amination Using Hydroxylamine-Derived Reagents	1
	<i>Zhe Zhou and László Kürti</i>	
1.1	Introduction	1
1.2	Cu-Catalyzed Reactions	2
1.3	Electrophilic Amination Reactions Catalyzed by Other Transition Metals	15
1.4	Electrophilic Amination with Hydroxylamine-derived Metallanitrenes	18
1.5	Transition-Metal-Free Electrophilic Amination Reactions	22
1.6	Conclusion	28
	References	28
2	Remote Functionalizations Using Nitrogen Radicals in H-Atom Transfer (HAT) Reactions	31
	<i>Ji Hye Kim, Elizabeth M. Dauncey and Daniele Leonori</i>	
2.1	Introduction	31
2.2	Intramolecular 1,5-H-Atom Transfer (1,5-HAT)	31
2.3	Photoinduced Strategies	34
2.3.1	Reductive Strategies	34
2.3.1.1	1,5-HAT via Iminyl Radicals	34
2.3.1.2	1,5-HAT via Amidyl and Sulfamidyl Radicals	37
2.3.2	Oxidative Strategies	39
2.3.2.1	1,5-HAT via Iminyl Radicals	39
2.3.2.2	1,5-HAT via Amidyl and Sulfamidyl Radicals	40
2.3.3	Photoinduced Bond Homolysis	43
2.4	Thermal Strategies	46
2.5	Summary and Conclusions	50
	References	50

3	Radical-Based C—N Bond Formation in Photo/Electrochemistry	53
	<i>Binbin Huang, Yating Zhao and Wujiong Xia</i>	
3.1	Introduction	53
3.2	C—N Bond Formation via N-radical Species Addition	54
3.2.1	Radical Addition to C—C Double/Triple Bonds	55
3.2.1.1	Amidyl Radical Addition	55
3.2.1.2	Hydrazonyl Radical Addition	62
3.2.1.3	Aminium Radical Cation Addition	64
3.2.2	Radical Species Addition to Aromatic Rings	71
3.3	Amination via N-atom Nucleophilic Addition	77
3.3.1	Aromatic C(sp ²)—H Bond Amination	77
3.3.2	Olefinic C(sp ²)—H Bond Amination	82
3.3.3	Activated C(sp ³)—H Bond Amination	85
3.3.3.1	Benzylic C(sp ³)—H Bond Amination	85
3.3.3.2	N- α -C(sp ³)—H Bond Amination	87
3.4	Amination via Radical Cross-coupling	90
3.4.1	Aryl C(sp ²)—N Bond Formation via Radical Cross-coupling	90
3.4.1.1	Aryl C(sp ²)—N Bond Formation Using Diarylamines	91
3.4.1.2	Aryl C(sp ²)—N Bond Formation Using Azoles	94
3.4.2	Other C—N Bond Formation via Radical Cross-coupling	96
3.5	Summary and Conclusions	98
	References	100
4	Propargylamines: Recent Advances in Asymmetric Synthesis and Use as Chemical Tools in Organic Chemistry	103
	<i>Fei Zhao, Seong-Heun Kim and Daniele Castagnolo</i>	
4.1	Introduction	103
4.2	Metal-Catalyzed Asymmetric Synthesis of Propargylamines	104
4.2.1	Enantioselective A ³ Coupling	104
4.2.1.1	Enantioselective A ³ Coupling Involving Primary Amines	106
4.2.1.2	Enantioselective A ³ Coupling Involving Secondary Amines	108
4.2.2	Enantioselective Propargylic Amination of Propargylic Esters with Amines	112
4.2.3	Cu-Catalyzed Enantioselective Ring Opening of Alkynyl-Substituted Epoxides/Lactones/Carbonates	116
4.2.4	Enantioselective Addition of Terminal Alkynes to Enamines/Enamides	118
4.2.5	Rh/Ru-Catalyzed Enantioselective Hydrogenation of Alkynyl-Substituted Enamides/Imines	120
4.2.6	Enantioselective C—H Activation: Synthesis of Cyclic Propargylamines	123
4.3	Enzymatic Synthesis of Propargylamines	126
4.4	Photoredox Synthesis of Propargylamines	130
4.5	Organocatalyzed Asymmetric Synthesis of Propargylamines	134
4.6	Propargylamines as Building Blocks in the Synthesis of Heterocycles	137
4.6.1	Synthesis of Pyrroles from Propargylamines	138
4.6.2	Synthesis of Pyrrolines from Propargylamines	139

4.6.3	Synthesis of Pyridines from Propargylamines	140
4.6.4	Synthesis of Quinolines from Propargylamines	142
4.6.5	Synthesis of Oxazoles from Propargylamines	145
4.6.6	Synthesis of Thiazoles from Propargylamines	148
4.7	Conclusions	150
	References	150
5	Transition-Metal-Catalyzed Chiral Amines Synthesis	155
	<i>Anton Cunillera, Carmen Claver, Cyril Godard, Martine Urrutigoity and Philippe Kalck</i>	
5.1	Introduction	155
5.2	Asymmetric Reductive Amination	156
5.3	Asymmetric Hydroamination	162
5.4	Asymmetric Hydroaminoalkylation	168
5.5	Asymmetric Hydroaminomethylation	171
5.6	Coupling on a Chiral Metal Center	176
5.7	Conclusion	177
	References	178
6	Industrial Relevance of Asymmetric Organocatalysis in the Preparation of Chiral Amine Derivatives	187
	<i>Luca Bernardi, Armando Carlone and Francesco Fini</i>	
6.1	Introduction	187
6.2	Organocatalysis in Manufacture: Representative Examples	189
6.3	Case Studies	200
6.3.1	Pregabalin	200
6.3.1.1	Pathway A: Desymmetrization of Glutaric Anhydride	53 202
6.3.1.2	Pathway B: Addition of an Amino α -Carbanion	55 to Michael Acceptors 204
6.3.1.3	Pathway C: Addition of Acetate Enolate Equivalents to Nitroalkene	56 206
6.3.2	Bicyclic α -Amino Acid Core of Telaprevir	214
6.3.3	5-(Trifluoromethyl)-2-isoxazolines as Antipest Agents	223
6.4	Summary and Conclusions	230
	References	231
7	Biocatalytic Synthesis of Chiral Amines Using Oxidoreductases	243
	<i>Sebastian C. Cosgrove, Jeremy I. Ramsden, Juan Mangas-Sanchez and Nicholas J. Turner</i>	
7.1	Introduction	243
7.2	Amine Oxidases	243
7.2.1	Introduction	243
7.2.2	(S)-Selective Amine Oxidases	244
7.2.2.1	Monoamine Oxidase from <i>Aspergillus niger</i>	244
7.2.2.2	Directed Evolution of MAO-N	244
7.2.2.3	Synthetic Applications and Cascades	247
7.2.2.4	Monoamine Oxidase from <i>Pseudomonas monteilii</i> ZMU-T01	249

7.2.2.5	Cyclohexylamine Oxidase from <i>Brevibacterium oxydans</i> (CHAO)	249
7.2.3	(R)-Selective Amine Oxidases	252
7.2.3.1	D-Amino Acid Oxidase (pkDAO)	252
7.2.3.2	6-Hydroxy-D-nicotine Oxidase (6-HDNO) from <i>Arthrobacter nicotinovorans</i>	252
7.3	Amine Dehydrogenases	252
7.3.1	Introduction	252
7.3.2	Discovery and Engineering of AmDH	253
7.3.2.1	Leucine Dehydrogenase	253
7.3.2.2	Phenylalanine Dehydrogenase and Chimeric Amine Dehydrogenase	254
7.3.2.3	Native Amine Dehydrogenase	254
7.3.3	Synthetic Applications of AmDH	256
7.3.3.1	Primary Amine Synthesis with Engineered AmDH	256
7.3.3.2	Primary Amine Synthesis with Natural AmDH	258
7.3.3.3	Substrate Promiscuity in AmDH	259
7.3.3.4	Cascade Reactions that Use AmDH	260
7.4	Imine Reductases	262
7.4.1	From Biosynthesis to Biocatalysis	262
7.4.2	Biocatalytic Application of Imine Reductases	264
7.4.2.1	IREDs in Cascade and Chemoenzymatic Synthesis	265
7.4.3	IREd Engineering	267
7.4.4	Imine Reductases Catalyzing Reductive Amination	267
7.4.5	Imine Reductase-Catalyzed Amine Alkylation Cascades	270
7.4.6	Engineering of Reductive Aminases	270
7.5	Engineered Cytochrome P450s	270
7.6	Conclusions and Perspectives	274
	References	274

8 Engineering Functional Nanomaterials Through the Amino Group 285

Giacomo Filippini, Paolo Pengo, Susanna Bosi, Giulio Ragazzon, Lucia Pasquato and Maurizio Prato

8.1	Abbreviations	285
8.1	Introduction	287
8.2	Quantification of Nanomaterial-Bound Amino Groups	288
8.3	Exploiting Amino Compounds for the Functionalization of Carbon-Based Nanomaterials	290
8.3.1	Historical Backgrounds: Allotropes of Carbon	290
8.3.2	Use of Amines for the Functionalization of Carbon Nanostructures	290
8.3.3	Other Functionalization Procedures of Common Carbon Nanostructures	297
8.3.4	Exfoliation of Graphite with Melamine	299
8.3.5	Other Carbon Nanomaterials	301
8.3.5.1	Carbon Nanohorns	301
8.3.5.2	Carbon Nanodiamonds	305
8.3.5.3	Carbon Nano-onions	307

8.3.6	Amino-Functionalized Carbon-Based Nanomaterials for Analytical Applications	308
8.4	Amines in the Synthesis and Functionalization of Carbon Dots	309
8.4.1	Amines as CD Constituents	310
8.4.2	Amine-Rich CDs from Arginine and Ethylenediamine (NCDs)	312
8.4.2.1	One-Pot Functionalization of NCDs	312
8.4.2.2	Postfunctionalization of NCDs	313
8.4.2.3	Use of CD-Supported Amines in Organocatalysis	315
8.5	Amines for the Engineering of Hybrid Organic–Inorganic Nanomaterials	316
8.5.1	Amines as Head Groups or End Groups on Self-assembled Monolayers on Flat Surfaces	316
8.5.2	Alkylamines in the Preparation of Semiconductor Quantum Dots	319
8.5.2.1	Sulfur–Amine and Selenium–Amine Systems	319
8.5.2.2	Capping Ligands for Quantum Dots and Ligand Exchange by Amines	321
8.5.3	Alkylamines as Reagents for the Synthesis and Passivation of Metal Nanoparticles	322
8.5.3.1	Alkylamines as Capping Agents for Metal Nanoparticles	322
8.5.3.2	Displacement of Amines from the Surface of Metal Nanoparticles	325
8.5.4	Amines on the Outer Surface of Organic–inorganic Hybrid Nanoparticles	325
8.5.5	Postfunctionalization of Amine-Terminated Organic–Inorganic Hybrid Nanoparticles	328
	References	329

9 Recent Advances in the Synthesis of Nitrogen Compounds from Biomass Derivatives 341

Ana C. Fernandes

9.1	Introduction	341
9.2	Synthesis of Nitrogen Compounds from Chitin and Its Derivatives	341
9.3	Synthesis of Amines and Formamides from α -Amino Acids	345
9.4	Synthesis of Nitrogen Compounds from Cellulosic Biomass Derivatives	348
9.5	Synthesis of Nitrogen Compounds from Lignin Derivatives	366
9.6	Synthesis of Nitrogen Compounds from Triglycerides and Fatty Alcohols	370
9.7	Conclusion	373
	References	373

10 Recent Advances in the Synthesis of Arylamines in the Light of Application in Pharmaceutical and Chemical Industry 377

Dino Berthold, Alexander M. Haydl, Joyce C. Leung, Ulrich Scholz, Qing Xiao and Zhibin Zhu

10.1	Modern Approaches to Transition-Metal-Catalyzed C–N Coupling in Industry	377
10.1.1	Introduction	377
10.1.2	Transition-Metal-Catalyzed C–N Bond Formation	377
10.1.2.1	Ullmann-Type Amination	378
10.1.2.2	Buchwald–Hartwig Amination	383

10.2	New Methodologies in the Synthesis of Arylamines on the Brink of Industrial Application	388
10.2.1	Introduction	388
10.2.2	Catalytic C–H Amination	389
10.2.2.1	Catalytic C–H Amination under Standard Conditions	389
10.2.2.2	Photoredox Catalysis	399
10.2.2.3	Electrochemical Approaches	402
10.2.3	Decarboxylative Aryl Amination	405
10.2.4	Nickel-Catalyzed C–N Coupling	407
10.2.5	Other Metal-Catalyzed Cross-Couplings	414
10.2.6	Reductive Amination	417
10.2.7	Hydroamination	419
10.2.8	Summary and Conclusions	424
10.3	Advances to Arylamine Formation Using Intensified and More Sustainable Process Technologies	424
10.3.1	Introduction	424
10.3.2	Flow Chemistry	425
10.3.2.1	Pd-Catalyzed C–N Bond Forming Reaction	425
10.3.2.2	Nucleophilic Aromatic Substitution	426
10.3.2.3	Telescoped Sequence of Nitration and Hydrogenation in Flow Synthesis	428
10.3.2.4	Chan–Lam Coupling	428
10.3.3	Immobilization of Catalysts/Supported Catalysts	429
10.3.4	Personal Accounts from Contract Manufacturing Companies on Utility of Modern Flow Amination Methods	431
10.4	Miscellaneous Aspects of Aromatic Amination Reactions in the World of Active Pharmaceutical Ingredients	431
10.4.1	Cohort of Concerns and its Regulatory Impact on Amine-Based Active Pharmaceutical Compounds	431
10.4.2	Role of Control of Elemental Impurities in Human Pharma Applications	432
10.4.3	Transition Metal Accounting	433
10.4.4	Recycling of Metals, Ligands, and Other Cost Drivers of Aromatic Amination	434
10.4.5	Regulatory Requirements for the Submission of a Catalytic Reaction in a New Drug Application	434
	References	435

Preface

The books “Modern Amination Methods” and “Amino Group Chemistry – From Synthesis to the Life Sciences” edited by one of us for Wiley-VCH in 2000 and in 2007, respectively, were intended to provide the reader with exhaustive overviews of the most advanced methodologies for the C—N bond formation and with the role played by the amino function in those processes that are closely related to the life sciences.

These books were well received by the chemical community and highlighted the importance of keeping scientists to be continuously aware of the progress in the field of amino group chemistry. In 2018, setting up to discuss the prospect of coediting a new book on amino group chemistry, we asked ourselves if the past decade witnessed sufficient breakthroughs to make such update worth to be read by academicians or industrialists. It did not take too much time to convince ourselves that such endeavor would have been worthwhile and timely. In the past years, most of the breakthrough discoveries in synthetic organic chemistry embed efficient preparations of nitrogen-containing compounds. Arguably, amino group chemistry lies at the core of recent methodological trends. Furthermore, the disclosure of new materials, from medicine to nanoscience, has often been grounded on nitrogen derivatives. Thus, we embarked in this adventure, approaching the selection of the topics with an unbiased and wide-range attitude. In this new book, we aimed at providing not only an overview over specific aspects of amino chemistry but also a unique journey through modern chemistry.

With the very useful insights into the topic selection provided by Wiley anonymous reviewers, we commenced to approach several prospective authors. We have been very lucky to find contact authors among the most authoritative of their fields, who, together with their coworkers, assembled chapters characterized by clear structures, schemes, mechanisms, and figures accompanying the text together with exhaustive and up-to-date reference sections.

Throughout the 10 chapters of the book, amines are discussed with respect to the most advanced methodologies for their preparation, from design to scope surveys to applications to biologically active targets or precursors thereof, including large-scale industrial settings. Because the pharmaceutical and agrochemical industries are nowadays mostly away from the development of racemic compounds, several of the book chapters refer to enantioselective synthesis of chiral amines. It is worth noting that the contributions to this book present interlinks that, as a fil rouge crossing the whole book, allow the reader to face some of the topics under different perspectives.

The book starts with a topic that constitutes a continuum between the 2000s monographies and this one: electrophilic amination. Indeed, substitution-type electrophilic amination has gained a tremendous progress in the decade. This unconventional, yet synthetically appealing approach to C—N bond formation is nowadays performed under transition metal (TM)- and metal-free catalysis aiming to employ bench-stable and easily obtainable reagents for forging new C—N bonds. In this context, Chapter 1 summarizes the most recent achievements complying with the requirements for an ideal simple and versatile synthetic methodology.

The following part of the book brings the reader in the domain of nitrogen radicals as reactive intermediates, a reactivity stream displaying high complementarity to classical ionic and TM mediated processes. The use of photo- and electrochemical induced catalytic strategies is widely discussed (Chapters 2 and 3), overviewing the recent dramatic advances enabling to couple challenging C—H activation processes with C—N bond forming events. Although aminations with high enantioselectivity are yet rarely accessed, these radical methodologies targeting the synthesis of a wide range of heterocyclic systems and an array of complex N-containing structures outline a new and highly promising frontier.

As an enabling methodology, the development and application of asymmetric catalysis is now an integral part of the work of any major research unit in academia and industry. In the past decade, organo- and biocatalysis have flanked TM catalysis as key players for the enantioselective synthesis of chiral amino compounds, as overviewed in the ensuing Chapters 4–7. Worth noting is the possibility that these various catalytic approaches might concur toward attaining the same targets. This is exemplified in Chapter 4 devoted to the synthesis of chiral propargylamines. Many of these compounds possessing pharmaceutical properties have found applications in the treatment of diseases and can be considered among the most efficient building blocks for the synthesis N-containing heterocycles.

The relevance of high atom efficiency and very limited waste production (high atom economy) is of paramount importance for the development of methodologies with industrial prospects. Reactions such as TM-catalyzed asymmetric hydroamination, reductive amination, and hydroaminomethylation, reported in Chapter 5, meet these goals, an additional advantage being the simplicity and the easy availability of the starting materials, which is only partially counterbalanced by the complexity of the used ligands. Because of their relevance, these reactions are occasionally mentioned elsewhere in this book, although in the frame of different contexts, thus providing the reader with an exhaustive set of complementary information.

In the following chapters, metal-free stereoselective catalytic methodologies at the service of amine synthesis, represented by organo- and biocatalysis, are treated with a greater focus on industrial applications reflecting the dynamic nature of these catalytic streams. The central goal of organocatalysis (Chapter 6) is herein addressed to the emerging utilization of this methodology as an exceedingly useful synthetic tool for the large-scale preparation of active pharmaceutical ingredients (APIs) in industrial settings. An emphasis is placed on three case studied in which organocatalysis stands out with respect to other enantioselective strategies, for its versatility and potential for scale-up. The revolutionary approach of enzyme engineering, in the context of oxidoreductase enzymes applied to the synthesis of N-containing biologically relevant intermediates and products, is described in Chapter 7.

The intrinsic selectivity of enzymes provides an obvious advantage in the key challenge of developing new synthetic platforms amenable to industrialization.

The following two chapters stand out for their uniqueness, targeting topics that were not even touched in the previous two books on amino chemistry but of exceptional prominence and timeliness. Chapter 8 deals with the use of amines in the synthesis, stabilization, and functionalization of organic–inorganic hybrid nanomaterials, highlighting through a large number of case studies the pivotal role played by the amino group to unlock practical applications in the fields of biology, medicine, and energy production. Conversely, the synthesis of a number of valuable amino compounds is reported in Chapter 9, from the transformation of renewable biomass resources that already incorporate the amino groups such as chitin, chitosan, and amino acids or by modifying bio-based compounds, followed by amination. Rare aminosugars, precursors of medicinal compounds, and a wide range of heterocycles are obtained avoiding the use of fossil-based feedstocks, thus providing a remarkable step forward in the ongoing shift from depleting to renewable resources.

The final Chapter 10 addresses current applications of TM-catalyzed aromatic amination in industrial settings by discussing a large number of case studies related to the manufacturing process of pharmaceutical compounds. In addition, with reference to the seminal work by Ullmann, Buchwald, and Hartwig, this contribution points on new concepts still at academic level, but either further extending the applicability of new methodologies, or on the brink of being industrially used. Also approaches with a focus on process intensification and sustainability (flow chemistry and catalyst immobilization) are presented, together with a view of the accompanying questions when applying the methodology of aromatic amination in the pharmaceutical industry. To make aware the reader about these challenges, themes such as the control of elemental impurities, the TM accounting, and the metal recycling are treated as well. Besides being a highly useful and up-to-date source of information on the TM-catalyzed aromatic amination in industry, this contribution will hopefully provide inspiration for academic research in developing new methodologies amenable to industrialization.

We warmly thank all the distinguished scientists and their coauthors for their rewarding and highly instructive contributions. Without their effort, even more valuable considering it partially coincided with a difficult period at the international level, this volume would have not been possible. Grateful acknowledgments are also addressed to the Wiley-VCH editorial staff, and in particular to Anne Brennführer, Aruna Pragasam, Elke Maase, and Katherine Wong, who encouraged us at project outset and helped us in a very competent manner in all the phases of the preparation of this book.

Bologna
07 April 2020

Alfredo Ricci and Luca Bernardi

1

Substitution-type Electrophilic Amination Using Hydroxylamine-Derived Reagents

Zhe Zhou and László Kürti


Rice University, Department of Chemistry, 6500 Main Street, Houston, TX 77030, USA

1.1 Introduction

Electrophilic amination is a class of organic reactions where C—N bonds are formed via the use of electrophilic aminating reagents [1–4]. Depending on the specific reaction pathway, electrophilic amination reactions can be classified either as substitution or addition. This chapter focuses on substitution reactions. Aminating reagents for the substitution-type electrophilic aminations are essentially NR_2^+ synthons, and the electrophilicity on the nitrogen atom is generally achieved by attaching a more electronegative functionality (X) to the nitrogen atom that can serve as a leaving group. Common structural motifs for this class of reagents include chloramines, hydroxylamines, and oxaziridines (i.e. cyclic hydroxylamine derivatives). Because of the safety hazards associated with the use of chloramines, recent developments in this area have been focused on the use of more stable hydroxylamine-type reagents (Scheme 1.1).


Substitution-type electrophilic amination reactions can operate under either uncatalyzed or catalyzed conditions. The majority of catalytic substitution-type electrophilic amination reactions are catalyzed by complexes of transition metals (TMs). In the uncatalyzed reactions, the carbon nucleophile directly attacks the electrophilic nitrogen atom and a new C—N bond is formed. In the TM-catalyzed reactions, the transition metal first enters into the N—X bond via an oxidative addition, and the new C—N bond is formed after sequential ligand exchange and reductive elimination (Scheme 1.2). The major difference between TM-catalyzed substitution-type electrophilic amination reactions and TM-catalyzed C—N cross-coupling reactions (i.e. Buchwald–Hartwig coupling) is the role of the nitrogen source: it acts as an electrophile in the former while as a nucleophile in the latter.

The majority of the literature in this area concerns the TM-catalyzed versions of substitution-type electrophilic amination. Therefore, they will be discussed first in this chapter.




Scheme 1.1 General structure of electrophilic aminating reagent.

Mechanism of uncatalyzed electrophilic amination



Mechanism of TM-catalyzed electrophilic amination




Scheme 1.2 Mechanisms of two main types of electrophilic amination.


1.2 Cu-Catalyzed Reactions

Narasaka and coworkers reported an early iteration of Cu-catalyzed substitution-type electrophilic amination of Grignard reagents utilizing *O*-sulfonyloximes as aminating reagents. The reactions can also give products without copper catalysis, albeit with much lower yields. Because of the nature of the aminating reagents, subsequent acidic hydrolysis is needed to convert the imine products to the desired amines [5, 6]. A similar amination process of organozinc reagents was reported by the Erdik group around the same time. In this case, in addition to the *O*-sulfonyloximes, the authors showed that the reaction can also proceed with methoxyamine when excess of the organometallic reagent was used (Scheme 1.3) [7, 8]. These early reactions have several drawbacks that affect their utilizations, including a very limited substrate scope, low conversion rate due to many side reactions, and, most importantly, the need to use a strong acid to hydrolyze the initial imine

Narasaka et al. [5,6]




Erdik et al. [7,8]



M = ZnCl, ZnBr, 1/2 Zn, 1/3 ZnMgBr

Scheme 1.3 Early examples of Cu-catalyzed electrophilic amination. Source: Erdik and Ay [2] and Tsutsui et al. [5].




Scheme 1.4 Cu-catalyzed electrophilic amination of organozinc reagents.

products. The use of strong acidic conditions in the hydrolysis makes these procedures unsuitable for substrates containing acid-labile functionalities.

In 2004, the Johnson group at UNC reported the Cu-catalyzed electrophilic amination of diorganozinc reagents using acyl hydroxylamines as aminating reagents [9]. This is the first report of Cu-catalyzed electrophilic amination reactions that give tertiary amines as products (Scheme 1.4). Compared to the *O*-sulfonyloximes, *O*-acyl hydroxylamines are more synthetically accessible and have better atom economy. A limitation, however, is that the nitrogen must be fully substituted (i.e. no acidic N—H bond is tolerated). In place of the diorganozinc substrate, this reaction can also use a Grignard reagent as the nucleophile, which is arguably more accessible and convenient to use [10].

Greaney and coworkers [11]

**Scheme 1.5** Cu-catalyzed electrophilic amination via aryne intermediate.


After the initial disclosure of Johnson and coworkers, subsequent research showed that other organometallic reagents can also serve as the nucleophile.

A unique reaction was reported by Greaney and coworkers at the University of Manchester in the UK [11]. In this case, an aryne intermediate is first generated in situ via iodine–magnesium exchange, followed by elimination. A nucleophile subsequently attacks the aryne, and the resulting arylmetal undergoes Cu-catalyzed electrophilic amination with the hydroxylamine-derived electrophilic aminating reagent to furnish aryl amines as the final products (Scheme 1.5). This reaction gives access to unique aromatic products with a 1,2-bis substitution pattern. The nucleophiles that can be used in this reaction include arylamines, thiophenols, and arylselenenols. The overall transformation exhibits excellent regioselectivity; however, the isolated yield is lowered in those cases in which the reactants are sterically encumbered.

One important consideration in these types of reactions is that the nucleophile has to be able to coexist with the aminating reagent in order for the catalytic cycle to be established. Otherwise, the nucleophile may directly react (i.e. protonation and/or uncatalyzed nucleophilic attack) with the aminating reagent and may lead to the formation of undesired by-products. This puts limitations on both the nucleophilicity and basicity of the organometallic nucleophiles. Organozinc compounds, Grignard reagents, and organoaluminum [12] reagents have been shown to be suitable nucleophiles for these types of reactions, while organolithium reagents are generally unsuitable because of their strong basicity and tendency to participate in side reactions that involve electron transfer pathways.

To address this issue and provide a solution for the direct electrophilic amination of organolithium reagents via copper catalysis, the group of A. B. Smith III (University of

Smith and coworker [13]


**Scheme 1.6** Cu-catalyzed electrophilic amination of organolithium reagents.

Pennsylvania, 2013) developed a protocol in which a siloxane transfer reagent is used to modulate the reactivity of the organolithium reagents (Scheme 1.6) [13]. The siloxane reagent acts as an “attenuator” for the more reactive organolithium reagents and lowers its reactivity by forming a less basic complex that can more readily participate in the catalytic cycle and, in the meantime, prevents the direct attack on the aminating reagent by the organolithium.

This type of C—N bond-forming reaction can also take place directly between cuprates and electrophilic aminating reagents. One reason to justify the use of stoichiometric amounts of a copper salt (i.e. full transmetalation) is that the low basicity of the resulting cuprate can tolerate the presence of N—H protons in the aminating reagent, thereby enabling the synthesis of primary and secondary amine products without the use of excess organometallic reagent. The Kürti group has demonstrated this possibility with the use of sterically hindered N—H oxaziridines (Scheme 1.7) [14]. Compared to the tertiary amines, unprotected primary amines are more versatile building blocks for further functionalization.

These sterically hindered N—H oxaziridines can be readily synthesized on multigram scale from the corresponding N—H imines and *meta*-chloroperbenzoic acid (*m*CPBA). The N—H oxaziridines are bench-stable compounds and can also be readily purified via flash column chromatography and using regular silica gel as the stationary phase. The steric hindrance created by the bulky alkyl groups reduces the kinetic acidity of the oxaziridine N—H bond, thus allowing the cuprates to be aminated as opposed to suffering unproductive proton transfer.


Kürti and coworkers [14]

**Scheme 1.7** Electrophilic amination of arylcuprates using an NH-oxaziridine.

Since the advent of direct C–H cupration of arenes, it is now also possible to utilize cuprate nucleophiles without first going through a separate transmetalation step. Uchiyama and coworkers (at RIKEN, Japan) showed that it was indeed possible to directly aminate aryl cuprates with *O*-benzyl hydroxylamine. The directed C–H cupration of arenes was achieved using a strong base $(\text{TMP})_2\text{Cu}(\text{CN})\text{Li}_2$, which can selectively deprotonate at the ortho position of the amide directing group. The resulting aryl cuprates can be directly primary aminated with benzyl hydroxylamine (Scheme 1.8) [15]. The hygroscopic nature of *O*-benzyl hydroxylamine necessitates its use as a stock solution in an organic solvent.


The instability of organometallic reagents used in the aforementioned reactions imposes limits on their widespread utilization, especially in an industrial setting. Efforts have been made to replace the air- and moisture-sensitive organometallic reagents with more stable alternatives that can be conveniently stored and used. Miura and coworkers (Osaka University, Japan) have shown that both arylboronates (Scheme 1.9) [16] and arylsilanes (Scheme 1.10) [17] can serve as starting materials in Cu-catalyzed electrophilic amination reactions. These reactions can proceed under ambient temperature and furnish the corresponding anilines in good to excellent isolated yields. The enhanced stability of arylboronates and arylsilanes reduces the complexity of the operation and, at the same

Uchiyama and coworkers [15]



Scheme 1.8 Electrophilic amination via directed C–H cupration.

Miura and coworkers [16]




Scheme 1.9 Cu-catalyzed electrophilic amination of arylboronates.

time, the wide commercial availability of arylboronates also adds convenience to these types of reactions.

Hirano and Miura discovered that ambident nucleophiles such as silyl ketene acetals are also suitable nucleophiles for the Cu-catalyzed electrophilic amination reactions. These reactions result in the formation of α -amino esters as products. The first generation of this reaction uses chloramines as aminating reagents [18], while the second generation

Miura and coworkers [17]



Scheme 1.10 Cu-catalyzed electrophilic amination of aryl silanes. Source: Modified from Miki et al. [17].

can proceed with the more stable and much safer *O*-benzoyl hydroxylamines (Scheme 1.11) [19]. This method provides a potential route for the syntheses of unnatural as well as modified natural amino acids.

Weak nucleophiles such as styrenes and some electron-deficient heterocycles can also participate in Cu-catalyzed electrophilic amination reactions.


In the case of styrenes, the substrates can undergo hydroamination or aminoboration depending on the specific reaction conditions. Hirano, Miura, and coworkers have demonstrated that styrenes can be stereoselectively functionalized with benzoyl hydroxylamine and bis(pinacolato)diboron under Cu catalysis (Scheme 1.12) [20]. The resulting products can further participate in transition-metal-catalyzed cross-coupling reactions.

When polymethylhydrosiloxane (PMHS) is used instead of bis(pinacolato)diboron, hydroamination products can be obtained under similar reaction conditions (Scheme 1.13). In these cases, it is proposed that the reaction proceeds with an initial CuH addition across the C—C bond of the olefins, followed by the electrophilic amination of the resulting cuprates [21].

With chiral ligands, the hydroamination reactions can give enantiomerically enriched products. Both the Miura (Scheme 1.14) and Buchwald (Scheme 1.15) groups have developed conditions using chiral phosphine ligands [21, 22].


Buchwald and coworkers have also reported the hydroamination of aryl acetylenes. The reaction is highly stereoselective, giving *E*-enamines as the major products. The enamine products can be further reduced to give alkyl amines, which are important building blocks in organic synthesis (Scheme 1.16) [23].

Miura and coworkers [19]




Scheme 1.11 Cu-catalyzed electrophilic amination of silyl enol ethers. Source: Modified from Matsuda et al. [19].

Miura and coworkers




Scheme 1.12 Cu-catalyzed electrophilic catalyzed aminoboration of styrenes. Source: Modified from Matsuda et al. [20].


Miura and coworkers [21]

**Scheme 1.13** Cu-catalyzed electrophilic hydroamination of styrenes. Source: Modified from Miki et al. [21].


Miura and coworkers [21]

**Scheme 1.14** Enantioselective Cu-catalyzed electrophilic hydroamination of styrenes. Source: Miki et al. [21].


Buchwald and coworkers [22]

**Scheme 1.15** Enantioselective Cu-catalyzed electrophilic hydroamination of styrenes. Source: Modified from Zhu et al. [22].

Buchwald and coworker [23]

**Scheme 1.16** Cu-catalyzed electrophilic amination of alkynes. Source: Shi and Buchwald [23].

Miura and coworkers [24]




Scheme 1.17 Cu-catalyzed annulative electrophilic amination. Source: Modified from Matsuda et al. [24].

ortho-Alkynyl phenols and anilines can also undergo annulative amination with electrophilic aminating reagents under Cu catalysis. Miura and coworkers have developed conditions for the synthesis of aminated benzofurans and indoles (Scheme 1.17) [24]. The transformation is operationally simple and proceeds at room temperature. The mechanism was probed and the authors concluded that the most plausible pathway is a nonradical electrophilic amination of the heteroarylcuprate species in the C—N bond-forming step.

Similar intramolecular reactions can also take place with substrates containing unactivated terminal alkenes. In 2015, the Wang group (Duke University) reported the copper-catalyzed vicinal diamination of unactivated alkenes with hydroxylamines that is both regio- and stereoselective. The first iteration of this reaction takes place on unsaturated amides and gives 4-amino-2-pyrrolidones as the products (Scheme 1.18) [25]. This transformation is considered to be the first metal-catalyzed alkene 1,2-diamination that enables the direct incorporation of an electron-rich amino group.

In 2016, Wang and coworkers successfully expanded the substrate scope to include unsaturated carboxylic acids, which undergo amino-lactonization under the reaction conditions (Scheme 1.19) [26]. The overall transformation allows the practitioner to access quickly and efficiently a wide range of amino-substituted γ - and δ -lactones as well as 1,2-amino alcohol derivatives, which are of significant value in the synthesis of natural products and active pharmaceutical ingredients.


Wang [25]

**Scheme 1.18** Cu-catalyzed electrophilic diamination. Source: Modified from Shen and Wang [25].


An unusual case of ring-opening amination of cyclopropanols has been reported by the Dai group [27]. In this reaction, a base-initiated ring-opening of cyclopropanol generates a carbanion nucleophile, which participates in the Cu-catalyzed electrophilic amination and affords β -aminoketones as products (Scheme 1.20). The catalytic cycle involves the oxidation of the Cu(I) complex to the corresponding Cu(III) species by the hydroxylamine reagent. Next, the Cu(III) intermediate promotes the ring-opening of the cyclopropanol substrate and the resulting copper-homoenolate undergoes reductive elimination to form the new C—N bond and to regenerate the catalytically active Cu(I) species. Overall, the transformation proceeds under mild reaction conditions and it is also compatible with a number of sensitive functionalities such as esters, epoxides, and unsaturated carbonyl compounds.

With electron-deficient arenes, direct C—H amination is also possible. Miura and coworkers have reported the Cu-catalyzed direct C—H amination using benzoyl hydroxylamines as aminating reagents (Scheme 1.21) [28]. Electron-deficient aromatic substrates such as fluoroarenes, oxadiazoles, and thiazoles can be directly aminated to furnish the corresponding aryl and heteroaryl amines. The Yotphan group later expanded the substrate scope to include benzoxazoles [29], while the Li group applied the reaction to enable the C—H amination of quinoline *N*-oxide [30].

Wang and coworkers [26]

**Scheme 1.19** Cu-catalyzed electrophilic amino-lactonization. Source: Modified from Hemric et al. [26].

Dai and coworkers [27]

**Scheme 1.20** Cu-catalyzed ring-opening amination. Source: Modified from Ye and Dai [27].