

CLINICAL CASES IN PARAMEDICINE

EDITED BY

SAM WILLIS | IAN PEATE | ROD HILL



WILEY Blackwell

Clinical Cases in Paramedicine



Dedication

This text is being finalised for publication during the COVID-19 global pandemic. Many healthcare workers have lost their lives during this time while looking after those who contracted the SARS-CoV-2 virus. Therefore this text is dedicated to all those healthcare workers and paramedics who continue to look after the sick when faced with adversity and at risk to their own lives.

Also in memory of Brian Mfula, lecturer in mental health nursing at Swansea University, Wales, UK and co-author of Chapter 12, who tragically lost his life to COVID-19.

Clinical Cases in Paramedicine

Edited by

Sam Willis

Lecturer in Paramedicine
Curtin University
Western Australia
Australia

Ian Peate

Head of School, School of Health Studies
Gibraltar Health Authority, St Bernard's Hospital
Gibraltar

Rod Hill

Head of School of Biomedical Sciences
Charles Sturt University
NSW, Australia

WILEY Blackwell

This edition first published 2021
© 2021 John Wiley & Sons Ltd

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by law. Advice on how to obtain permission to reuse material from this title is available at <http://www.wiley.com/go/permissions>.

The right of Sam Willis, Ian Peate and Rod Hill to be identified as the author(s) of the editorial material in this work has been asserted in accordance with law.

Registered Office(s)

John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, USA
John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Office

9600 Garsington Road, Oxford, OX4 2DQ, UK

For details of our global editorial offices, customer services, and more information about Wiley products visit us at www.wiley.com.

Wiley also publishes its books in a variety of electronic formats and by print-on-demand. Some content that appears in standard print versions of this book may not be available in other formats.

Limit of Liability/Disclaimer of Warranty

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting scientific method, diagnosis, or treatment by physicians for any particular patient. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. While the publisher and authors have used their best efforts in preparing this work, they make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives, written sales materials or promotional statements for this work. The fact that an organization, website, or product is referred to in this work as a citation and/or potential source of further information does not mean that the publisher and authors endorse the information or services the organization, website, or product may provide or recommendations it may make. This work is sold with the understanding that the publisher is not engaged in rendering professional services. The advice and strategies contained herein may not be suitable for your situation. You should consult with a specialist where appropriate. Further, readers should be aware that websites listed in this work may have changed or disappeared between when this work was written and when it is read. Neither the publisher nor authors shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

Library of Congress Cataloging-in-Publication Data

Names: Willis, Sam, 1978– editor. | Peate, Ian, editor. | Hill, Rod (Rodney), editor.

Title: Clinical cases in paramedicine / edited by Sam Willis, Ian Peate, Rod Hill.

Description: Hoboken, NJ : Wiley-Blackwell, 2021. | Includes bibliographical references and index.

Identifiers: LCCN 2020026984 (print) | LCCN 2020026985 (ebook) | ISBN 9781119619253 (paperback) | ISBN 9781119621041 (adobe pdf) | ISBN 9781119621034 (epub)

Subjects: MESH: Emergency Treatment | Emergency Medical Technicians | Emergencies | Problems and Exercises | Case Reports

Classification: LCC RC86.9 (print) | LCC RC86.9 (ebook) | NLM WB 18.2 | DDC 616.02/5076–dc23

LC record available at <https://lccn.loc.gov/2020026984>

LC ebook record available at <https://lccn.loc.gov/2020026985>

Cover Design: Wiley

Cover Image: Courtesy of Sarah Angelou

Set in 9/12pt Minion by SPi Global, Pondicherry, India

Contents

Preface, vi

List of contributors, vii

Chapter 1 Respiratory emergencies, 1

Jennifer Stirling, Clare Sutton and Georgina Pickering

Chapter 2 Cardiac emergencies, 25

Michael Porter and Joel Beake

Chapter 3 Neurological emergencies, 47

Kristina Maximous

Chapter 4 Abdominal emergencies, 74

Tania Johnston and Mark Hobson

Chapter 5 Palliative and end-of-life care, 99

Alisha Hensby and Samantha Sheridan

Chapter 6 Medical emergencies, 118

Tom E. Mallinson

Chapter 7 Non-technical skills, 142

Georgette Eaton

Chapter 8 Trauma cases, 161

Tom E. Mallinson and Fenella Corrick

Chapter 9 Paediatric cases, 186

Erica Ley

Chapter 10 Patient-centred care in complex cases, 203

Yasar Gunaratne and David Krygger

Chapter 11 Legal and ethical cases, 225

Ruth Townsend

Chapter 12 Mental health cases, 242

David Davis, Tom Hewes, Lynne Walsh and Brian Mfula

Chapter 13 Older adults, 261

Sam Taylor

Chapter 14 Obstetric cases, 280

Aimee Yarrington

Chapter 15 Remote area cases, 301

Steve Whitfield and Kerryn Wratt

Chapter 16 Mining emergencies, 323

Paul Grant and Curtis Northcott

Index, 348



Preface

Paramedicine is a fast-paced, ever-changing profession and those who practise as paramedics must be able to keep up to date with changes using the latest evidence and expert opinion where evidence does not exist. Paramedic education precedes clinical practice and therefore high-quality learning materials are essential to prepare student paramedics for employment.

The case studies in this book bring together a diverse range of examples that accurately represent the caseload experienced by contemporary paramedics all in one place. They use a mix of evidence-based cases and expert opinion supplied by leaders in the industry.

Case-based learning (CBL) and problem-based learning (PBL) rely on high-quality, well-written case studies that paramedic educators, students and clinicians can use to aid their understanding of out-of-hospital care. Not only is the content contemporary, but the cases are structured in a manner that reflects a systematic approach to each scene, allowing students to develop a sense of structure to the way they proceed in each case. Each chapter has a range of interactive learning activities that allow students to stop and think about what is going on, and the questions throughout the cases provide students with additional learning opportunities.



List of contributors

Joel Beake

Registered Advanced Care Paramedic
Queensland Ambulance Service
Brisbane, QLD, Australia

Curtis Northcott

Registered Advanced Care Paramedic
RMA Medical Rescue and Registered Paramedic
Mount Isa, QLD, Australia

Fenella Corrick

GP Registrar
Western Isles, Scotland, UK

David Davis

College of Paramedics, Bridgewater, UK

Georgette Eaton

Clinical Practice Development Manager
Advanced Paramedic Practitioner (Urgent Care)
London Ambulance Service, NHS Trust, London, UK

Paul Grant

Registered Paramedic
Mines Emergency Rescue and Response (MERR)
QLD, Australia

Yasaru Gunaratne

Advanced Care Paramedic
Queensland Ambulance Service, Gold Coast, QLD
Australia

Alisha Hensby

Lecturer in Paramedicine
Charles Sturt University, Bathurst, NSW, Australia

Tom Hewes

Senior Lecturer and Programme Director of Paramedic
Sciences, Swansea University, Wales, UK

Mark Hobson

Clinical Practice Educator for Specialist Practice,
Paramedic
South Central Ambulance Service NHS Foundation
Trust, Bicester, UK

Tania Johnston

Lecturer in Paramedicine
Charles Sturt University, Bathurst, NSW, Australia

David Krygger

Advanced Care Paramedic with Specialist Training in
Low Acuity Referral Services
Queensland Ambulance Service
Gold Coast, QLD, Australia

Erica Ley

Senior HEMS Paramedic
Lincolnshire and Nottinghamshire Air Ambulance
Lincoln, UK
Associate Lecturer in Paramedic Science
University of East Anglia, Norwich, UK

Tom E. Mallinson

Rural GP & Paramedic
BASICS Scotland, Auchterarder, UK

Kristina Maximous

Lecturer in Paramedicine
Charles Sturt University, Bathurst, NSW, Australia

Brian Mfula

Lecturer in Mental Health Nursing
Swansea University, Wales, UK

Georgina Pickering

Lecturer in Paramedicine
Charles Sturt University, Bathurst, NSW, Australia

Michael Porter

Paramedic
Queensland Ambulance Service
Brisbane, QLD, Australia

Samantha Sheridan

Lecturer in Paramedicine
Charles Sturt University, Bathurst
NSW, Australia

Jennifer Stirling

Lecturer in Paramedicine
Charles Sturt University, Bathurst, NSW, Australia

Clare Sutton

Lecturer in Paramedicine, Discipline Group Leader
Charles Sturt University, Bathurst, NSW, Australia

Sam Taylor

HEMS Paramedic
Air Ambulance Kent Surrey Sussex, Chatham, UK

Ruth Townsend

Senior Lecturer in Paramedicine
Charles Sturt University, Bathurst, NSW, Australia

Lynne Walsh

Senior Lecturer in Mental Health/Public Health
Swansea University, Wales, UK

Steve Whitfield

Lecturer/Course Convenor
Griffith University, School of Medicine
(Paramedicine), Gold Coast, QLD, Australia
Registered Paramedic
Queensland Ambulance Service
Brisbane, QLD, Australia

Kerryn Wratt

Registered Paramedic
President, Australasian Wilderness and Expedition
Medicine Society (AWEMS), Omeo, VIC, Australia
CEO, RescueMED, Omeo, VIC, Australia

Aimee Yarrington

Registered Paramedic and Registered Midwife
Shropshire, UK

Chapter 1

Respiratory emergencies

Jennifer Stirling, Clare Sutton and Georgina Pickering
Charles Sturt University, Bathurst, NSW, Australia

CHAPTER CONTENTS

- Level 1:** Asthma
- Level 1:** Chronic obstructive pulmonary disease (COPD)
- Level 2:** Pulmonary embolism (PE)
- Level 2:** Life-threatening asthma
- Level 3:** Respiratory sepsis
- Level 3:** Smoke inhalation

LEVEL 1 CASE STUDY

Asthma

Information type	Data
Time of origin	17:08
Time of dispatch	17:10
On-scene time	17:20
Day of the week	Friday
Nearest hospital	30 minutes
Nearest backup	15 minutes
Patient details	Name: Betsy Booper DOB: 10/09/2002

CASE

You have been called to an outdoor running track for an 18-year-old female with shortness of breath. The caller states she has taken her inhaler to no effect.

Pre-arrival information

The patient is conscious and breathing. You can access the area via the back gate of the sports field and drive right up to the patient, who is sat down on the track.

Windscreen report

The location appears safe. Approx. 10 people around the patient. Environment – warm summer evening and good light.

Entering the location

The sports coach greets you as you get out of the ambulance and informs you that the patient suffers with exercise-induced asthma, but this is worse than normal and her inhaler has been ineffective.

On arrival with the patient

The patient is sat on a bench on the side of the track. She is leaning forward, resting her elbows on her thighs (tripodding). She says hello as you introduce yourself to her.

Patient assessment triangle**General appearance**

Alert. Speaking in short sentences. She looks panicked.

Circulation to the skin

Flushed cheeks.

Work of breathing

Breathing appears rapid and shallow. An audible wheeze is noted.

SYSTEMATIC APPROACH**Danger**

None at this time.

Response

Alert on the AVPU scale.

Airway

Clear.

Breathing

RR: 28. Regular and shallow. No accessory muscle use. Expiratory wheeze on auscultation.

Circulation

HR: 100. Regular and strong. Capillary refill time <2 seconds. Flushed cheeks and peripherally warm.

Disability

Moving all four limbs.

Pupils equal and reactive to light (PEARL).

Exposure

Bystanders have left. Next of kin are now on scene.

Temperature: warm summer evening – approx. 20 °C.

Vital signs

RR: 28 bpm

HR: 100 bpm

BP: 125/74 mmHg

SpO₂: 93%

Blood glucose: 5.2 mmol/L

Temperature: 36.9 °C

PEF: 300 L/min

GCS: 15/15

4 Lead ECG: sinus tachycardia

TASK

Look through the information provided in this case study and highlight all of the information that might concern you as a paramedic.

Aside from auscultation, which you have already done, what examination techniques should you incorporate into this patient assessment?

- Inspection – observe the chest for abnormalities such as wounds, scars, bruising, asymmetry and recession.
- Palpation – feel for any asymmetry, vocal fremitus and tenderness.
- Percussion – hyper- or hypo-resonance.

What adventitious (added) sounds might indicate asthma and why?

Expiratory wheeze. This sound is made when air has a restricted path through the bronchi, due to inflammation and muscle spasm in the airways.

What medicine (pharmacology) is likely to relieve the patient's symptoms and why?

Nebulised salbutamol – it is a Beta2, adrenergic agonist that relaxes smooth muscle in the bronchi.

Case Progression

You treat the patient with 5 mg of nebulised salbutamol and 6 L of oxygen. The nebuliser finishes and you remove the mask.

Patient assessment triangle

General appearance

The patient is now speaking in full sentences.

Circulation to the skin

Flushed.

Work of breathing

Normal effort of breathing.

SYSTEMATIC APPROACH

Danger

None at this time.

Response

Alert.

Airway

Clear.

Breathing

RR: 16. Regular. Normal depth. No accessory muscle use. No wheeze or adventitious sounds.

Circulation

HR: 105. Regular and strong. Capillary refill time <2 seconds. Flushed cheeks and peripherally warm.

Disability

No change.

Exposure

No change.

Vital signs

RR: 16 bpm

HR: 105 bpm

BP: 128/78 mmHg

SpO₂: 97%

Blood glucose: not repeated

Temperature: not repeated

PEF: 380 L/min

GCS: 15/15

4 lead ECG: sinus tachycardia

What kinds of questions would you ask this patient specifically related to asthma as part of the history-taking process?

See Table 1.1.

Table 1.1 History-taking questions**Asthma history**

Does this feel like your normal asthma?

Is this the worst it's ever been?

What time did this episode start today?

Do you take your asthma medication regularly?

What were you doing when it started today?

What usually triggers your symptoms?

When was the last time you visited your GP and/or went to hospital with these symptoms?

Have you ever been intubated or been in ICU with these symptoms?

Medication history

What asthma medications do you take?

How frequently do you have to take your medication?

Do you usually have to take your inhaler while exercising?

When was the last time you had a medication review with your GP?

Have you had any recent changes in medication?

Do you take any other medications?

Have you had any coaching on the best way to take your inhaler?

F/SH (family and social history)

Does anyone else in your family experience asthma?

Do you smoke? If so, how frequently?

Do you drink or take any drugs recreationally?

Who do you live with?

What do you do for work?

Do you exercise regularly?

Are you under any particular stress at the moment?

Past medical history (PMH)

Do you have any other medical problems?

Do you have any allergies?

Have you had a cough or cold recently?

The patient is 160 cm tall, what should her predicted peak expiratory flow reading (PEFR) be? Her first reading was 300 – what percentage is that from predicted?

(Hint: you will be required to look this up using the Australian National Asthma Council chart found here: http://www.peakflow.com/pefr_normal_values.pdf or by doing an internet search.)

- 400 L/min.
- 75%.

LEVEL 1 CASE STUDY

Chronic obstructive pulmonary disease (COPD)

Information type	Data
Time of origin	07:09
Time of dispatch	07:12
On-scene time	07:30
Day of the week	Wednesday
Nearest hospital	15 minutes
Nearest backup	40 minutes
Patient details	Name: Dave Beater DOB: 21/09/1954

CASE

You have been called to a residential address for a 66-year-old male with difficulty in breathing. The caller states he has been breathless all night and has had a cough recently. He has seen his GP who prescribed antibiotics and steroids but he feels his breathing has got worse overnight.

Pre-arrival information

The patient is conscious and breathing and is in a first-floor flat/unit.

Windscreen report

The location appears safe. Greeted at the main door by the patient's wife.

Entering the location

Wife escorts you up in the lift to the patient's flat.

On arrival with the patient

Patient is sat in the tripod position and appears distressed. He makes eye contact when you arrive, but does not speak as is so short of breath. He has a productive cough that results in a string of green-looking sputum that he manages to capture in his handkerchief to show you.

Patient assessment triangle

General appearance

Alert, and makes eye contact, but is acutely distressed. Can only speak in single words and is reluctant to talk. In tripod position, coughing.

Circulation to the skin

Pink face, breathing through pursed lips.

Work of breathing

Increased work of breathing – rapid and shallow breaths with accessory muscle use.

SYSTEMATIC APPROACH**Danger**

None at this time.

Response

Alert.

Airway

Clear.

Breathing

RR: 36. Rapid and shallow, with accessory muscle use. Widespread bilateral wheeze noted on auscultation.

Circulation

HR: 110. Radial palpable – irregular. Capillary refill time 2 seconds.

Disability

Pupils equal and reactive to light (PEARL).

Exposure

The patient is in his own home.

Vital signs

RR: 36 bpm

HR: 110 bpm

BP: 150/90 mmHg

SpO₂: 86%

Blood glucose: 4.5 mmol/L

Temperature: 37.8 °C

PEF: unable to record

GCS: 15/15

4 Lead ECG: atrial fibrillation

Allergies: nil

TASK

Look through the information provided in this case study and highlight all of the information that might concern you as a paramedic.

What is COPD?

COPD is a progressive disease and is characterized by air flow obstruction that is not fully reversible. The airway obstruction results from damage to alveoli, alveolar ducts and bronchioles due to chronic inflammation.

List the features of an acute exacerbation of COPD.

- Increased dyspnoea.
- Increased sputum production.
- Increased cough.
- Upper airway symptoms, such as a cold and sore throat.
- Increased wheeze.

- Reduced exercise tolerance.
- Fluid retention.
- Increased fatigue.
- Acute confusion.
- Worsening of previously stable condition.

Case Progression

After administration of 5 mg salbutamol via nebuliser, the patient's condition improves slightly and he hands you a medical card that his 'breathing doctor' gave to him. The card states the patient is at risk of retaining CO₂ and should only be administered with 28% oxygen to achieve saturations between 88 and 92%.

Patient assessment triangle

General appearance

Alert and more interactive.

Circulation to the skin

Pink.

Work of breathing

Increased work of breathing – breathing rapid, but not as shallow as before.

SYSTEMATIC APPROACH

Danger

None at this time.

Response

Alert.

Airway

Clear.

Breathing

RR: 30. Audible wheeze on auscultation.

Circulation

HR: 120. Palpable radial. Capillary refill time 2 seconds.

Disability

Moving all four limbs.

Exposure

Normal temperature in the ambulance.

Vital signs

RR: 30 bpm

HR: 120 bpm

BP: 148/78 mmHg

SpO₂: 90%

Blood glucose: not repeated

Temperature: not repeated

GCS: 15/15

4 lead ECG: atrial fibrillation

Allergies: nil

When the nebuliser has finished, you notice that the patient's SpO₂ is dropping so you decide to keep the patient on oxygen. What percentage of oxygen would you administer to this patient and why?

28% oxygen through a nasal cannula. The patient is at risk of developing hypercapnia respiratory failure, so it is important the oxygen is titrated to maintain saturations between 88 and 92%. Research suggests that over-oxygenation increases the mortality and morbidity of COPD patients and that titration of oxygen administration can reduce mortality.

What is meant by the term hypercapnia?

- 'A condition of abnormally elevated carbon dioxide (CO₂) levels in the blood, caused by hypoventilation, lung disease, or diminished consciousness' (NAEMT, 2015, p. 92).
- 'Alveolar hypoventilation with increased alveolar carbon dioxide limits the amount of oxygen available for diffusion into the blood, leading to secondary hypoxemia' (McCance et al., 2010, p. 1269).

LEVEL 2 CASE STUDY

Pulmonary embolism (PE)

Information type	Data
Time of origin	17:55
Time of dispatch	18:01
On-scene time	18:10
Day of the week	Friday
Nearest hospital	30 minutes
Nearest backup	15 minutes
Patient details	Name: Jasmine Wallis DOB: 27/12/2000

CASE

You have been called to a car park for a 20-year-old female who is complaining of feeling dizzy and faint.

Pre-arrival information

She is conscious and breathing.

Windscreen report

The car park is behind a row of shops and is poorly lit. The patient is hard to spot at first, as she is sitting on the metal fire escape steps with her head in her hands at the back of a building. She is alone. The car park is full, which prevents you parking near to the patient.

Entering the location

You park your ambulance as near as possible and cross the car park to get to your patient.

On arrival with the patient

The patient is able to raise her head and make eye contact.

Patient assessment triangle

General appearance

The patient looks at you when you speak and is able to speak in full sentences.

Circulation to the skin

Mildly pale.

Work of breathing

Increased. The patient looks mildly short of breath.

SYSTEMATIC APPROACH**Danger**

None at this time.

Response

Alert.

Airway

Clear.

Breathing

RR: 26. Mildly increased effort, no accessory muscle use. Auscultation – clear.

Circulation

HR: 120. Tachycardic, weak and regular pulse. Capillary refill time >2 seconds.

Disability

Pupils equal and reactive to light (PEARL).

Exposure

The patient is sitting on metal fire escape stairs, in a dark, cold car park in an undesirable part of town.

Vital signs

RR: 26 bpm

HR: 120 bpm

BP: 90/60 mmHg

SpO₂: 90%

Blood glucose: 4.4 mmol/L

Temperature: 36.5 °C

ECG: sinus tachycardia

Allergies: nil

TASK

Look through the information provided in this case study and highlight all of the information that might concern you as a paramedic.

List your differential diagnoses for this patient.

- Musculoskeletal pain.
- Pericarditis.
- Hyperventilation.
- Chest infection.
- Syncope.
- Pneumothorax.

List as many predisposing factors associated with PE as you can. Which could assist you with working through your differential diagnosis and history taking?

See Table 1.2.

Table 1.2 Pulmonary embolism predisposing factors**Surgery, especially recent**

Abdominal
Pelvic
Hip or knee
Post-operative intensive care

Obstetrics

Pregnancy

Cardiac

Recent acute myocardial infarction

Limb problems

Recent lower limb fractures
Varicose veins
Lower limb problems secondary to stroke or spinal cord injury

Malignancy

Abdominal and /or pelvic, in particular advanced metastatic disease
Concurrent chemotherapy

Other

Risk increases with age
>60 years of age
Previous proven deep vein thrombosis (DVT)/PE
Immobility
Thrombotic disorder
Neurological disease with extremity paresis
Thrombophilia
Hormone replacement therapy and oral contraception
Prolonged bed rest >3 days
Other recent trauma

Source: JRCALC (2019), p. 367.

What validated assessment tool could assist you with assessing the probability of PE in this patient?

See Table 1.3.

Table 1.3 Wells' criteria for PE

Criteria	Score
Clinical signs and symptoms of DVT (leg swelling and pain with palpation of the deep veins)	3
An alternative diagnosis is PE is less likely	3
Pulse rate >100 bpm	1.5
Immobilisation or surgery in the previous 4 weeks	1.5
Previous DVT/PE	1.5
Haemoptysis	1
Malignancy (treatment ongoing or within the last 6 months or palliative)	1
Clinical probability	
High	>6 points
Moderate	2–6 points
Low	<2 points

Source: JRCALC (2019), p. 368.

Note: When using the Wells' criteria, a low probability does not rule out PE.

Case Progression

You decide to move your patient to the back of the ambulance to continue the examination in a warm and private environment. On standing, the patient complains of feeling dizzy and faint and is unable to walk even a couple of steps. You instruct your crewmate to fetch the carry chair as you can't get the stretcher close enough to the patient.

Patient assessment triangle**General appearance**

Patient feels better when lying flat.

Circulation to the skin

Normal.

Work of breathing

Increased. Patient complains of not being able to 'catch her breath'.

SYSTEMATIC APPROACH**Danger**

None at this time.

Response

Alert.

Airway

Clear.

Breathing

RR: 30.

Circulation

HR: 128. Weak radial.

Disability

Moving all four limbs.

Exposure

Normal temperature in the ambulance.

Vital signs

RR: 30 bpm

HR: 128 bpm

BP: 88/60 mmHg

SpO₂: unable to obtain

Blood glucose: not repeated

Temperature: not repeated

GCS: 15/15

12 lead ECG: sinus tachycardia with right bundle branch block (RBBB)

What is the most common ECG finding in PE? What other ECG changes are associated with PE?

The most common ECG finding in PE is sinus tachycardia. PE can cause any of the following ECG changes:

- T-wave inversion.
- New-onset atrial fibrillation.
- Right bundle branch block.
- Right axis deviation.

- S1Q3T3 (this is a specific pattern that is seen rarely in PE):
 - S waves in lead I.
 - Q waves in lead III.
 - T-wave inversion in lead III.

Explain why females taking the oral contraceptive pill are at greater risk of developing a PE.

Virchow's triad explains the three broad categories that play a part in thrombus formation:

1. Hypercoagulability.
2. Hemodynamic changes (stasis, turbulence).
3. Endothelial injury/dysfunction.

Taking contraceptive drugs that contain oestrogen can actually change the constitution of the blood, increasing plasma and other clotting factors. This causes the woman to be in a hypercoagulative state, increasing the risk of developing DVT/PE.

LEVEL 2 CASE STUDY

Life-threatening asthma

Information type	Data
Time of origin	07:13
Time of dispatch	07:15
On-scene time	07:26
Day of the week	Monday
Nearest hospital	20 minutes
Nearest backup	10 minutes
Patient details	Name: Billy Bob DOB: 01/06/1995

CASE

You have been called to a residential address for a 25-year-old male with difficulty in breathing. Caller states he has been breathless all night and has had a cough recently.

Pre-arrival information

The patient is conscious and breathing and is located in a third-floor flat/unit – there is no lift.

Windscreen report

The location appears safe and you are greeted at the communal entrance by the patient's partner.

Entering the location

The partner appears agitated and hurries you up the stairs, stating that the patient was having his breakfast and his breathlessness got a lot worse.

On arrival with the patient

The patient is sat leaning forward and appears panicked. He does not say hello when you introduce yourself and states repeatedly that he cannot breathe, in short sharp breaths.

Patient assessment triangle

General appearance

Alert, but does not acknowledge your presence. Acutely distressed. Unable to speak in full sentences, leaning forward with clear dyspnoea.

Circulation to the skin

Pale and peripherally cyanosed.

Work of breathing

He has increased breathing effort and only giving 1 word answers.

SYSTEMATIC APPROACH**Danger**

None at this time.

Response

Alert.

Airway

Clear.

Breathing

RR: 32. Rapid and shallow. No accessory muscle use. Minimal air movement bilaterally on auscultation.

Circulation

HR: 130. Radial weak and barely palpable, regular. Capillary refill time 3 seconds. Nail beds appear bluish.

Disability

Pupils equal and reactive to light (PEARL) – 5 mm.

Exposure

The chest is exposed to conduct an assessment. The patient is in a private residence and the unit has a warm temperature.

Vital signs

RR: 32 bpm

HR: 130 bpm

BP: 100/54 mmHg

SpO₂: 87%

Blood glucose: 4.3 mmol/L

Temperature: 37.2 °C

Peak expiratory flow reading (PEFR): unable to record

GCS: E4, Verbal – not complying with your questioning, only stating he cannot breathe, M6

4 Lead ECG: sinus tachycardia, regular

Allergies: nil

TASK

Look through the information provided in this case study and highlight all of the information that might concern you as a paramedic.

Using the latest guidelines from the Australia and New Zealand Thoracic Society (ANZTS), the British Thoracic Society (BTS) or a source that draws on these resources, compare and contrast the differences between life-threatening asthma and anaphylaxis, and explain why this is more likely to be asthma than any other differential diagnosis.

Similarities: asthma and anaphylaxis both present with respiratory distress and a wheeze. Both are due to an inflammatory response. And both may appear flushed – from exertion in asthma, and in anaphylaxis the skin's reaction to the allergen.

Differences: in anaphylaxis the whole airway can be affected, producing particular symptoms not associated with asthma, such as voice changes, stridor, inspiratory wheeze and tongue and lip swelling. Also asthma is predominantly a respiratory problem, whereas anaphylaxis can present with gastrointestinal problems and hypotension, which can lead to distributive shock.

Although this did occur after eating, the patient seems to be presenting with symptoms limited purely to the respiratory system. There are no dermatological, gastrointestinal or cardiovascular changes that would indicate anaphylaxis.

Is this patient suffering from moderate, severe or life-threatening asthma, and why?

Life-threatening asthma. See Table 1.4.

Table 1.4 Comparison of asthma severity

Near-fatal asthma	Raised PaCO ₂ and/or requiring mechanical ventilation with raised inflation pressures
Life-threatening asthma	In a patient with severe asthma any one of: PEF <33% best or predicted SpO ₂ <92% PaO ₂ <8 Kpa 'Normal' PaCO ₂ (4.6–6.0 Kpa) Altered conscious level Exhaustion Arrhythmia Hypotension Cyanosis Silent chest Poor respiratory effort
Acute severe asthma	Any one of: PEF 33–50% best or predicted Respiratory rate ≥25/min Heart rate ≥110/min Inability to complete sentences in one breath
Moderate acute asthma	Increasing symptoms PEF >50–75% best or predicted No features of acute severe asthma

Source: British Thoracic Society (2019).

List your treatment, route and dosages.

- Adrenaline – 500 µg IM.
- Salbutamol – 5 mg nebulised.
- Ipratropium bromide – 500 µg nebulised.
- Oxygen – 6/8 L.
- Hydrocortisone – 100 mg IV (IM possible if unable to gain IV access).

Case Progression

You treat this patient rapidly with 500 µg of intramuscular (IM) adrenaline while your crewmate administers 5 mg of salbutamol and ipratropium bromide via a nebulizer, on 6 L of oxygen. After nebuliser therapy and 1 dose of IM Adrenaline, you rapidly extricate your patient to the ambulance. You deliver a pre-alert to the nearest emergency department.

Patient assessment triangle**General appearance**

Alert and now looking at you and nodding or shaking his head in response to your questions.

Circulation to the skin

Pale.

Work of breathing

Increased work of breathing – breathing still rapid, but less shallow.

SYSTEMATIC APPROACH**Danger**

None at this time.

Response

Alert.

Airway

Clear and peripherally cyanosed.

Breathing

RR:28. Audible wheeze on auscultation.

Circulation

HR: 128. Palpable radial. Capillary refill time 2 seconds. Nail beds appear bluish.

Disability

Moving all four limbs.

Exposure

Normal temperature in the ambulance.

Vital signs

RR: 28 bpm

HR: 128 bpm

BP: 110/78 mmHg

SpO₂: 91%

Blood glucose: not repeated

Temperature: not repeated

GCS: 15/15

4 lead ECG: sinus tachycardia

This type of incident may lead to high levels of stress during the time you are with the patient. Name at least four short-term effects of stress.

-
- Increased heart rate
 - Increased blood pressure
 - Pupil dilation
 - Sweating
 - Increased blood sugar levels
 - Inhibitions of digestive secretions
 - Peripheral vasoconstriction
 - Bronchodilation
-

Source: ANZ (2015).

It is important to recognise symptoms of long-term (chronic) stress in yourself or others. Name at least two long-term effects of stress.

-
- Behaviour changes:
 - Difficulty sleeping.
 - Altered eating habits.
 - Smoking/drinking more.
 - Avoiding friends and family.
 - Sexual problems.
 - Physical responses:
 - Tiredness.
 - Indigestion and nausea.
 - Headaches.
 - Aching muscles.
 - Palpitations.
 - Mental responses:
 - Increased indecision.
 - Difficulty concentrating.
 - Poor memory.
 - Feeling inadequate.
 - Low self-esteem.
 - Emotional responses:
 - Mood swings, becoming irritable or angry.
 - Increased anxiety.
 - Feeling numb.
 - Hypersensitivity.
 - Feeling drained and listless.
-

Source: Ambulance care practice (2019).

LEVEL 3 CASE STUDY

Respiratory sepsis

Information type	Data
Time of origin	09:15
Time of dispatch	09:30
On-scene time	09:43
Day of the week	Sunday
Nearest hospital	20 minutes
Nearest backup	40 minutes
Patient details	Name: Nicholas Beaumont DOB: 01/01/1947

CASE

You have been called to a residential address for a 73-year-old male complaining of weakness and shortness of breath.

Pre-arrival information

Patient is conscious and breathing. Upstairs in bed.

Windscreen report

The scene is safe. You are met at the door by the patient's wife.

Entering the location

The wife tells you her husband has had a productive cough for 3 days and is now unable to get out of bed.

On arrival with the patient

The patient is lying in bed and appears lethargic.

Patient assessment triangle*General appearance*

Alert but lethargic.

Circulation to the skin

Flushed, warm to touch and clammy.

Work of breathing

Increased work of breathing.

SYSTEMATIC APPROACH**Danger**

None at this time.

Response

Alert on the AVPU scale.

Airway

Clear.

Breathing

RR: 34. Rapid. Mild accessory muscle use. Right basal crackles on auscultation.

Circulation

HR: 130. Radial palpable but weak – regular. Capillary refill time 3 seconds.

Disability

Pupils equal and reactive to light (PEARL).

Exposure

The patient is in his own bed and the ambient temperature is warm.

Vital signs

RR: 34 bpm

HR: 130 bpm

BP: 108/54 mmHg

SpO₂: 87%

Blood glucose: 8.3 mmol/L

Temperature: 38.4 °C

GCS: 15/15

4 lead ECG: sinus tachycardia

Allergies: nil

TASK

Look through the information provided in this case study and highlight all of the information that might concern you as a paramedic.

Case Progression

You administer high-flow oxygen titrated to maintain an SpO₂ of 94–98%. Fluid is not indicated at this time and your local guidelines do not allow for the administration of prehospital antibiotics. You commence rapid transport to the Emergency Department with a sepsis pre-alert.

Patient assessment triangle*General appearance*

Alert.

Circulation to the skin

Flushed.

Work of breathing

Increased work of breathing.

SYSTEMATIC APPROACH**Danger**

None at this time.

Response

Alert.

Airway

Clear.

Breathing

RR: 30. Right basal crackles.

Circulation

HR: 126. Weak radial. Capillary refill time 3 seconds.

Disability

Moving all four limbs.

Exposure

Normal temperature in the ambulance. Patient covered with sheet not blanket to assist with ambient cooling.

Vital signs

RR: 30 bpm

HR: 126 bpm

BP: 100/58 mmHg

SpO₂: 90%

Blood glucose: not repeated

Temperature: 38.3 °C

GCS: 15/15

4 lead ECG: sinus tachycardia

What is sepsis?

'Sepsis is characterised by a life-threatening organ dysfunction due to a dysregulated host response to infection' (UKST, 2019, p. 14).

Outline the pathophysiology of sepsis.

Sepsis is when the body's natural inflammatory immune response to a localised infection becomes systemic, setting off a chain of physiological responses that quickly become life-threatening. It is an exaggerated response involving both the complement system (immune response) and the coagulation cascades.

The body tries to keep up with the increased demand for oxygen by raising the respiratory rate (RR) to increase the level of oxygen in the blood and to oxygenate the extra blood flow through the lungs. The heart rate (HR) and stroke volume (SV) are raised, leading to an increased cardiac output (CO). Vasodilation occurs, allowing the blood vessels to transport a greater blood volume, which eventually leads to reduced preload and reduced SV. The HR increases further to compensate, resulting in tachycardia. Some patients may be on medications that mask tachycardia (e.g. betablockers).

Profound vasodilation leads to a 'relative loss' in circulating volume and the increased permeability of the blood vessels following the release of histamine results in an 'absolute loss' as fluid escapes into the extravascular space. 25% of patients with septic shock present with a normal BP (cryptic shock or occult hypoperfusion) and others may present with relative hypotension (systolic BP >40 mmHg lower than normal systolic BP).

In the early stages patients may be warm and flushed as vasodilation leads to an increased blood volume in the peripheries. Heat generated soon becomes lost through the skin, reducing the temperature. In the later stages, the patient begins to peripherally shut down as the body attempts to redirect the blood to its core organs, which results in a further cooling of the skin. Hypothermia/cold sepsis occurs in 10–20% of patients and is more common in elderly patients. The mortality rate for these patients is double that of those with pyrexia.

Which groups are most at risk of developing sepsis?

- Elderly patients (>75 years or frail).
- Young patients (under 1 year).
- Immunocompromised patients whose immune system is impaired by medication or illness (e.g. chemotherapy patients) or where immune function is impaired due to medical conditions (diabetes and sickle cell) or medications (immunosuppressants or steroids).
- Post-surgery (within the last 6 weeks).
- Open wounds.
- Patients with indwelling medical devices (catheters or cannulas).
- Intravenous drug users.
- Pregnant women with recent history of miscarriage or termination and post-delivery.

What prompts or tools are used to determine when to screen for sepsis?

Guidelines used to recommend use of the modified systemic inflammatory response syndrome (SIRS) criteria, whereby patients presenting with two of more SIRS criteria with a confirmed or suspected infection were deemed to require further investigation to confirm or exclude a diagnosis of sepsis. This screening tool captured those patients presenting with 'uncomplicated' sepsis who were otherwise well and were at low risk for clinical deterioration. The definition of sepsis has now been updated so only those with a degree of organ dysfunction or clinical compromise are included. The SIRS criteria are no longer used as a screening tool.

The red flag system was developed to be used in conjunction with the SIRS criteria as a guide to which patients needed early intervention. This was to ensure responsible antibiotic stewardship due to the sensitivity of the SIRS criteria. The red flag system is quick to apply and is used by over 90% of UK hospitals.

The revised version of the National Early Warning Score (NEWS2) track and trigger system has been shown to be the most effective screening tool for predicting adverse outcomes for patients presenting with sepsis. This has

now been incorporated into many systems, where screening is recommended for those with a NEWS2 of greater than 5 with identified risk factors or clinician concerns.

Which components of the Sepsis Six apply to the prehospital environment?

- Oxygen: titrate to maintain SpO₂ at 94–98%.
- Fluids: bolus of 500 mL over 15 minutes if indicated (systolic BP <90 mmHg).
- Antibiotics: benzylpenicillin for meningococcal septicaemia. Refer to local guidelines regarding the use of broad-spectrum antibiotics. Not routinely recommended.
- Lactate: measure lactate if indicated by local guidelines. Not routinely recommended.

LEVEL 3 CASE STUDY

Smoke inhalation

Information type	Data
Time of origin	02:24
Time of dispatch	02:25
On-scene time	02:30
Day of the week	Friday
Nearest hospital	15 minutes
Nearest backup	10 minutes
Patient details	Name: Sam Bryant DOB: 09/09/1990

CASE

You have been called to a fire at a residential address for a 30-year-old male with smoke inhalation.

Pre-arrival information

The patient is conscious and breathing and has extricated himself from the fire. He is at the neighbour's house when you arrive.

Windscreen report

Fire and police units are on scene. The incident has been contained.

Entering the location

The patient is sat on the couch at a neighbour's house.

On arrival with the patient

The patient is talking to a police officer and appears distressed.

Patient assessment triangle

General appearance

He is alert and has soot around his mouth and nose. He is coughing quite badly.

Circulation to the skin

Normal skin colour.

Work of breathing

Increased work of breathing.

SYSTEMATIC APPROACH

Danger

None at this time – the hazard has been contained.