

# Principles of Orthopedic Practice for Primary Care Providers

Andrew J. Schoenfeld  
Cheri A. Blauwet  
Jeffrey N. Katz  
*Editors*

*Second Edition*

 Springer

# Principles of Orthopedic Practice for Primary Care Providers

Andrew J. Schoenfeld  
Cheri A. Blauwet • Jeffrey N. Katz  
Editors

# Principles of Orthopedic Practice for Primary Care Providers

Second Edition

 Springer

*Editors*

Andrew J. Schoenfeld  
Department of Orthopedic Surgery  
Brigham and Women's Hospital  
Harvard Medical School  
Boston, MA  
USA

Cheri A. Blauwet  
Department of Physical Medicine &  
Rehabilitation and Orthopedic Surgery  
Brigham and Women's Hospital  
Harvard Medical School  
Boston, MA  
USA

Jeffrey N. Katz  
Department of Orthopedic Surgery  
Brigham and Women's Hospital  
Harvard Medical School  
Boston, MA  
USA

ISBN 978-3-030-74624-7

ISBN 978-3-030-74625-4 (eBook)

<https://doi.org/10.1007/978-3-030-74625-4>

© The Editor(s) (if applicable) and The Author(s), under exclusive license to Springer Nature Switzerland AG 2021

This work is subject to copyright. All rights are solely and exclusively licensed by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG  
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

# Preface

It is a rare privilege to be able to engage in the creative process of generating a second edition of a text, as we have tried to do here with *Principles of Orthopaedic Practice for Primary Care Providers*. The original title was devised as an homage to the original *Principles of Practice* developed by Sir William Osler and which still today serves as a fundamental cornerstone in general medical care. Our hope was that we could provide similar guidance in the arena of orthopaedics and musculoskeletal medicine and the invitation to compose a second edition may signal that we were able to achieve this goal, even to some small degree. In this second edition, the editors and authorship team have sought to provide a balance of new and classic information relevant to decision-making in the orthopaedic realm, from spinal disorders to sports medicine, joint replacement, and the management of patients with osteoporosis. The second edition also introduces new chapters on pain management, adult spinal deformity, stress and running injuries, and the management of costs that we feel will be of value to primary care clinicians. We hope that the presentation of old and new in this second edition will meet the needs of primary care providers and help to inform the care of patients with orthopaedic and musculoskeletal conditions.

Boston, MA, USA

Andrew J. Schoenfeld  
Cheri A. Blauwet  
Jeffrey N. Katz

# Acknowledgments

I recognize that there is insufficient space and time to thank all of those who have contributed to, and aided me in, my life's journey and work—of which this text represents but a small part. That said, I want to thank my wife Erin for her ever-present love, dedication, and support—as well as the innumerable sacrifices made as we have pursued this career in academic medicine together. I also want to thank my children: Roman, Alyssa, and Leo, who continue to inspire, entertain, and drive me to do better. All of my accomplishments in life stand in the light and grace of God, who I know has given me more than I can ever hope to repay. I also want to thank the patient faculty and members of the Department of Orthopaedic Surgery at Brigham and Women's, many of whom graciously contributed their time and expertise to the composition of this text. I also could not have asked for a better and more energizing team of co-editors as Jeff and Cheri. As always, I further acknowledge the inspiration and support provided in words and ways seen and unseen by Patricia and David Schoenfeld; Lena and Abraham Schoenfeld; Eneida and Carlos Weber; Laura Ortiz Weber; the Andrew who is always with me; Papito, Madrina, and Madama Isabel Monsanto; Doña Juana Mercader; Josefa de Soler; Amelia Mercader; Judah Loew; Israel Baal Shem Tov; and Nachman, the son of Feiga. I dedicate my portion of this second edition to all of you.

Andrew J. Schoenfeld, MD, MSc

Dedicated to Eli, Stella, and Spencer who bring light to my day and remind me to live with gratitude.

Cheri A. Blauwet, MD

Dedicated to the memory of Dr. Robert Sheon, a brilliant, humble, inspiring pioneer in the evaluation and management of musculoskeletal problems.

Jeffrey N. Katz, MD, MSc

# Contents

## Part I The Spine

- 1 Axial Neck and Back Pain** ..... 3  
Jay M. Zampini
- 2 Sacroiliac Joint Dysfunction and Piriformis Syndrome** ..... 17  
Erika T. Yih and Danielle L. Sarno
- 3 Pain Management for Chronic Musculoskeletal Disorders** ..... 39  
Alexander J. Kim, Tonnison Malcolm, and Ehren R. Nelson
- 4 Adult Spinal Deformity** ..... 65  
Caleb M. Yeung, Harry M. Lightsey IV, and Melvin C. Makhni
- 5 Cervical Radiculopathy and Myelopathy** ..... 81  
Kevin M. Hwang, Amandeep Bhalla, and James D. Kang
- 6 Lumbar Disc Herniation and Radiculopathy** ..... 95  
Christopher M. Bono and Andrew K. Simpson
- 7 Degenerative Lumbar Spinal Stenosis and Spondylolisthesis** ..... 107  
Daniel G. Tobert and Mitchel B. Harris

## Part II Osteoporosis

- 8 Osteoporosis, Vertebral Compression Fractures,  
and Vertebral Cement Augmentation** ..... 123  
Marco L. Ferrone and Andrew J. Schoenfeld

## Part III The Hip

- 9 Hip Soft Tissue Injuries** ..... 139  
Cheri A. Blauwet and David M. Robinson

<b>10</b>	<b>Femoroacetabular Impingement, Labral Tears, and Hip Arthroscopy</b> .....	159
	Matthew J. Best and Scott D. Martin	
<b>11</b>	<b>Total Hip Arthroplasty and the Treatment of Hip Osteoarthritis</b> ...	173
	Michael J. Weaver	
<b>12</b>	<b>Hip-Spine Syndrome</b> .....	189
	Christina Y. Liu, Jeffrey Lange, and Richard Wilk	
<b>Part IV The Shoulder and Elbow</b>		
<b>13</b>	<b>Shoulder Soft Tissue Injuries</b> .....	201
	Courtney K. Dawson	
<b>14</b>	<b>Shoulder Instability</b> .....	221
	Laura Lu, Marie Walcott, and Arnold B. Alqueza	
<b>15</b>	<b>Glenohumeral Osteoarthritis</b> .....	237
	Daniel Plessl, Laurence Higgins, Michael Messina, and Carolyn M. Hettrich	
<b>16</b>	<b>Elbow Osteoarthritis and Soft Tissue Injuries</b> .....	251
	George S. M. Dyer and Stella J. Lee	
<b>Part V The Hand and Wrist</b>		
<b>17</b>	<b>Hand and Wrist Soft Tissue Conditions</b> .....	279
	Christina Y. Liu and Brandon E. Earp	
<b>18</b>	<b>Hand and Wrist Arthritis</b> .....	301
	Dafang Zhang and Barry P. Simmons	
<b>19</b>	<b>Upper Extremity Nerve Entrapment</b> .....	323
	Philip E. Blazar and Ariana N. Mora	
<b>Part VI Sports and Activity Related Injuries and Orthobiologics</b>		
<b>20</b>	<b>Bone Stress Injuries</b> .....	339
	Erin E. Finn and Adam S. Tenforde	
<b>21</b>	<b>Hamstring and Calf Injuries</b> .....	351
	David M. Robinson and Kelly C. McInnis	
<b>22</b>	<b>Running Injuries</b> .....	369
	Matthew Zinner and Rebecca G. Breslow	
<b>23</b>	<b>Orthobiologics</b> .....	395
	Kristian von Rickenbach, Alp Yurter, and Joanne Borg-Stein	



**Part VII The Knee**

**24 Knee Osteoarthritis** ..... 413  
 Jeffrey N. Katz, Kaetlyn R. Arant, and Thomas S. Thornhill

**25 Surgical Approaches to Advanced Knee OA (TKA, UKA, Osteotomy)** ..... 425  
 Adam S. Olsen and Vivek M. Shah

**26 Cartilage Defects in the Knee: Clinical, Imaging, and Treatment Aspects** ..... 437  
 Chilán B. G. Leite, Gergo Merkely, and Christian Lattermann

**27 Meniscal and Ligamentous Injuries of the Knee** ..... 453  
 Simon Goertz, Emily M. Brook, and Elizabeth G. Matzkin

**28 Anterior Knee Pain: Diagnosis and Treatment** ..... 473  
 Natalie A. Lowenstein and Elizabeth G. Matzkin

**Part VIII The Foot and Ankle**

**29 Ankle Arthritis** ..... 491  
 Eric M. Bluman, Jeremy T. Smith, Christopher P. Chiodo, and Elizabeth A. Martin

**30 Soft Tissue Disorders of the Ankle** ..... 501  
 Jeremy T. Smith, Eric M. Bluman, Christopher P. Chiodo, and Elizabeth A. Martin

**31 Midfoot Arthritis and Disorders of the Hallux** ..... 525  
 Christopher P. Chiodo, Jeremy T. Smith, Elizabeth A. Martin, and Eric M. Bluman

**32 Plantar Fasciitis** ..... 535  
 James P. Ioli

**33 Foot and Ankle Injuries** ..... 543  
 Elizabeth A. Martin, Eric M. Bluman, Christopher P. Chiodo, and Jeremy T. Smith

**Part IX Cost and Quality in Musculoskeletal Care**

**34 Managing Cost and Quality in Musculoskeletal Care** ..... 569  
 Cameron R. Egan, Adam E. Roy, and Richard Iorio

**Index** ..... 581

# Contributors

**Arnold B. Alqueza, MD** Department of Orthopedics, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Kaetlyn R. Arant, BA** Department of Orthopedic Surgery, Brigham and Women's Hospital, Boston, MA, USA

**Matthew J. Best, MD** Department of Orthopaedic Surgery, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

**Amandeep Bhalla, MD** Department of Orthopaedic Surgery, Harbor-UCLA Medical Center, Torrance, CA, USA

**Cheri A. Blauwet, MD** Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital/Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Philip E. Blazar, MD** Department of Orthopedic Surgery, Hand and Upper Extremity, Brigham and Women's Hospital, Boston, MA, USA

**Eric M. Bluman, MD, PhD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Christopher M. Bono, MD** Harvard Medical School, Boston, MA, USA  
Department of Orthopaedic Surgery, Massachusetts General Hospital, Boston, MA, USA

**Joanne Borg-Stein, MD** Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital/Harvard Medical School, Charlestown, MA, USA  
Spaulding/Newton-Wellesley Spine Center, Wellesley, MA, USA

**Rebecca G. Breslow, MD** Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Emily M. Brook, PA-C** Department of Surgery, Dartmouth Geisel School of Medicine, Hanover, NH, USA

**Christopher P. Chiodo, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Courtney K. Dawson, MD** Sports Medicine and Orthopedic Surgery, Brigham and Women's Hospital, Boston, MA, USA

**George S. M. Dyer, MD** Department of Orthopedic Surgery, Brigham and Women's Hospital, Boston, MA, USA

**Brandon E. Earp, MD** Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Cameron R. Egan, MD** Department of Orthopaedics, Massachusetts General Hospital, Boston, MA, USA

**Marco L. Ferrone, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Erin E. Finn, MD** University of Michigan Medical School, Ann Arbor, MI, USA

**Simon Goertz, MD** Brigham and Women's Hospital, Boston, MA, USA

**Mitchel B. Harris, MD** Department of Orthopaedic Surgery, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

**Carolyn M. Hettrich, MD** Sports Medicine and Shoulder Service, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Laurence Higgins, MD, MBA** Vice President of Strategic Development, Arthrex, Naples, FL, USA

**Kevin M. Hwang, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**James P. Ioli, DPM** Department of Orthopedics, Brigham and Women's Hospital, Boston, MA, USA

**Richard Iorio, MD** Department of Orthopaedics, Brigham and Women's Hospital, Boston, MA, USA

**James D. Kang, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Jeffrey N. Katz, MD, MSc** Department of Orthopedic Surgery and Division of Rheumatology, Immunity and Inflammation, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Alexander J. Kim, MD** Department of Anesthesiology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Jeffrey Lange, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Christian Lattermann, MD** Cartilage Repair Center, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Stella J. Lee, MD** Seacoast Orthopedics, Newburyport, MA, USA

**Chilan B. G. Leite, MD** Cartilage Repair Center, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

Instituto de Ortopedia e Traumatologia, Hospital das Clinicas, HCFMUSP, Faculdade de Medicina, Universidade de Sao Paulo, Sao Paulo, SP, Brazil

**Harry M. Lightsey IV, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Christina Y. Liu, MD** Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Natalie A. Lowenstein, BS** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Laura Lu, MD** Department of Orthopedics, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Melvin C. Makhni, MD, MBA** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Tennison Malcolm, MD** Department of Anesthesiology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Scott D. Martin, MD** Department of Orthopaedic Surgery, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

**Elizabeth A. Martin, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Elizabeth G. Matzkin, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Kelly C. McInnis, MD** Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

**Gergo Merkely, MD** Cartilage Repair Center, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Michael Messina, MD** Tower Health Medical Group, Orthopaedic Surgery and Total Joint Program at Phoenixville Hospital, Phoenixville, PA, USA

**Ariana N. Mora, MD** Department of Orthopedic Surgery, Hand and Upper Extremity, Brigham and Women's Hospital, Boston, MA, USA

**Ehren R. Nelson, MD** Department of Anesthesiology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Adam S. Olsen, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Boston, MA, USA

**Daniel Plessl, MD** Sports Medicine and Shoulder Service, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Kristian von Rickenbach, MD, MSc** Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital/Harvard Medical School, Charlestown, MA, USA

**David M. Robinson, MD** Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital/Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Adam E. Roy, MD** Department of Orthopaedics, Massachusetts General Hospital, Boston, MA, USA

**Danielle L. Sarno, MD** Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Department of Neurosurgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Andrew J. Schoenfeld, MD, MSc** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Vivek M. Shah, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Boston, MA, USA

**Barry P. Simmons, MD** Department of Orthopedic Surgery, Hand and Upper Extremity Service, Brigham and Women's Hospital, Boston, MA, USA

**Andrew K. Simpson, MD, MBA, MHS** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Jeremy T. Smith, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Adam S. Tenforde, MD** Spaulding National Running Center, Spaulding Rehabilitation Hospital, Cambridge, MA, USA

**Thomas S. Thornhill, MD** Department of Orthopedic Surgery, Brigham and Women's Hospital, Boston, MA, USA

Harvard Medical School, Boston, MA, USA

**Daniel G. Tobert, MD** Department of Orthopaedic Surgery, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

**Marie Walcott, MD** Agility Orthopedics, Stoneham, MA, USA

**Michael J. Weaver, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Richard Wilk, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Caleb M. Yeung, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Erika T. Yih, MD** Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Alp Yurter, MD** Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital/Harvard Medical School, Charlestown, MA, USA

**Jay M. Zampini, MD** Department of Orthopaedic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

**Dafang Zhang, MD** Department of Orthopedic Surgery, Hand and Upper Extremity Service, Brigham and Women's Hospital, Boston, MA, USA

**Matthew Zinner, BA** Charles E. Schmidt College of Medicine, Boca Raton, FL, USA

**Part I**  
**The Spine**

# Chapter 1

## Axial Neck and Back Pain



Jay M. Zampini

### Introduction

Greater than 80% of all adults will, at one time or another, experience back pain debilitating enough to impair activities of daily living, occupational performance, or quality of life. Although the lumbar spine is affected more frequently than the cervical or thoracic regions, pain that affects any segment of the spine can be termed “axial spinal pain” and should be distinguished from conditions with neurogenic pain, such as neurogenic claudication and radiculitis. The pathophysiology and treatment of axial spinal pain differ from that of the neurogenic conditions, though the two may be present concomitantly. This chapter will review the pathophysiology, evaluation, and treatment of axial pain in the neck and back.

### Definition and Epidemiology

Axial pain is defined as pain localized to one or more regions of the spine and/or SI joints without radiation into the lower extremities. It typically is present at all times and not necessarily aggravated by ambulation or activity. Pain may be lessened with rest or lying flat, but this does not have to be the case and is not required for a diagnosis. There are a number of factors that may be responsible for axial pain including joint dysfunction, degenerative changes, trauma, tumor or infection, myofascial structures, and non-organic pain generators.

---

J. M. Zampini (✉)

Department of Orthopaedic Surgery, Brigham and Women’s Hospital, Harvard Medical School, Boston, MA, USA

e-mail: [jzampini@bwh.harvard.edu](mailto:jzampini@bwh.harvard.edu)

© The Author(s), under exclusive license to Springer Nature Switzerland AG 2021

A. J. Schoenfeld et al. (eds.), *Principles of Orthopedic Practice for Primary Care Providers*, [https://doi.org/10.1007/978-3-030-74625-4\\_1](https://doi.org/10.1007/978-3-030-74625-4_1)



With greater than 80% of the adult population experiencing axial spinal pain at some point in life, and many not seeking medical care, it is difficult to make conclusive epidemiologic statements about populations at risk. It can almost be stated that anyone who lives long enough is at risk for back pain. Certain factors are known to associate with a higher risk of chronic axial pain, including obesity, tobacco use, total body vibration as may occur in long-distance truck driving or using a jackhammer, and repetitive hyperextension activities of the lumbar spine.

## **Clinical Presentation**


### ***Pain History***

The evaluation of axial spinal pain is no different than any other pain evaluation and should include the time of onset, location of maximal pain, duration, severity, and associated symptoms. An inciting event should be noted if possible. A patient should be asked to consider events in the 2–3 days preceding the onset of pain since the inflammation, which often causes axial spinal pain, will increase over this time period. Body positions or maneuvers that exacerbate or alleviate the pain should be sought as should other associated symptoms. Patients should also be queried as to whether similar symptoms have presented in the past.

A thorough axial pain evaluation is then performed, with consideration given to the structures that may be pain generators. All spinal structures can potentially cause pain. These structures include the vertebral body and disc in the anterior spine; facet joints, other bony processes, interspinous and supraspinous ligaments, and SI joints posteriorly; as well as the myofascial tissue in all spinal regions (Fig. 1.1). As each of these structures performs a unique function, they also possess characteristic patterns of pain that may be elucidated through the history and physical exam. The pain patterns typically associated with dysfunction of each key spinal structure are summarized in Table 1.1.

The history of axial pain should clearly document the presence or absence of any “red flag,” signs, and symptoms. A history of acute, high-energy trauma, such as car accidents or falls from greater than standing height, would suggest the need for emergent evaluation. Constitutional symptoms, such as unintended weight loss in excess of 10% of body weight or unexplained fevers or chills, would suggest the need for a neoplastic or infectious work up. Other neurologic “red flags,” such as bowel or bladder retention or incontinence, should be sought to identify potential neurologic emergencies.

**Fig. 1.1** Schematic of the human spine. The spine contains four zones: cervical, thoracic, lumbar, and sacrum



**Table 1.1** Pain patterns typically associated with dysfunction of key spinal structures

	Myofascial	Fracture	Discogenic	Facetogenic	Sacroiliac
Injury identified	No	Yes	No	No	No
Tenderness	Trigger point	Focal	No	Focal	Focal
Exacerbating factors	Muscle stretch or activation	Spinal motion	Prolonged sitting or standing	Spinal hyperextension	Forced SI joint motion
Alleviating factors	Muscle rest	Immobilization	Recumbency	Recumbency	Recumbency
Neurologic symptoms	None	Possible	Possible	Possible	None
Referred pain <sup>a</sup>	None	Possible	Possible	Possible	Possible

<sup>a</sup>Cervical spine conditions can cause referred pain between the occiput and the lower scapulae, depending on the spinal level of the condition. Lumbar conditions can cause referred pain to the buttock and posterior thighs

### ***Physical Examination***

A specific diagnosis of axial pain can be made most often by the history alone. The physical examination serves to confirm the expected diagnosis. For most patients, it is useful to examine all aspects of the spine not expected to be painful before focusing on the structure anticipated to be the pain generator, since the examination is sure to exacerbate the pain at least temporarily. Any involuntary guarding associated with increased pain can obscure other aspects of the evaluation such as the

neurologic examination. Examination of the sensory, motor, and reflex functions can often be performed first and without any additional discomfort to the patient. This should be followed by a standing examination of the spine. Spinal curvature and posture should be evaluated with attention to shoulder height, pelvic obliquity, and any deviation of spinal balance. Spinal balance generally means that the patient's head is centered over the pelvis in both the sagittal and coronal planes. Gait should be examined from this position as well; attention should be paid to voluntary and involuntary alteration of gait to avoid pain and to any assistance device required for mobility. In the standing position, the spine should be palpated in the midline to determine if any bony tenderness is present. The musculature should be palpated next, again focusing on areas not expected to be tender before palpating potentially painful muscles. Spinal motion should be assessed last as this is often most painful for the patient. Objective measurements of spinal flexion, extension, lateral bending, and rotation, while valuable to document objective responses to treatment, are typically not as helpful for diagnostic purposes.

Next, provocative maneuvers should be performed for diagnostic confirmation if necessary. For axial spinal pain, provocative maneuvers are most useful for confirming the SI joint as the source of pain. A patient should be supine for most of these tests. One sensitive test of the SI joint is performed by passively flexing the hip on the painful side and then abducting and externally rotating the hip while the contralateral leg remains on the examination table. This maneuver—flexion abduction external rotation (FABER) test—compresses the ipsilateral SI joint and reproduces pain as a result. The test is positive if pain near the SI joint is reproduced. The test is nonspecific, however, since several structures are manipulated simultaneously (the hip joint, SI joint, lumbar spine, musculature), and should be followed by other confirmatory tests. If pain at the SI joint can be reproduced by compressing the pelvis either by using bilateral, posteriorly directed pressure on the anterior superior iliac spines (ASIS) in the supine position, the AP pelvic compression test, or by pressure on the greater trochanter with the patient in the lateral decubitus position, the lateral pelvic compression test, then the painful structure can be confirmed to be the SI joint.

Provocative testing of the facet joints or palpation that reproduces pain in this area, or over myofascial structures, can also be helpful in formulating a differential diagnosis. Extension of the neck and lumbosacral region that reproduces axial pain may also indicate the facet joints as a potential source of symptoms. Pain exacerbated on forward flexion at the lumbosacral junction and also reproduced with axial loading of the shoulders may be indicative of discogenic pain. A final aspect of the physical examination includes evaluation of other potentially painful joints in the upper or lower extremities to rule out these structures as additional pain generators or contributors to the overall constellation of symptoms. It is important to realize that there may be more than one clinical entity responsible for symptoms, and there is emerging appreciation for the interplay between the spinal and pelvic structures, as well as the neck and shoulder girdle, in pain syndromes. These clinical conditions are now frequently referred to as “neck-shoulder syndrome” or “hip-spine syndrome.”

One further consideration in the examination of a patient with axial pain is the impact of psychological somatization and symptom magnification. These patients will perceive pain that is either present without any physical disruption of a spinal structure or out of proportion to what would be expected by the physical condition. To make this determination requires a nuanced approach to patient evaluation; several classic findings, termed Waddell's findings, have been reported to correlate with somatization and symptom magnification. Gentle downward compression of a patient's head does not cause any motion of the lumbar spine and should, therefore, cause no low back pain. Similarly, if spinal motion is simulated—with rotation of the shoulders, back, and pelvis at the same time—the spine itself is not affected, and no pain should be experienced. Finally, light touch of the skin overlying the spine should not produce pain. Observation of pain with any of these maneuvers should alert the clinician that non-organic factors are contributing to the patient's pain and should be taken into account when planning further evaluation and treatment.

## **Differential Diagnosis and Diagnostic Testing**

### ***Myofascial Pain***

Muscles are the structures most susceptible to fatigue and overuse injury as well as to injuries resulting from acute demand exceeding muscle capacity. These injuries collectively comprise the most common cause of spinal pain and are generically called strains. Activation or passive stretch of the injured muscle will exacerbate the pain. Palpation will reveal focal, typically unilateral tenderness at the site of muscle injury. Multiple painful triggers may be encountered in the paraspinal musculature of patients with myofascial pain syndromes, such as fibromyalgia. Imaging does not help confirm a diagnosis but does rule out other potential etiologies as a cause of pain.

### ***Pain Associated with Fractures and Ligamentous Injuries***

In both young and old patients, referred pain can be felt in a pattern characteristic of the level of injury. Injuries close to the upper cervical spine will have referred pain to the occiput; injuries of the lower cervical spine will have referred pain even as far distally as the lower aspect of the scapulae. Similarly, lumbar fracture patients can complain of referred pain to the buttocks or upper thighs. Dermatomal symptoms to the hands or feet do not represent referred pain and suggest that a full neurologic exam should be included. Palpation reveals focal tenderness at the site of injury. Plain film and computed tomography (CT) imaging are used to diagnose or confirm a fracture. Magnetic resonance imaging (MRI) may be required if these initial studies are negative to evaluate for concomitant disc or ligamentous injury or to assess the acuity of a particular fracture.


## ***Discogenic Pain***


Several painful conditions have been shown to localize to the disc: tears of the annulus, herniated discs, and degenerative disc disease (Fig. 1.2). With an annular tear, patients complain of axial pain deep inside the spine and focally at or near the injury site. Pain is typically increased with lumbar flexion or sitting and relieved with lumbar extension or lying flat. Plain film images may be read as negative depending on the extent of degenerative changes involving the disc space (Fig. 1.3). MRI is the diagnostic test of choice and will accurately display the amount of disc degeneration at various levels within the spine (Fig. 1.4). As a result, this imaging modality is nonspecific and cannot identify which, if any of the degenerative discs identified, is the cause of a patient's axial pain.

## ***Facetogenic Pain***

Patients with painful, degenerative facet joints will complain of morning pain and stiffness of the back. Spinal extension increases the load borne by the facet joints, and patients will complain that this maneuver exacerbates the pain. Referred pain is often present with painful facets: upper cervical facet referred pain may be perceived along the occiput with lower cervical referred pain felt in the shoulders or scapulae. Lumbar referred pain is perceived within the buttocks, pelvis, or posterior thighs. Spinal extension may increase the sensation of referred pain. It should be noted that the discs

**Fig. 1.2** This sagittal, T2-weighted MRI of the lumbar spine shows normal (*white arrow*) and degenerative discs. The degenerative discs show decreased disc height and low disc signal from loss of disc hydration (*white arrow head*) and annular tearing (*black arrow head*)





**Fig. 1.3** Planar radiographs of the lumbar spine are ideal to identify and monitor scoliosis (a), spondylolisthesis (b), and compression fractures (c)

and facet joints age or degenerate concomitantly and may be symptomatic simultaneously. These patients will note that prolonged sitting and standing both exacerbate pain. Plain film, CT, and MR imaging can all demonstrate evidence of facet arthrosis, although none of these imaging modalities is considered a specific test.

### ***Sacroiliac Pain***

The SI joints form the link between the spine and pelvis. The joints are extremely stable as a result of strong ligaments on both the posterior and anterior aspects of the joint. Patients with painful sacroiliac joints complain of pain just medial to the posterior superior iliac spines, the bony prominences at the top of the buttocks. Patients



**Fig. 1.4** MRI is useful for identifying the source of axial spinal pain including occult fractures (a) and ligament sprains (b). The occult fracture (a) is identified by the high STIR signal in the vertebral body (arrow) compared to low signal in an uninjured vertebra (arrow head). The ligament injury (b) is shown at the arrow compared to a normal-appearing ligamentum flavum seen at the level below (arrow head)

may experience pain with lumbosacral range of motion, ambulation, or single-leg stance. The unique location and function of the SI joints allows for a somewhat more focused examination than for other degenerative spinal conditions. At least three other provocative maneuvers (FABER test, thigh thrust, Gaenslen's test, and/or pelvic compression) should be positive to confirm SI pain with relative certainty. Plain film images and CT scans may show joint degeneration, while active inflammation or synovitis can be appreciated on MRI. The extent of findings localized to the SI joint does not necessarily correlate with the degree of a patient's SI-related pain.

### *Conditions Causing Referred Pain to the Spine*

All evaluations of axial spinal pain should consider non-spinal sources as well. Visceral, vascular, autoimmune, neoplastic, and infectious conditions are responsible for 2–3% of all axial spine pain. These conditions often cause non-mechanical pain, or pain that does not change with spinal motion. Patients will report that they “Just can't get comfortable in any position.” Red flag signs and symptoms should be sought in these patients with a concomitant vascular examination as deemed necessary.



## Nonoperative Management

A large majority of patients with newly diagnosed axial pain will return to their baseline state of spinal health within 4–6 weeks, oftentimes with little to no treatment. For this reason, noninvasive, nonoperative modalities are the preferred choice for the treatment of axial spinal pain.

For patients with acute spinal pain—whatever the underlying origin—a short period of rest from aggravating maneuvers is indicated. A patient should not be placed on complete bed rest for more than 1–2 days. After even a few days of bed rest, the musculature of the entire body, including the paraspinal muscles, will begin to atrophy, making effective rehabilitation a challenge. The patient should be advised to return to activity as soon as possible with avoidance of the most painful activities. Additionally, nonsteroidal anti-inflammatory drugs should be prescribed at an appropriate dose for the purposes of pain relief. An oral steroid taper can also be used but should be used with caution, as several reports have suggested that oral steroids may reduce the efficacy of later, more invasive treatments such as injections.

By 2–4 weeks following symptom onset, most patients will have recovered sufficiently to resume most activities of daily living and even more strenuous activities such as exercise. It is at this point that physical therapy (PT) can be helpful to further reduce pain and to begin rehabilitation and prevention of future exacerbations. Therapists can perform pain-relieving treatments including massage, stretch, and spinal manipulation to accelerate pain reduction. This phase of treatment may also include chiropractic care and acupuncture. The long-term goals of PT should focus on improving muscle strength. Patients with muscle strains require strengthening of the injured muscle and all muscles that support the spine (known as the “core” musculature) to become better able to participate in the activities that initially precipitated the pain. Even patients with annular tears, herniated discs, and degenerative conditions can benefit from the trunk stability provided by strengthening the paraspinal musculature. Using one or more of these three noninvasive treatments, greater than 90% of patients should experience relief of acute axial pain, and many should experience long-term maintenance of spinal health.

Patients who fail to achieve relief of axial spinal pain through activity modification, oral agents, and therapy often can be treated with spinal injections. Injection techniques vary and are chosen for the specific pathology to be treated. Chronic muscle strains or muscle spasm may benefit from trigger point injections at the point(s) of maximal muscle tenderness. Recalcitrant cases of muscle spasm, particularly with cervical torticollis, are sometimes treated with injection of botulinum toxin (Botox, Allergan, Dublin, Ireland).

Axial pain thought to result from the disc or facet joints can be treated with epidural and perifacet injections, respectively. Epidural injections typically involve localization of the affected spinal level on fluoroscopy followed by injection of lidocaine and a corticosteroid. Immediate reduction of the pain with the effect of the topical anesthetic agent confirms the target as a pain generator. Epidural injections are best reserved for pathology within the spinal canal—disc herniations and occasionally annular tears. Patients with facet pathology benefit from perifacet



injections. These injections can be placed directly into the facet capsule; however, most pain specialists now inject anesthetic cranially and caudally to the facet to block the medial branch of the dorsal primary ramus of the nerve root, the main innervation of the joint. These medial branch blocks have been found to be safer and more effective for reduction in pain emanating from the facets. Additionally, medial branch blocks can be used to plan radiofrequency denervation of the facet joint, a technique that offers longer-term relief of facet-based pain in well-selected patients.

Aside from pain relief, two other benefits are provided through spinal injections. First, if a patient experiences partial relief with the injection, he or she may be better able to participate in therapy. The two modalities can then work synergistically to accelerate recovery and prevent future recurrence. Second, the application of a topical anesthetic agent or corticosteroid can help to predict if a patient will respond favorably to surgery. Temporary but substantial relief of symptoms implies that a more permanent treatment option, namely, surgery, could be considered in select clinical scenarios.

## **Indications for Surgery**

Surgery is not indicated for the vast majority of patients with axial neck and back pain for several reasons: the condition is often not amenable to surgery (e.g., muscle strain, ligament sprain), the condition is stable and self-limited (e.g., most compression fractures and nearly all spinous process and transverse process fractures), or imaging findings are too diffuse to determine which process represents the main pain generator (e.g., multilevel degeneration with axial pain). Surgical treatment of axial pain is currently well indicated for patients with scoliosis and kyphosis, spondylolisthesis, and spinal instability resulting from fractures and dislocations. Surgical intervention for degenerative disease with axial pain in the absence of neurogenic symptoms is rarely indicated, and only if the degeneration is localized, patients have failed to achieve sustained pain relief with nonoperative modalities, and significant clinical information can confirm that the degenerative conditions identified are the sole pain generators. The clinical information best able to predict a positive outcome following surgery is the observation of complete (or near complete) resolution of axial pain with focal spinal injections coupled with consistent, reproducible physical examination findings pointing to the degenerative structures as pain generators. Additionally, the patient's history should be free of other psychosocial factors that could confound treatment. These factors include psychiatric conditions with predominant somatization symptoms, presence of active litigation related to an injury associated with the pain (e.g., car accidents, work-related injuries), and the presence of an active workers' compensation claim.

## **Operative Management**

One of the most compelling reasons to avoid surgery for axial pain, if at all possible, is that fusion-based procedures are the primary treatment for these conditions. The main rationale for fusion follows the logic that pain from a moving structure can be controlled by eliminating motion at the structure. In all segments of the spine and SI joints, fusion involves preparing the environment surrounding two bones to be conducive for the growth of a new bone. The bridging bone will then join the two initially independent segments into a single structure.

### ***Anterior Spinal Fusion***

Spinal fusion can be performed from an anterior approach to the disc space between the vertebral bodies. These operations are termed “interbody” or “intervertebral” fusions for this reason. The technique is most often used for anterior cervical spine surgery and in the lumbar spine for discogenic back pain. Anterior fusion enjoys the advantage of a large space for the placement of bone graft for fusion between the well-vascularized vertebral bodies. Cervical spine surgery is readily accomplished in this manner with a relatively minimally invasive approach that exploits natural anatomic planes between the trachea, esophagus, and major neurovascular structures in the neck. Thoracolumbar surgery, however, has the disadvantage of requiring exposure through the thoracic and abdominal cavities with attendant risk of injury to the visceral and vascular structures contained therein. Bone graft, either from a cadaveric donor or from the anterior iliac crest, is impacted into the space previously occupied by the intervertebral disc to achieve the fusion. This is typically stabilized using a metal plate affixed to the anterior aspect of the vertebrae with bone screws, as such instrumentation has been shown to provide more immediate stability and enhance the likelihood of fusion.

Postoperatively, patients often use a cervical collar or brace to protect the spine until pain begins to resolve. The fusion site will heal over the course of several months and is monitored using periodic radiographs. Visualization of bone bridging between the intended vertebrae signifies complete healing of the fusion.

### ***Posterior Spinal Fusion***

Thoracolumbar fusion is most commonly performed using a posterior approach. The advantage of the posterior approach in the thoracic and lumbar regions is that long segments of the spine can be accessed without violating the thoracic

and abdominal cavities and complication rates are reduced as a result. Fusion can be achieved by placing an interbody graft using carbon fiber or titanium cages, cadaver bone, or autograft from the iliac crest or elsewhere. Stabilization is achieved via bone screws anchored to the vertebrae through channels created in the pedicles and connected by rods. Patients may be given a back brace to assist mobilization after thoracolumbar posterior fusion. The brace is typically used only until a patient's pain resolves and the muscles once again become able to assist stability. In patients with osteopenia or osteoporosis, a rigid brace may be prescribed for use until the fusion site shows signs of consolidation on radiographs.

### ***SI Joint Fusion***

Fusion of the SI joint requires debridement of the cartilage of the joint with replacement of the cartilage with bone graft. The SI joint can be accessed anteriorly or posteriorly with bone graft taken directly from the ilium. Stabilization is achieved using a plate bridging from the sacrum to the ilium or via percutaneously placed screws that span the joint space.

After SI fusion, patients are instructed to use crutches or a walker to assist in mobilization. Weight bearing on the operative limb is restricted to the so-called “toe-touch” or “touchdown” weight bearing for several weeks following surgery.

### **Expected Outcomes**

The vast majority of patients (up to 90%) with acute axial pain can be expected to experience pain relief within 6 weeks of symptom onset. Patients with initial episodes of pain can, therefore, be reassured that the pain will resolve and not result in a chronic condition. In general, the longer a patient experiences activity-limiting axial pain, the longer treatment will take to relieve the pain, and the less likely he or she will be to experience complete pain relief. This observation was recently confirmed in an analysis of the multicenter Spine Patient Outcomes Research Trials (SPORT). Patients with lumbar disc herniations who experienced functional limitations for greater than 6 months were found to have inferior results, irrespective of treatment, as compared to patients in pain for less than 6 months. It is unclear if this finding suggests that patients developed chronic pain syndromes independent of the initial pain generator or if permanent structural damage to the spine was responsible.

If a patient is unable to achieve satisfactory relief through nonoperative measures, fusion-based procedures have been shown to result in long-term reductions in pain and improvement in function for only 60–70% of well-selected patients

with axial neck and back pain. Reports of randomized trials and observational studies have shown that some well-selected patients could achieve pain relief and functional improvement following surgery. The selection process must be rigorous, however, in order to assure the best outcome possible. Ideally, patients should be free from nicotine products and should not be involved in litigation over the cause of pain to assure optimal outcomes. Patients must additionally be prepared to expect that no treatment will completely eliminate back pain. They should be counseled that pain reduction will approximate what was achieved with spinal injections and should be willing to accept that a 50% reduction in pain may be the best that can be achieved. Patients expecting full alleviation of pain following surgery should have their expectations appropriately adjusted through counseling from primary care physicians and surgeons prior to agreeing to any procedure (Table 1.2).

**Table 1.2** Summary of axial neck and back pain disorders with synopsis of presentation, diagnostic testing, and suggested management options

Clinical entity	Presentation	Diagnostic testing	Conservative management	Surgical indications and operative management
Myofascial pain	Trigger point tenderness Limited or no focal pain	Primarily clinical	Rest, ice, NSAIDS PT Trigger point injection	N/A
Fracture/ligamentous injury	History of trauma Focal tenderness to palpation over injured region	Plain films/CT MRI—if there is concern for ligamentous injury	Rest, ice, NSAIDS PT Spinal bracing	Spinal instability or failure of nonoperative management with persistent pain Spinal stabilization procedures often require instrumented fusion
Discogenic back pain	Pain worse with sitting or standing Forward flexion exacerbates the pain	MRI-degenerative changes involving the discs (may not be diagnostic)	NSAIDS PT Spinal injections	Reserved for select cases where nonoperative treatment fails Fusion-based procedure
Facetogenic pain	Pain worse with standing and ambulation Extension exacerbates the pain	MRI-degenerative changes involving the facet joints (may not be diagnostic)	NSAIDS PT Facet injections, radiofrequency lesioning, rhizotomy	Reserved for select cases where nonoperative treatment fails Fusion-based procedure

*PT* physical therapy, *CT* computed tomography, *MRI* magnetic resonance imaging, *NSAIDs* non-steroidal anti-inflammatory drugs

## Suggested Reading

- Fritzell P, Hagg O, Wessberg P, Nordwall A. 2001 Volvo Award Winner in Clinical Studies: Lumbar fusion versus nonsurgical treatment for chronic low back pain: a multicenter randomized controlled trial from the Swedish Lumbar Spine Study Group. *Spine (Phila Pa 1976)*. 2001;26(23):2521–32.
- Kaye AD, Manchikanti L, Abdi S, et al. Efficacy of epidural injections in managing chronic spinal pain: a best evidence synthesis. *Pain Physician*. 2015;18(6):E939–1004.
- Khan JM, Harada GK, Basques BA, Nolte MT, Louie PK, Iloanya M, Tchalukov K, Berkowitz M, Derman P, Colman M, An HS. Patients with predominantly back pain at the time of lumbar fusion for low-grade spondylolisthesis experience similar clinical improvement to patients with predominantly leg pain: mid-term results. *Spine J*. 2020;20(2):276–82.
- Kleimeyer JP, Cheng I, Alamin TF, Hu SS, Cha T, Yanamadala V, Wood KB. Selective Anterior Lumbar Interbody Fusion for Low Back Pain Associated With Degenerative Disc Disease Versus Nonsurgical Management. *Spine (Phila Pa 1976)*. 2018;43(19):1372–80.
- Pearson AM, Lurie JD, Tosteson TD, Zhao W, Abdu WA, Weinstein JN. Who should undergo surgery for degenerative spondylolisthesis? Treatment effect predictors in SPORT. *Spine (Phila Pa 1976)*. 2013;38(21):1799–811.
- Riew KD, Ecker E, Dettori JR. Anterior cervical discectomy and fusion for the management of axial neck pain in the absence of radiculopathy or myelopathy. *Evid Based Spine Care J*. 2010;1(3):45–50.

# Chapter 2

## Sacroiliac Joint Dysfunction and Piriformis Syndrome



Erika T. Yih and Danielle L. Sarno

### Part I: Sacroiliac Joint Dysfunction

#### *SI Joint Anatomy, Innervation, and Function*

##### **Anatomy**

The sacroiliac (SI) joint is a large axial joint connecting the spine to the pelvis. Lying at the junction of the sacrum and the ilium, the average surface area of the joint is 17.5 cm<sup>2</sup>, and the average volume is 0.6–2.5 mL [1]. However, the SI joint varies widely in size, shape, and contour between individuals and sometimes even between sides within a single individual [2]. Although the SI joint is typically characterized as a large synovial joint, only the anterior third is a true synovial joint, while the rest of the junction has an absent or only rudimentary posterior capsule and is instead supported by an intricate ligamentous system [3]. The SI joint is also supported by a large network of pelvic and lower extremity muscles, some of which (e.g., gluteus maximus, piriformis, biceps femoris) are functionally connected to the SI joint ligaments and therefore affect joint mobility [4].

The surface of the SI joint is flat until puberty, when the iliac surface starts to become rougher and duller and develop some fibrous plaques [4, 5]. In the third and

---

E. T. Yih

Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA  
e-mail: [eyih@partners.org](mailto:eyih@partners.org)

D. L. Sarno (✉)

Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Department of Neurosurgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA  
e-mail: [dsarno@bwh.harvard.edu](mailto:dsarno@bwh.harvard.edu)