


Surgery of Exotic Animals

Edited by

R. Avery Bennett | Geoffrey W. Pye



WILEY Blackwell

Surgery of Exotic Animals

Surgery of Exotic Animals

Edited by

*R. Avery Bennett, DVM, MS, DACVS
Department of Small Animal Clinical Services Louisiana State University
School of Veterinary Medicine
Baton Rouge, LA, USA*

*Geoffrey W. Pye, BVSc, MSc, DACZM
Disney's Animals, Science, and Environment Team
Disney's Animal Kingdom
Bay Lake, FL, USA*

WILEY Blackwell

This edition first published 2022

© 2022 John Wiley & Sons, Inc.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by law. Advice on how to obtain permission to reuse material from this title is available at <http://www.wiley.com/go/permissions>.

The right of R. Avery Bennett and Geoffrey W. Pye to be identified as the authors of the editorial material in this work has been asserted in accordance with law.

Registered Office

John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, USA

Editorial Office

111 River Street, Hoboken, NJ 07030, USA

For details of our global editorial offices, customer services, and more information about Wiley products visit us at www.wiley.com.

Wiley also publishes its books in a variety of electronic formats and by print-on-demand. Some content that appears in standard print versions of this book may not be available in other formats.

Limit of Liability/Disclaimer of Warranty

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting scientific method, diagnosis, or treatment by physicians for any particular patient. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. While the publisher and authors have used their best efforts in preparing this work, they make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives, written sales materials or promotional statements for this work. The fact that an organization, website, or product is referred to in this work as a citation and/or potential source of further information does not mean that the publisher and authors endorse the information or services the organization, website, or product may provide or recommendations it may make. This work is sold with the understanding that the publisher is not engaged in rendering professional services. The advice and strategies contained herein may not be suitable for your situation. You should consult with a specialist where appropriate. Further, readers should be aware that websites listed in this work may have changed or disappeared between when this work was written and when it is read. Neither the publisher nor authors shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

Library of Congress Cataloging-in-Publication Data

Names: Bennett, R. Avery, editor. | Pye, Geoffrey W., editor.

Title: Surgery of exotic animals / edited by R. Avery Bennett, Geoffrey W. Pye.

Description: First edition. | Hoboken, NJ : Wiley-Blackwell, 2022. |

Includes bibliographical references and index.

Identifiers: LCCN 2021028715 (print) | LCCN 2021028716 (ebook) |

ISBN 9781119139584 (hardback) | ISBN 9781119139591 (adobe pdf) |

ISBN 9781119139607 (epub)

Subjects: MESH: Animals, Exotic | Surgery, Veterinary--methods

Classification: LCC SF997.5.E95 (print) | LCC SF997.5.E95 (ebook) | NLM

SF 997.5.E95 | DDC 636.089/7--dc23

LC record available at <https://lcn.loc.gov/2021028715>

LC ebook record available at <https://lcn.loc.gov/2021028716>

Cover Design: Wiley

Cover Image: © Kerri A. D'Ancicco, Michael Karlin, Geoffrey W. Pye, R. Avery Bennett

Set in 9.5/12.5pt STIXTwoText by Straive, Pondicherry, India

10 9 8 7 6 5 4 3 2 1

Avery

Since childhood I have been fascinated with veterinary medicine and the diversity of species. Because of various events I abandoned my dream of becoming a veterinarian and obtained a Bachelor of Science degree in Secondary Education. After being a teacher for three years, I joined the Peace Corps and served in Belize. It was while serving, I decided to try again to complete my preveterinary sciences to be eligible to apply. During Veterinary School, my interests were surgery and zoological medicine. While I applied for residencies in both, I was granted a residency in Small Animal Surgery. After graduation, I went into my Rotating Internship at Texas A&M University where I met Elizabeth A. Russo, DVM, MS, DACVIM. Liz was my first mentor in zoological medicine. She had done a residency in Zoological Medicine at the Bronx Zoo, but I remember her telling me she felt she had not learned medicine. So, she completed a residency in Small Animal Medicine and Surgery and became a Diplomate of the American College of Veterinary Internal Medicine. Her goal was to apply the principles she had learned in her training to become a board certified specialist in Internal Medicine to exotic animal species. While she passed away before I finished my training in Veterinary Surgery, I modeled my career after her, applying the principles and techniques I learned in my training becoming a boarded Veterinary Surgeon to exotic animals. I dedicate this book to Claudio Ribeiro da Silva, my husband and life partner; and to Elizabeth A Russo, DVM, MS, DACVIM for inspiring me to advance the field of surgery in exotic animals. For many years Geoff has pushed me to write this book and, in all honesty, without him this book would never have come to be. All my thanks to you, Geoff!

Geoff

There are always pivotal moments and people that change the trajectory of life. For me, I have been lucky to have both hugely influence my life and career. I dedicate this book to my parents, Cath and Bill, who instilled in me a love for nature, and to my family, Dodi, Hannah, and Zach, who instill in me a love for life and have been immensely supportive in this endeavor. I wish to recognize Drs. Jim Carpenter, Pat Morris, Don Janssen, and Scott Terrell who took leaps of faith based on what they saw I could be rather than who I was and have provided me with so many opportunities and learnings that have helped me grow my career and as a person. Finally I would like to thank Avery for being a patient and amazing mentor and encouraging my passion for surgery. "May I make a suggestion?," as I incorrectly held an instrument, chose an inappropriate suture, planned to cut in the wrong place/grab the wrong tissue, or looked to create too much surgical trauma, lives fondly in my many memories of Avery. Thank you sir.

Contents

List of Contributors ix

Preface xi

About the Editors xii

- 1 General Principles, Instruments, and Equipment** 1
R. Avery Bennett
- 2 Suture Materials** 11
Michael S. McFadden
- 3 Magnification Surgery** 23
Heidi Phillips
- 4 Invertebrate Surgery** 35
Gregory A. Lewbart
- 5 Fish Surgery** 46
Claire Vergneau-Grosset and E. Scott Weber, III
- 6 Amphibian Surgery** 63
Claire Vergneau-Grosset and E. Scott Weber, III
- 7 Reptile Orthopedic Surgery** 74
Michael S. McFadden
- 8 Surgical Approaches to the Reptile Coelom** 85
Geoffrey W. Pye and R. Avery Bennett
- 9 Reptiles: Soft Tissue Surgery** 96
Stephen J. Mehler and R. Avery Bennett
- 10 Avian Orthopedics** 112
Brett Darrow and R. Avery Bennett
- 11 Approaches to the Caudal Coelom (Abdomen) of Birds** 154
Michael B. Mison and R. Avery Bennett
- 12 Avian Reproductive Procedures** 163
Stephen J. Mehler and R. Avery Bennett

- 13 Surgery of the Avian Gastrointestinal Tract 175**
Michael B. Mison and R. Avery Bennett
- 14 Surgery of the Avian Respiratory System and Cranial Coelom 190**
Geoffrey W. Pye
- 15 Minimally Invasive Surgery Techniques in Exotic Animals 204**
Stephen J. Mehler and R. Avery Bennett
- 16 Orthopedic Surgery in Small Mammals 227**
Michael S. McFadden
- 17 Rabbit Soft Tissue Surgery 240**
R. Avery Bennett
- 18 Ferret Soft Tissue Surgery 277**
Catriona MacPhail
- 19 Rodent Soft Tissue Surgery 297**
R. Avery Bennett
- 20 Soft Tissue Surgery in Hedgehogs 322**
Daniel J. Duffy and R. Avery Bennett
- 21 Surgery of the Sugar Glider 332**
Geoffrey W. Pye
- 22 Small Mammal Dental Surgery 338**
Estella Böhmer
- 23 Large Mammal Dental Surgery 369**
Allison D. Woody, David A. Fagan, and James E. Oosterhuis
- 24 Primate Surgery 383**
Celia R. Valverde and Elizabeth Bicknese
- 25 Marine Mammal Surgery 402**
Jennifer L. Higgins, Carmen M. H. Colitz, and Dean A. Hendrickson
- 26 Megavertebrate Laparoscopy 418**
Mark Stetter and Dean A. Hendrickson
- 27 Zoo Animal Surgery 426**
Geoffrey W. Pye
- 28 Surgical Oncology in Exotics 432**
Elizabeth A. Maxwell
- Index 448**

List of Contributors

R. Avery Bennett, DVM, MS, DACVS

Adjunct Professor of Companion Animal Surgery
Department of Small Animal Clinical Services
School of Veterinary Medicine, Louisiana State
University
Baton Rouge, LA, USA

Elizabeth Bicknese, DVM, MPVM, cVMA

Senior Veterinarian, Veterinary Services, San Diego
Zoo Wildlife Alliance
San Diego, CA, USA

Estella Böhmer, DVM, Dr. med vet

Clinic of Small Animal Surgery and Reproduction,
Center of Veterinary Clinical Medicine
Ludwig-Maximilians-University Veterinärstr
Munich, Germany

Carmen M. H. Colitz, DVM, PhD, MBA, DACVO, DCLOVE (hon)

All Animal Eye Care in Jupiter Pet Emergency and
Specialty Center
Jupiter, FL, USA

Brett Darrow, DVM, DACVS, CCRP

Staff Veterinary Surgeon
Capital Veterinary Specialists
Department of Veterinary Surgery
Jacksonville, FL, USA

Daniel J. Duffy BVMS(Hons.), MS, FHEA, MRCVS, DACVS-SA, DECVS

Assistant Professor Orthopaedic Surgery
North Carolina State University College of Veterinary
Medicine
Department of Clinical Sciences
Raleigh, NC, USA

David A. Fagan, DDS

Executive Director
The Colyer Institute
San Diego, CA, USA

Dean A. Hendrickson, DVM, MS, DACVS

Professor of Equine Surgery
Colorado State University College of Veterinary
Medicine and Biomedical Sciences
Department of Clinical Sciences, Veterinary Teaching
Hospital
Fort Collins, CO, USA

Jennifer L. Higgins, DVM, PhD

Colorado State University College of Veterinary Medicine
and Biomedical Sciences
Department of Clinical Sciences, Veterinary Teaching
Hospital
Fort Collins, CO, USA

Gregory A. Lewbart VMD, MS, DACZM, DECZM (ZHM)

Professor of Aquatic, Wildlife, and Zoological
Medicine
North Carolina State University
College of Veterinary Medicine
Department of Clinical Sciences
Raleigh, NC, USA

Catriona MacPhail, DVM, PhD

Professor of Small Animal Surgery
Colorado State University
Department of Clinical Sciences
Fort Collins, CO, USA

Elizabeth A. Maxwell, DVM, MS, DACVS-SA, CVPP

Clinical Assistant Professor, Surgical Oncology
University of Florida
Department of Small Animal Clinical Sciences
College of Veterinary Medicine
Gainesville, FL, USA

Michael S. McFadden, MS, DVM, DACVS-SA

Owner, Houston Mobile Veterinary Surgery
Magnolia, TX, USA

Stephen J. Mehler, DVM, DACVS-SA

Chief Medical Officer
Veterinarian Recommended Solutions
Blue Bell, PA, USA
Co-Founder
Main Line Veterinary Specialists
Devon, PA, USA

Michael B. Mison, DVM, DACVS-SA

Professor of Clinical Surgery
University of Pennsylvania School of Veterinary Medicine
Department of Clinical Sciences and Advanced Medicine
Philadelphia, PA, USA

James E. Oosterhuis, DVM

Principal Veterinarian
San Diego Zoo Safari Park
Escondido, CA, USA
Director of Research and Medical Care
The Colyer Institute
San Diego, CA, USA

Heidi Phillips, DVM, DACVS-SA

Associate Professor, Small Animal Surgery
University of Illinois College of Veterinary Medicine
Department of Veterinary Clinical Medicine
Urbana, IL, USA

Geoffrey W. Pye, BVSc, MS, DACZM

Animal Health Director
Disney's Animals, Science, and Environment Team
Disney's Animal Kingdom, Bay Lake, FL, USA

Mark Stetter, DVM, DACZM

Dean & Professor
Colorado State University
Department of Clinical Sciences
College of Veterinary Medicine & Biomedical Sciences
Fort Collins, CO, USA

Claire Vergneau-Grosset, DVM, IPSAV, CES, DACZM

Assistant Professor – Zoological Medicine Service,
Veterinarian of the Aquarium du Québec
Faculté de médecine vétérinaire, Université de Montréal
Saint-Hyacinthe, QC, Canada

Celia R. Valverde, DVM, DACVS

SAGE Veterinary Referral Centers-Dublin
Dublin, CA, USA

E. Scott Weber, III, MSc., VMD, Dipl ACVPM, CertAqVet

White Marsh Hollow
Conestoga, PA, USA

Allison D. Woody, DVM, DAVDC

Clinical Veterinarian
The Colyer Institute
San Diego, CA, USA
San Diego Animal Dentistry
San Diego, CA, USA

Preface

We have two main goals with the “Surgery of Exotic Animals”: (i) collate surgical information on exotic species into one book rather than have it scattered throughout the literature and (ii) ensure that every chapter is written and/or edited by a surgery specialist. Exotic animal surgery developed along a different pathway compared to domestic species because surgery specialists have felt out of depth with species with which they had no familiarity. Consequently, surgical techniques were developed by veterinarians with a special interest in exotics, but no specialized training in surgery. Principles instilled during a formal training program in surgery apply across species, including humans, but are only touched on during veterinary school. Veterinary surgeons are able to perform the procedures; however, they feel uncomfortable with the husbandry, anesthesia, and preoperative and postoperative patient care. Occasionally, a specialist surgeon is found with a special interest in exotic species. For the majority of this book, it is these rare individuals that have written chapters using sound surgical principles paired with knowledge of exotic species. An exotic specialist with a special interest in surgery or the pairing of a surgery specialist with an

exotic animal specialist is used for some chapters. The latter was also used for editing the book to ensure both aspects were considered in every chapter. These partnerships should be encouraged as it can be hugely beneficial for the animal as each person will bring their skillset to the table. The surgeon can bring strong surgical principles and the exotic animal specialist can help them extrapolate those to the unique anatomy of and what may or may not work for the exotic species. Diplomates of the American College of Veterinary Surgeons, Diplomates of the American Veterinary Dental College, and specialist surgeons from the human medicine field are great resources for exotic animal veterinarians and are at the cutting edge of new techniques, equipment, and products that could be used in exotic patients. These specialists need to be encouraged to take the leap into the unknown that exotic animal veterinarians take on a routine basis. Solid surgical principles used in domestic species and humans can be readily extrapolated for use in exotic animals in most cases.

Avery and Geoff
FL, USA
2020

About the Editors



Dr. R. Avery Bennett recently retired from Louisiana State University where he served for four years as professor of Companion Animal Surgery and now remains as an adjunct professor. He has served in senior positions in academia and the public sector in both surgical and exotic animal fields at the University of Florida, University of Pennsylvania, University of Illinois, Animal Medical Center – New York, San Francisco Zoo, and private surgical referral practices in California and Florida. Avery earned his doctorate of Veterinary Medicine with high honors from Michigan State University (1983). He completed a rotating internship in small animal medicine and surgery at Texas A&M University (1984) and a small animal surgery residency at Colorado State University along with a master's degree in veterinary clinical sciences (1987). Avery is a Diplomate of the American College of Veterinary Surgeons (1988). While Avery is competent in both soft tissue and orthopedic surgery, his areas of interest include surgery of exotic animals, minimally invasive surgery – laparoscopic and thoracoscopic surgery, oncologic surgery, and microsurgery.



Dr. Geoffrey W. Pye serves Disney's Animals, Science, and Environment team as the Animal Health Director and joined the team in 2014 to lead the animal health and animal nutrition teams at Disney's Animal Kingdom, the Seas with Nemo and Friends at Epcot, the TriCircle D Ranch, Castaway Cay, and the Aulani Resort. He has also served as a senior veterinarian for the San Diego Zoo, Werribee Open Range Zoo, and Currumbin Sanctuary. Geoff completed his bachelor of veterinary science at the University of Melbourne (1988) and earned a master of science in wild animal health from the University of London in association with the London Zoo (1995). He completed an internship in zoological medicine at Kansas State University (1998) and a residency in zoological medicine at the University of Florida (2001). Geoff is a Diplomate of the American College of Zoological Medicine (2001).

1

General Principles, Instruments, and Equipment

R. Avery Bennett

Introduction

Many surgeries in exotic animal are analogous to those performed in other species. The size of some patients makes surgery more challenging. Appropriate preoperative work up, patient preparation, surgeon preparation, perioperative antibiotic therapy, thermal support, and hemostasis are essential for a successful outcome. Preemptive, multimodal analgesia has long been known to improve recovery from surgery and is especially important with wildlife and exotic prey species. Additionally, in recent years, patients' anxiety has become an important consideration when these patients have to be hospitalized. Use medications evaluated for the species having surgery to develop a preemptive analgesic and antianxiety treatment plan. When prey species experience anxiety, stress, fear, and pain, they often die for no apparent reason. Alternatively, they may recover from the surgical event only to die a day or two later, likely from these stresses. In pet species, it appears that if they are accustomed to being handled by humans they are more likely to survive the perioperative period. If multiple procedures need to be accomplished, as a general rule, it is better to perform multiple short anesthetic events and surgeries than to try to do everything under one long anesthetic event. This is especially true when imaging is needed for surgical planning. It is best to anesthetize the patient for imaging, then recover it and evaluate the study. Once a diagnosis is made and a plan developed, anesthetize the patient the next day for the surgical procedure.

It is important that patients resume eating for nutritional support as soon as possible after surgery. Many small patients are not able to undergo long periods of anorexia because they do not have the energy stores to support themselves because in their natural environment they are constantly eating. The nutritional needs of the patient must be addressed and supplemented as needed either per os, using a feeding tube, or intravenously.

Presurgical Considerations

Prior to performing surgery, it is vital to evaluate the patient and address any abnormalities. In many situations, fasting is recommended; however, sometimes it is not necessary. Hemodynamic support is important for all but the shortest surgical procedures. Patient and surgeon aseptic techniques should be followed for all surgeries regardless of size and species. Perioperative or therapeutic antibiotic therapy needs to be considered. Because small patients become hypothermic quickly, thermal support is essential even for diagnostic procedures done under anesthesia.

Patient Support

Small exotic animals are especially prone to developing perioperative complications such as hypovolemia from blood loss, hypothermia, and renal and respiratory compromise. It is important to evaluate the patient systemically prior to anesthesia and surgery and to address any abnormalities preoperatively. During anesthesia and surgery, provide fluid therapy to support the cardiovascular system. Monitor body temperature and take measures to minimize hypothermia. Take appropriate steps to minimize blood loss and take precautions to minimize the risk of surgical site infections.

Data Base

The ideal preoperative data base includes a complete blood count, serum or plasma biochemistry panel, and a urinalysis. Additional diagnostics may be indicated based on the species and the medical problem being addressed. For example, mice and rats are prone to mycoplasmosis pneumonia, and preoperative chest radiographs are indicated prior to anesthesia in these species. While it may be difficult to obtain a urine sample from a small rodent, they are prone to developing renal insufficiency later in life, so it is important to evaluate urine specific gravity and a dipstick for proteinuria. These can be done with two drops of urine

that can be obtained by placing the patient in a plastic or glass container for several minutes. In ferrets, it is important to determine the blood glucose level prior to anesthesia because they are prone to forming insulinomas. In very small patients, it may not be feasible to take enough blood for all of the abovementioned diagnostics. At a minimum determine a hematocrit, total protein, blood urea nitrogen (azotemia test strip), glucose, and urine specific gravity.

Preoperative Fasting

Various species such as equids, rabbits, and rodents are not able to vomit for physiologic reason, so fasting to prevent aspiration pneumonia is not necessary. Additionally, if attempting to decrease gastrointestinal contents for a surgery, it can take days to make a difference because the majority of ingesta is within the hindgut. A short fast is recommended to allow these species to swallow any food material to reduce the risk of food entering the trachea during intubation. A prolonged fast in small mammals can result in a negative energy balance which increases their risk for developing complications after surgery (Jenkins 2000). Many small patients have low hepatic glycogen stores and may develop hypoglycemia during a prolonged fast (Harkness 1993; Redrobe 2002). Administer fluids containing dextrose subcutaneously (SC), intravenously (IV), or intraosseously (IOs) in patients prone to developing hypoglycemia. The gastrointestinal transit time in ferrets is rapid, and a prolonged fast is not recommended. An hour fast in ferrets is long enough for the stomach to empty minimizing the risk of developing aspiration pneumonia. On the other end of the spectrum, some reptiles may only eat once a week or even less often so there is no need for a fast.

Hemodynamic Support

Small patients have a small total blood volume and what may appear to be minimal hemorrhage can be life-threatening. If the patient is anemic and surgery can be postponed, it should be postponed until the hematocrit is into the normal range. It would be a rare event that surgery made a hematocrit increase, typically the opposite is the norm. Consider a blood transfusion from a conspecific, if more than minimal hemorrhage is anticipated or if the patient is anemic preoperatively. Strict attention to intraoperative hemostasis is essential when performing surgery on any small patient.

In patients experiencing serious blood loss during surgery, crystalloid or colloid fluid therapy should be administered as quickly as possible for cardiovascular support. More ideal, blood from a conspecific should be used, but often this is not available. Preplanning by having a conspecific blood donor available can be life-saving. In ferrets, there are no blood types and no reports of transfusion reactions. It is safe to use any ferret as a blood donor. In many species, blood typing

may not be known. If it is unknown whether a species has blood types, a crossmatch should be performed prior to administering a blood transfusion.

Anesthesia results in loss of fluids because of dry gases making parenteral fluid administration vital for most surgical procedures. It can be difficult to achieve vascular access in small patients. Vascular access provides a route for the administration of fluids during anesthesia at the standard rate of 10 ml/kg/hr and, maybe more importantly, provides a route for administration of emergency drugs in the event of a crisis. An IOs catheter can be placed relatively easily in most species even in small patients. SC administration of fluids is much less effective than IV or IOs and is an ineffective route for administration of emergency drugs. A single dose of 10 ml/kg SC of 4% dextrose has been recommended for short procedures in healthy small exotic mammals (Redrobe 2002). Fluids administered subcutaneously or intraperitoneally are slowly absorbed and not appropriate for treatment of severely ill, dehydrated, or shocky patients.

Maintain vascular access in the postoperative period if at all possible. Continue to administer fluid at least at a maintenance rate until the patient has completely recovered and is eating and drinking well.

Perioperative Antibiotic Therapy

The aim in administering perioperative antibiotics is that the blood level of antibiotic will be effective in preventing incision site infection from target organisms. In most cases, the target organisms are normal skin flora that cannot be completely eliminated during patient skin preparation. The antibiotic should be administered prior to making an incision, which is when the first exposure occurs and should continue until the surgery is complete so any blood clots that form will have therapeutic levels of antibiotic. If the patient is already receiving antibiotics for treatment of an infection and has therapeutic circulating levels of antibiotics effective against the target organisms, additional IV perioperative antibiotic is not needed. If the therapeutic antibiotic is not expected to be effective against the surgical target organisms, perioperative antibiotic administration of an antibiotic expected to be effective against the surgical target organisms should be administered perioperatively. For example, if a patient is receiving cephalexin for a skin infection, but the surgery is in the perineal area where fecal contamination is a concern, adding a perioperative antibiotic against which fecal flora are likely to be sensitive is appropriate. There is no evidence that continuing to administer a perioperative antibiotic for a brief period of time postoperatively is beneficial for preventing surgical site infections (SSIs) and may select for resistant organisms.

Surgeon Aseptic Preparation

The goal of surgeon aseptic preparation is to reduce the incidence of SSI. In human medicine, SSI has been shown to delay wound healing, increase antibiotic use, increase hospital stay, and increase costs, and can result in fatal consequences. Minimizing the risk of SSI in exotic animals is essential for many reasons. Surgeons should wear a cap, mask, gloves, surgical gown, and shoe covers for all surgical procedures (Figure 1.1). Procedures should be done in the cleanest environment possible. A dedicated operating room is ideal, but avoid doing surgery in rooms where abscesses are treated, dirty procedures are performed, or there is fecal contamination.

In one study, glove perforation occurred in 67% of surgeries underscoring the need for hand antisepsis to reduce skin flora before gloving (Verwilghen et al. 2011). Scrubbing is *not* recommended because it removes protective mechanisms on the skin surface and exposes more potentially pathogenic bacteria. Additionally, it causes small abrasions and excoriations damaging the surgeon's skin and increasing the risk of colonization by pathogenic organisms.

Alcohol-based hand rubs are recommended because they have rapid action, are faster to use, and cause less skin damage compared with antiseptic soaps. Alcohol-based

hand rubs do not require water. It has been estimated that on average 20l of water are used per hand when using antiseptic soaps. Additionally, many water faucets and municipal water sources are contaminated with *Pseudomonas* sp. and other Gram negative organisms that can recontaminate the hands. There is no reason to do a one-minute hand wash with a neutral soap before applying the hand treatment, and it has been shown that omitting hand washing prior to applying Sterilium® (Medline Industries, www.medline.com) increases its efficacy (Verwilghen et al. 2011). The World Health Organization recommends alcohol-based hand rubs over antiseptic soaps because of their superior efficacy both *in vitro* and *in vivo*, better skin tolerance, lower environmental impact, and no risk of recontamination from rinsing with contaminated water.

Patient Preparation

Regardless of the patient size or species, standard aseptic technique is essential for reducing the risk of SSI. Commonly, exotic animals are more likely to develop SSI than domestic animals because their housing often require them to be in proximity to their urine and feces. Prepare an appropriate area around the proposed surgical incision. Some species like chinchillas and rabbits have fragile skin

Figure 1.1 The surgeon must adhere to proper aseptic technique wearing clean scrubs (a). A cap, mask, and shoe covers, and sterile gown and gloves are proper attire regardless of the size and species of the patient (b).



that can easily be damaged during clipping for surgery. For most mammals, a No. 40 clipper blade is used; however, in patients with very fine fur a No. 50 clipper blade (Oster Professional Products, McMinnville, TN) is better suited. The teeth of the No. 50 blade are closer together than those of the No. 40, making it more difficult for the patient's skin to get caught between the teeth and be cut. Move slowly over the skin and keep the skin in front of the clipper blade flattened. Clean and lubricate the blade frequently to help decrease the risk of damaging the skin.

In birds, it is best to pluck feathers which then regrow rapidly; however, it is best not to pluck primary flight and tail feathers as this can damage the follicle resulting in abnormal feather growth. Some birds, such as penguins, have very small dense feathers that are very difficult to pluck. Use a No. 10 clipper blade for these birds. The No. 10 blade teeth are farther apart and clip the feathers at the base. Cut or clipped feathers will not be replaced until the shaft is molted and replaced naturally.

For many years, it has been suggested that alcohol be avoided for skin preparation of small patients because evaporative cooling might cause clinically important hypothermia. A study in mice compared the effect of skin preparation on hypothermia (Skorupski et al. 2017). They compared alcohol alone, povidone iodine alone, povidone iodine alternating with alcohol or saline or warmed saline. Alcohol alone or alternating with povidone iodine caused a dramatic drop in surface and core temperatures, but it rebounded quickly. When alcohol was used alone, the these temperatures returned to nearly the level of the control animals within minutes; however, when alternating with povidone iodine, this was not observed to occur. The authors theorized that the alcohol evaporates very quickly and the cooling stops, whereas povidone iodine and saline evaporate over a longer period of time resulting in more profound hypothermia. Using warmed saline did not modify this effect. The coldest core temperatures were observed in all povidone iodine treatment groups, and this hypothermia persisted. The authors suggested the povidone iodine takes longer to evaporate than saline, resulting in these low body temperatures. The authors concluded that using alcohol for rodent aseptic preparation causes less hypothermia and should be encouraged.

In that study, three applications alternating povidone iodine with either saline or alcohol was used. In veterinary, surgery tradition has been to apply alternating patient skin preparation solutions such as povidone iodine or chlorhexidine alternating with alcohol or saline. The author has been unable to find any scientific evidence for this protocol. It is not applied to humans where a single application of a skin preparation solution is applied prior to making an incision. In fact, the instructions for use on the stock bottle of chlo-

rhexidine indicate to apply the solution and allow it to remain in place for two minutes, then wipe it off and be done.

In another study, an alcohol-based skin preparation was compared to chlorhexidine for reducing skin surface bacteria (Maxwell et al. 2018). The alcohol-based skin preparation sprayed onto the surgical site and allowed to dry was equally as effective as chlorhexidine used according to manufacturer instructions. Make sure the alcohol has completely evaporated prior to using electrosurgery or a CO₂ laser that can ignite the alcohol.

It is not necessary to scrub the patient's skin and actually may be contraindicated. Scrubbing not only irritates the skin, but it also exposes deeper and more pathogenic bacteria to the skin surface potentially contaminating the deeper tissues when the incision is made. A study in horses comparing mechanical and nonmechanical sterile preoperative skin preparation with chlorhexidine gluconate showed there was no difference between a five-minute mechanical scrub and a five-minute application with no scrubbing (Davids et al. 2015). The manufacturer's instructions for using chlorhexidine direct that it be applied to the site and allowed to remain for two minutes before wiping it off. Wiping it off with either saline or alcohol may remove the chlorhexidine negating its residual activity and should, therefore, not be done. A need for alternating between a preparation solution and saline or alcohol has never been established in any species. A newer skin preparation product, Chloraprep™ One Step (BD Medical, www.bd.com) is applied to the skin for 30 seconds and left to dry for 30 seconds after which the skin is ready for incision. The World Health Organization Global Guidelines for Prevention of Surgical Site Infections states "The panel recommends alcohol-based antiseptic solutions based on chlorhexidine gluconate for surgical skin preparation in patients undergoing surgical procedures" (www.who.int/gpsc/ssi-guidelines/en/).

Patient Draping

Proper aseptic technique is indicated for all surgical patients regardless of size and species. Place quarter drapes around the surgical site and then cover with a patient drape with a fenestration around the site. Currently, disposable paper drapes are commonly used for both quarter drapes and patient drapes. Avoid the tendency to make a small sterile field as it is then easy to contaminate yourself by accidentally touching objects on the table outside the sterile field. Create a sterile field of the entire table top and take appropriate action to reestablish a sterile field if contamination occurs (Figure 1.2). When working with very small patients, it may be beneficial to use a clear plastic drape over the patient to allow respiration to be monitored; however, other respiratory monitoring devices such as end-tidal CO₂ monitors allow monitoring of respiration more accurately and do not require the ability to visualize the patient.




Figure 1.2 This 1.2 kg patient is being draped for a thoracotomy. First quarter drapes of Huck towels are placed to isolate the surgical field (a). Then the patient is covered with a patient drape large enough to create a sterile field of the entire table (b). The instruments are on the back table, and this is pulled up to the patient drape to complete the sterile field (c).

Thermal Support

Hypothermia during anesthesia is a major concern for many reasons including decreased metabolic functions and excretion of anesthetic agents. Many small exotic animals have a large body surface area/volume ratio predisposing them to developing hypothermia during anesthesia. The body temperature of a rat can drop 18 °F (10 °C) after 20 minutes of anesthesia (Harkness et al. 2010). Use an esophageal or rectal temperature probe to continuously monitor the patient's body temperature. The normal temperature for a given species may not be known; however, once the patient is anesthetized insert a temperature probe to determine its body

temperature. Then monitor the temperature and how much it drops during anesthesia. Even in poikilotherms, this is very helpful and can help make decisions about the need to provide additional thermal support.

A short anesthesia and operative time accomplished by having all the necessary equipment ready and accessible will help to minimize hypothermia. It is best to use patient warming devices with temperature control such as forced warm air and warm water blankets. It can be difficult to place a warm air blanket around a small patient. These devices blow warm dry air, so pay particular attention to keeping the cornea lubricated to reduce the risk of creating corneal ulcers. This is

especially important with small patients where the head is hard to visualize under the blanket. It is considered beneficial to use more than one patient warming device. For example combine a circulating warm water blanket under the patient and a forced warm air device around and over the patient where possible. For most small exotic animals set the circulating warm water blanket and forced warm air device at 104°F (40°C) because these patients have a high body temperature. Drape the patient as quickly as possible to help hold heat in under the drapes.

Use warm (101–103°F; 39–40°C) saline for abdominal and thoracic lavage. Fill the body cavity with the warm saline and do not immediately remove it. Allow it to dwell within the body for several minutes. Repeat this process until the body temperature begins to rise. Once the downward trend in body temperature reverses and the body cavity has been closed, in most cases, the temperature will continue to rise assuming other patient warming devices are in place. For most species, if the body temperature drops below 96°F (35.5°C) during the surgery, stop the procedure and instill warm saline into the body cavity before continuing the procedure. Repeat the process before closing.

Instrumentation and Equipment

Hemostatic Aids

The average cotton-tipped applicator (CTA) holds approximately 0.1 ml of blood when completely soaked and a 4×4 gauze sponge holds 10±2 ml (Hughes et al. 2007). The total blood volume of small mammals is 57 ml/kg of body weight (Jenkins 2000; Bennett 2009). If a patient loses 10–15% of total blood volume (approximately 1% of body weight), it is generally considered safe. Most mammals experience hypovolemic shock and release large amounts of catecholamines with loss of 15–20% of the total blood volume. Loss of 20–30% of the total blood volume can have life-threatening consequences (Jenkins 2000; Bennett 2009). As an example, if a 50 g mouse loses more than five CTAs full of blood, it is equivalent to more than 20% of the blood volume (Harkness 1993). Table 1.1 shows approximately 20% of the blood volume of common rodent pets.

Magnification is an aid for hemostasis (see Chapter 3). What would seem a small amount of blood to the naked eye observer appears to be major blood loss when magnified drawing the surgeon's attention to the hemorrhage and the need to arrest it. Additionally, small vessels can be identified and controlled more easily when working under magnification.

Hemostatic clips are available in various sizes including microclips. These are very useful for controlling hemorrhage in small patients, where it is difficult to accurately place a ligature. Additionally, they can be applied in deep,

Table 1.1 Loss of approximately 20% of total blood volume results in cardiovascular compromise with potentially life-threatening consequences.

Species	20% of blood volume (ml)
Gerbil	1.2
Hamster	1.4
Rat	4.0
Guinea pig	4.5

Twenty percent of the approximate blood volume of rodents.

Source: Harkness (1993), Jenkins (2000), Harkness et al. (2010).

hard-to-reach locations. With some types of hemostatic clips, both straight and right-angled applicators are also available (Figure 1.3). This applicator makes it easier to get under a hemostat to apply a clip to a severed vessel.

Sterile CTAs are useful for atraumatic tissue dissection and manipulation. They can be used to apply pressure to damaged vessels allowing a clot to form. Use moistened CTAs when using them for tissue manipulation and dissection and dry ones to absorb fluids.

Absorbable gelatin sponge is made from treated purified gelatin solution. It is capable of absorbing many times its weight in blood and also provides a scaffolding for clot formation. It is completely absorbed in 4–6 weeks. Surgicel (Ethicon, Inc., Somerville, NJ) is oxidized regenerated cellulose resembling cloth. It is a hemostatic aid that adheres nicely to moist tissues, but is not capable of absorbing much fluid and does not adhere well to dry tissue. Gelatin sponges are thick and absorb more fluid than oxidized regenerated cellulose but are prone to becoming dislodged from the tissue.

Topical thrombin is commercially available and when applied uses the patient's fibrinogen to form a fibrin clot. It is most useful for oozing and minor bleeding from capillaries and venules where standard surgical techniques are ineffective at halting the hemorrhage. It can be applied directly to the source of hemorrhage or can be applied onto gelatin sponge that is then placed over the site of hemorrhage. A hemostatic “taco” can be made of gelatin sponge wrapped in oxidized regenerated cellulose and soaked in topical thrombin and is very effective for controlling hemorrhage (Figure 1.4).

Electronic Hemostatic Devices

Electrocautery

Electrocautery uses direct current electricity to heat metal until it is red hot. The red hot tip is applied to the source of hemorrhage to heat and coagulate tissues. Electrocautery causes heat damage to adjacent tissues that then undergo necrosis minimizing its value in many situations. Cautery pens (Convenient




Figure 1.3 Straight and right-angled hemostatic clip applicators are available. Compared with the standard applicator (a), the right-angled applicators (b) can be inserted into a small body cavity to place a hemostatic clip at nearly a right angle, while the standard applicator requires the handles to be almost perpendicular to the vessel, making it difficult to place through a small approach into a body cavity.




Figure 1.4 A “hemostatic taco” made of gelatin sponge (G) wrapped in oxidized, regenerated cellulose (S) and soaked in topical thrombin has been applied to the caudal vena cava (arrow) after right adrenalectomy.

Cautery Kit; Jorgensen Labs; www.jorvet.com) may be useful to cauterize small vessels in small exotic mammals if electro-surgery is not available. They have a fine wire tip and are battery-operated. Keep in mind that the longer the heat is applied to the tissue, the more thermal necrosis occurs.

Electrosurgery

Electrosurgery uses high-frequency alternating current to generate energy. There is an active electrode and an indifferent electrode or ground. With monopolar electrosurgery the current exits the generator, passes to the active electrode, and concentrates at the tip which contacts the patient. After creating a thermal event at the tip, the current disperses throughout the patient's body, exits through the ground plate, and is carried back to the generator to complete the electrical circuit. Therefore, the patient is within the electrical circuit. If the ground plate only contacts a small area of

the patient's skin burns and subsequent necrosis can result because the current concentrates where it exits the patient. The Surgitron® Dual Frequency 120 (Ellman International, Inc., Hewlett, NY) generates 4.0 MHz current which is in the radio frequency wavelength. Current is received by the indifferent electrode acting as an antenna. Because the current is in the radio frequency, the area of contact between the patient and the ground plate is irrelevant and thermal injury should not occur. This is helpful with small patients where it might be difficult to disperse the current exiting the patient adequately.

The bipolar electrosurgical forceps are primarily used for hemostasis, but can also be used for tissue dissection resulting in minimal hemorrhage. The same generator is used, but the patient is not within the electrical circuit so the current only affects tissue within the tips of the forceps. Bipolar forceps are most useful near structures that would be adversely affected by the current such as the heart, brain, spinal cord, etc., or for hemostasis deep within body cavities. When using bipolar to dissect or cut tissue, grasp the tissue in the forceps and activate the current while pulling the forceps. The tissue within the tips will be affected by the current providing hemostasis for small vessels within the tissue grasped. In birds, this technique is useful for making a skin incision. Grasp the skin with the forceps and activate the current to create a small defect in the skin. Then insert one limb of the forceps under the skin for 1–2 cm, oppose the other limb of the forceps, activate the current, and withdraw the forceps, thus cutting the skin. Bipolar mode is available on most electrosurgical units including the Surgitron®.

Vessel Sealing Devices

LigaSure™ (Medtronic, Minneapolis, MN) is a vessel/tissue sealing system that is commonly used in veterinary surgery. The ForceTriad™ (Medtronic, Minneapolis, MN) combines the LigaSure technology with monopolar and bipolar

electrosurgery. While the specific technology is proprietary, LigaSure™ is a type of bipolar electrosurgery. The generator senses the electrical impedance of the tissues within the tips and delivers the amount of current needed to melt the elastin and collagen; it then allows the elastin and collagen to re-form, creating a permanent seal in a single application. It generates less than 1 mm of heat lateral to the seal and is Food and Drug Administration (FDA)-approved to seal vessels up to 7 mm diameter. A wide range of handpieces are available for open surgical procedures and for minimally invasive surgeries.

Harmonic Scalpel

Ethicon® Harmonic Scalpel (Ethicon US, LLC, www.ethicon.com/na) is an ultrasonically activated cutting instrument that converts electrical energy to mechanical energy. The blade vibrates at 55,500 Hz creating heat that coagulates vessels and cuts tissue. Hand pieces are available in various sizes and types for both open surgical procedures and minimally invasive surgery. It is FDA-approved to seal vessels up to 5 mm diameter and does not create smoke. The tissue is compressed between a Teflon anvil and the vibrating blade. There are two settings, one to seal the tissue and another to cut through the tissue.

Carbon Dioxide Laser

The carbon dioxide (CO₂) laser (Aesculight; LuxarCare LLC, Bothell, WA) gained popularity in veterinary surgery in the 1990s and continues to be in common use today. CO₂ lasers produce a beam of intense light at a wavelength that is highly absorbed by water molecules. It cuts tissue when the beam is very focused and coagulates for hemostasis when it is more diffused. The CO₂ laser seals vessels <0.6 mm, so most skin incisions are bloodless with minimal or no bruising. It also seals lymphatic vessels and nerves so there is less swelling and reportedly less pain associated with incisions compared with those made using a scalpel. When used with correct technique, the amount of lateral heat damage is minimal (Harkness et al. 2010); however, when used incorrectly, lasers can cause significant thermal damage.

In addition to being able to incise skin and other tissues, CO₂ lasers can also be used to fulgurate or destroy tissue, such as a tumor or the tissue bed after a tumor has been removed in an effort to eliminate residual microscopic disease. Use a high-power setting and keep the tip as close to the tissue as possible when cutting tissue. Do not touch the tip to the tissue. If there is hemorrhage, move the tip farther from the tissue, which will diffuse the beam and coagulate vessels. Use a smaller tip to create a small spot of light which will then cut better with less collateral heat. A wider tip will not cut as well and will cause more heat damage, but will also control hemorrhage better by coagulating vessels. Proper safety training for all personnel is important when using surgical lasers.

Magnification

Magnification is a vital part of efficiently performing surgery on small exotic animals, but it is also useful for delicate procedures in any size animal (see Chapter 3).

Focal Light

Overhead surgery lights are not adequate for surgery in small exotic animals and are becoming obsolete being replaced by head-mounted focal light sources that illuminate wherever the surgeon is looking, including deep into body cavities. These lights illuminate a smaller area allowing better visualization of the tissues. Inexpensive head-mounted cool focal lights are available with or without magnifying loupes from many sources (LED Headlight; MDS, Inc., Brandon, FL). SurgiTel loupes (General Scientific Corporation, Inc., Ann Arbor, MI) are available with different light options that mount onto the frame to illuminate the surgeon's field of view.

General Scientific Corporation, Inc., also offers SurgiCam HD, a head-mounted digital video camera with 3× magnification. The procedure can be viewed on a monitor, and images, and videos can be captured onto a computer.

Retractors

In the past, spring-loaded eyelid retractors have been recommended for abdominal wall retractors in small patients. The tension cannot be adjusted and is often too much for small patients. Heiss and Alm Self-Retaining Retractors work well as abdominal and tissue retractors in companion exotic animals (Figure 1.5). The Lone Star™ Retractor (Veterinary Specialty Products, Inc., Mission, KS) consists of a plastic frame and silastic bands with tissue hooks on the ends (stays) (Figure 1.6). Be sure to look for the models with all components that are fully autoclavable. Place the stay hooks in the tissue and using an appropriate amount of tension, pull and insert the silastic bands into the notches of the frame to retract tissue. These retractors are very versatile and have a wide range of applications.

Instrumentation

In the past, many surgeons have turned to using ophthalmic instruments for surgery in small exotic animals; however, they are not well suited for this purpose. Being made for surgery on a superficial structure, the eye, they are short making them more difficult to control and to manipulate tissues deep in a body cavity. Microsurgical instruments are manufactured so they are of a standard length and only the tips are miniaturized (Bennett 2009) (see Chapter 3). They are long enough for the instrument to




Figure 1.5 An Alm retractor (a) uses a thumb screw to open the jaws, while the Heiss retractor (b) has a quick release ratchet mechanism.




Figure 1.6 The Lone Star retractor system consists of a plastic ring with notches around the perimeter (a) and silastic bands with tissue hooks that are placed in the tissue to be retracted (b). Insert the bands into the notches in the ring to maintain tissue retraction.

balance in the hand while the tips extend beyond the surgeon's hand into the patient's body cavity (Figure 1.7). They should be counterbalanced to minimize hand fatigue. Round handles are recommended because the instruments should be rolled between the thumb and first two fingers rather than using wrist action. Having round handles is most important for needle holders because the curved needle has to be passed through the tissue using this rolling action between the thumb and fingers. Even with scissors and forceps, it is easier to roll them if the handles are round. Many surgeons prefer needle holders without box lock for delicate tissue because when the lock is set, the needle can jump and tear tissues. Hold a microsurgical instrument as if you were holding a pen (Figure 1.8a). An across-the-palm grip is inappropriate (Figure 1.8b). As a starting point, a microsurgical pack should contain a microsurgical needle holder, a microsurgical scissors, and microsurgical thumb forceps (Figure 1.9). Add other microsurgical instruments to the pack with increasing experience.




Figure 1.7 Microsurgical instruments should be a standard length with miniaturized tips that extend beyond the hand into the tissue and should be counterbalanced to minimize hand fatigue.

Suction and Irrigation

Ophthalmic bulb syringes work well for tissue irrigation in small exotic animals (Figure 1.10). Alternatively, use a




Figure 1.8 Hold the instruments like a pen (a) and not across the palm (b) for better control and the ability to roll the instrument.



Figure 1.9 The basic microsurgical pack should consist of a needle holder, scissors, and forceps.

syringe with a needle attached and bend the needle at the hub repeatedly until it breaks off to minimize the risk damaging tissue with the needle. Fine Barron suction tips have a release hole providing two degrees of suction; stronger



Figure 1.10 Ophthalmic bulb syringe is useful for irrigation in small patients.

suction with the hole covered and more delicate suction with the hole open. If tissues suck up against the tip, place the corner of a gauze over the end of the suction tip so it is not able to suck up tissue.

References

- Bennett, R.A. (2009). Rodents: soft tissue surgery. In: *BSAVA Manual of Rodents and Ferrets* (eds. E. Keeble and A. Meredith), 73–79. Gloucester: British Small Animal Veterinary Association.
- Davids, B.I., Davidson, M.J., Tenbroeck, S.H. et al. (2015). Efficacy of mechanical versus non-mechanical sterile preoperative skin preparation with chlorhexidine gluconate 4% solution. *Veterinary Surgery* **44** (5): 648–652.
- Harkness, J.E. (1993). Anesthesia, surgery. In: *A Practitioner's Guide to Domestic Rodents* (ed. J.E. Harkness), 37–50. Denver, CO: American Animal Hospital Association.
- Harkness, J.E., Turner, P.V., VandeWoude, S., and Wheeler, C. (2010). *Harkness and Wagner's Biology and Medicine of Rabbits and Rodents*, 5e. Ames, IA: Wiley-Blackwell.
- Hughes, K., Chang, Y.C., Sedrak, J., and Torres, A. (2007). A clinically practical way to estimate surgical blood loss. *Dermatology Online Journal* **13** (4): 17.
- Jenkins, J.R. (2000). Surgical sterilization in small mammals. Spay and castration. *Veterinary Clinics of North America: Exotic Animal Practice* **3**: 617–627.
- Maxwell, E.A., Bennett, R.A., and Mitchell, M.A. (2018). Efficacy of application of an alcohol-based antiseptic hand rub or a 2% chlorhexidine gluconate scrub for immediate reduction in the bacterial population on the skin of dogs. *American Journal of Veterinary Research* **79** (9): 1001–1007.
- Redrobe, S. (2002). Soft tissue surgery of rabbits and rodents. *Seminars in Avian and Exotic Pet Medicine* **11**: 231–245.
- Skorupski, A.M., Zhang, J., Ferguson, D. et al. (2017). Quantification of induced hypothermia from aseptic scrub applications during rodent surgery preparation. *Journal of the American Association for Laboratory Animal Science* **56** (5): 562–569.
- Verwilghen, D., Grulke, S., and Kampf, G. (2011). Presurgical hand antisepsis: concepts and current habits of veterinary surgeons. *Veterinary Surgery* **40**: 515–521.

2

Suture Materials

Michael S. McFadden

Introduction

Suture materials play an important role in veterinary surgery, but many veterinarians overlook important details and try to use one suture material for too many applications. Most veterinarians with no postgraduate training in surgery select suture materials based on hospital policy or cost, whereas veterinarians with postgraduate training in surgery select suture material based on a more complete understanding of suture materials and indications. The ideal suture would be in place only as long as needed and would then immediately disappear. Tissues heal at different rates, so materials that are more slowly absorbed are best used only in tissues that take a long time to heal. Suture should cause minimal tissue reaction, and the surgeon should attempt to minimize the amount of suture buried because all suture is foreign material. The amount of inflammation caused by different suture materials is illustrated in Figure 2.1. With the exception of rodents and rabbits, there are relatively few studies specifically examining suture materials in exotic animals.

Suture Materials

The role of sutures is to maintain incised or injured tissue in apposition to allow the tissue to heal (Bellenger 1982; van Rijssel et al. 1989; Roush 2003). The ideal suture material would provide high tensile strength for a sufficient time to allow the tissue to heal, have good knot security, resist infection, and cause no inflammatory, immunogenic, or carcinogenic reactions. The reaction of tissue to sutures depends on several variables such as the type, quantity, and duration of suture implantation as well as the tissues into which suture is implanted.

Surgeons must be familiar with different suture materials in order to make the most appropriate choice for specific clinical situations (Ratner et al. 1994). It is important to have a thorough understanding of the tissues, healing time required, the time the suture retains sufficient tensile strength to support the tissues as they heal, and time to complete absorption. The suture that is chosen may affect wound healing, functional outcome, and cosmetics. The number of suture types available has increased dramatically, and each suture type has specific physical, handling, and tissue reaction characteristics. Sutures are classified in many ways based on their ability to be absorbed and whether they are single, stranded, or braided. Different suture properties can also affect the knot security which must also be considered when choosing suture materials (Marturello et al. 2014).

Absorbable Suture Materials

Absorbable suture materials are defined as materials that lose their tensile strength within 60 days of implantation (Tan et al. 2003). These sutures are either composed of different synthetic polymers or are of biologic origin. Absorbable sutures that are biologic in origin are broken down by phagocytosis, while synthetic polymers are absorbed by hydrolysis. Hydrolysis involves breaking down polymers into monomers by direct water cleavage, and the monomers are then absorbed and metabolized by the body (Tan et al. 2003). Common absorbable sutures include chromic gut, polydioxanone (PDS™, Ethicon Inc., Cincinnati, OH), poliglecaprone 25 (Monocryl™, Ethicon Inc., Cincinnati, OH), polyglactin 910 (Vicryl™, Ethicon Inc., Cincinnati, OH), polyglyconate (Maxon™, Covidien, Medtronic, Minneapolis, MN), synthetic polyester called Glycomer™ 631 (Covidien, Medtronic, Minneapolis, MN) comprised of glycolide (60%), dioxanone (14%), and trimethylene carbonate (14%) (Biosyn™,




Figure 2.1 Inflammatory reactions after implantation of different suture materials to demonstrate variation in inflammatory reactions. Samples were chosen as representative examples of zero (a), mild (b), moderate (c), and severe (d) inflammatory responses. Suture material was lost in (a) and (b) during processing, but surrounding inflammatory reaction is intact. Images are 20 $\times$ magnification.

Covidien, Medtronic, Minneapolis, MN), polyglycolic acid (DexonTM, Covidien, Medtronic, Minneapolis, MN), polyglytone 6211 (CaprosynTM, Covidien, Medtronic, Minneapolis, MN), and synthetic polyester called LactomerTM comprised of a copolymer of glycolide and lactide (PolysorbTM, Covidien, Medtronic, Minneapolis, MN).

Various companies manufacture generic versions of most of these suture materials. In one study comparing the same material made by two different companies (Ethicon, Inc., and Hyaiyin Medical Instruments Co. Ltd.), the materials had significantly different tensile strength and stiffness (De La Puerta et al. 2011). In some comparisons, the generic outperformed the brand name materials. Most studies of suture materials are conducted using brand name materials. Based on this study, direct comparisons between brand names and generic materials cannot be made.

Biologic Absorbable Suture Material

Chromic Gut

Chromic gut is a monofilament absorbable suture composed of purified connective tissue (mostly collagen) derived from either bovine or ovine intestine and is absorbed by phagocytosis and enzymatic digestion by macrophages (Bellenger 1982; Roush 2003). Chromic salts are used to coat the suture to help delay absorption, increase tensile strength, and decrease tissue reactivity. Chromic gut is rapidly absorbed in mammals, is completely absorbed in approximately 60–90 days, and only maintains tensile strength for 7–10 days (Ratner et al. 1994). Chromic gut has a tendency to cause increased inflammatory response compared to other absorbable sutures which makes it less desirable.

rhinoceros laparoscopy 422–423
 rockfish, enucleation 53
 rodents
 castration 430
 cutaneous and subcutaneous masses
 abscesses 307
 benign and malignant tumors 306
 cervical lymphadenitis 307
 mammary tumors 305–306
 scent glands 306–307
 Zymbal's glands 306
 ear surgery 315–316
 enucleation and exenteration 315
 gastrointestinal surgery 310–311
 cheek pouch eversion 309–310
 gastric trichobezoars 307–309
 hystrichomorph 299
 intraabdominal masses 313–314
 non-hystrichomorph 299–300
 pododermatitis 316
 recovery 316–318
 reproductive tract surgery
 cesarean section 304–305
 orchiectomy 297–301
 ovariectomy and ovariohysterectomy 301–304
 paraphimosis 301
 suture reactions in 16
 tail degloving 316
 thoracic surgery 314–315
 tumors 439–440
 urinary tract surgery 311–313
 rostral diverticulum, avian 190–191, 193–194
 rostral maxillary abscesses, rabbits 355–356

S

salpingohysterectomy, avian 170, 172–174, 216–217
 salpingotomy
 reptiles 98, 100
 avian 170
Scelopores 50–51
 scent glands
 ferrets 290
 rodents 306–307
 Schmidt prisms 28
 Schroeder-Thomas (ST) splints 117, 119, 233–234
 scissor beak, avian 144–147
 scorpions 40
 scrotal ablation, sugar gliders 332
 sea hare, suture materials 19
 sea horse, enucleation 53
 sea lions, lensectomy 404
 sea squirts 42

semitubular plates
 avian 122
 reptiles 80
 shell fractures
 repair, in apple snail 37–38
 reptile orthopedic surgery 81–82
 Shope fibromas, rabbits 436
 shoulder luxation, avian 135–136
 Siberian sturgeon 19
 silk, suture material 14
 Sils™ Port 208–209
 sinusitis, primates 387
 sinusotomy
 avian 190
 infraorbital diverticulum 191, 193
 mandibular diverticulum approach 193
 postorbital approach 193
 preauditory approach 193
 preorbital diverticulum 191, 195
 rostral diverticulum and maxillary chamber 190–191, 193–194
 suborbital approach 193
 skin preparation 4
 skin surgery
 amphibian 65
 fish 49–52
 skin tumors, rabbits 436–437
 small mammals
 abscess-treatment 353–355
 alveolectomy 364–365
 apicoectomy 353
 cheek tooth extraction
 extraoral approach 346–347
 intraoral approach 345–347
 mandibular osteotomy 347–349
 maxillary osteotomy 349
 condylectomy 364
 dorsal rhinostomy 357–358
 incisor extraction
 elodontoma treatment 341–345
 extraoral approach 338–343
 intraoral approach 338–339
 jaw fracture treatment 361–363
 lateral fenestration of nasolacrimal duct 356–357
 mandibular abscesses 349–353
 mandibular symphysiolysis stabilization 363–364
 mandibulectomy 364–365
 orthopedic surgery
 anatomy and physiology 227
 bone composition 230
 bone healing 230–231
 diagnostics 227–228

- femoral fractures 237–238
- fractures 231–236
- fracture stabilization principles 231
- humerus fractures 236
- joint disease 228–230
- metacarpal bones fractures 237
- pelvic fractures 237
- radius and ulna fractures 236–237
- stabilization of specific fractures 234
- tibia and fibula fractures 237–238
- trauma 229
- retrobulbar abscesses 358–361
- rostral maxillary abscesses 355–356
- small rodents, anatomy and physiology 227
- snakes
 - bone cortex 77
 - coelomic surgery 92–93
 - musculoskeletal system 74
 - ovariosalpingectomy 100
 - salpingotomy 100
- soft tissue sarcomas (STS), avian 434
- spica splint, avian 116
- spiders 40
- splay leg, rabbits 229
- spleen, ferrets 278
- splenectomy, ferrets 283–284
- splints
 - avian 114–119
 - reptiles 78–79
- sponges 35–36
- squamous cell carcinomas (SCC)
 - avian 433
 - hedgehog 326–327
- stainless steel
 - avian 17
 - plates 123
 - suture material 14
- stereopsis 24
- stifle joint instability, rabbit 229
- stifle luxation, avian 140–141
- stillbirth monodiscoid placenta 396
- subcutaneous ureteral bypass (SUB) device, rabbits 248
- sugar gliders
 - abdominal surgical approach 334
 - orchiectomy 332–333
 - ovariectomy and ovariohysterectomy 334–335
 - paracloacal glands removal 336
 - patagial surgery 335–336
 - penectomy 333–334
 - postsurgical self-mutilation 332
 - tail amputation 336
- surgical loupes
 - compound 27–28
 - illumination 30
 - magnification changer of 25
 - vs. operating microscope 29–30
 - principles 23
 - prismatic 28
 - selection 28–29
 - simple 27
 - two-lens system 23
- surgical oncology
 - applications 432–433
 - avian 432–435
 - alimentary tumors 434–435
 - bone tumors 434
 - hemangioma/hemangiosarcoma 434
 - integumentary and musculoskeletal tumors 432–434
 - lipomas/liposarcomas 434
 - soft tissue sarcomas 434
 - ferrets
 - endocrine tumors 437
 - integumentary tumors 437–438
 - reproductive tumors 438
 - rabbits
 - bone tumors 437
 - reproductive tract tumors 435–436
 - skin tumors 436–437
 - thymic neoplasia 436
 - urinary tract tumors 436
 - reptiles 440–442
 - rodents 439–440
- surgical site infections (SSIs) 3
- suture materials 11–19
 - absorbable 11–12
 - barbed and antibacterial-coated suture 14
 - biologic absorbable 12
 - biologic nonabsorbable 14
 - birds 17
 - ferrets 17
 - fish 18–19
 - inflammation caused by 11–12
 - invertebrates 19
 - nonabsorbable 14
 - rabbits 16–17
 - reptiles and amphibians 17–18
 - rodents 16
 - synthetic absorbable 13
 - tissue healing 15–16
- swim bladder surgery 54–55
- synthetic absorbable sutures 13
- synthetic polyester. *see* Glycomer™ 631
- syrinx, lateral approach in avian 198

t

tail amputation
 ferrets 291
 primates 431
 sugar gliders 336
 tail degloving, rodents 316
 tape leg splint, avian 117–118
 tarantulas 40
 tarsal luxation, avian 141
 tarsiers. *see* nonhuman primates (NHPs)
 tarsorrhaphy, amphibian surgery 67
 telemetry
 amphibians 70–71
 fish 58
 implantation, marine mammals 408–409
 tendons, healing time 15
 tenorrhaphy, luxation, avian 143
 thecal cell tumors, gerbils 301–302
 thermal support 5–6
 thoracic surgery
 rabbits 260
 lung lobe abscesses 261–264
 postoperative care 264–265
 thymoma and thymic lymphoma 260–261
 rodents 314–315
 thoracoscopy
 definition 204
 hilar lymph node 224
 lung lobectomy, partial lobectomy, and lung biopsy 223–224
 mediastinal mass biopsy and removal 224
 small mammals 222–223
 thoracotomy. *see also* cranial celiotomy, avian
 rabbit
 lateral 263–264
 ventral 262–263
 patient draping for 4–5
 rodents
 lateral intercostal 315
 ventral 315
 thorny-headed worms 392–394
 through-the-lens mounted (TTL) loupes 28–29
 thymic lymphoma, rabbit 260–261
 thymic neoplasia, rabbits 436
 thymoma, rabbit 260–261
 tibia fractures, small mammals 237–238
 tibiotarsus
 fractures, avian 117
 safety factor 112
 titanium plates 123
 tooth extraction, hippopotamuses 426

topical thrombin 6
 tracheal resection and anastomosis, avian 196, 198–199
 tracheotomy, avian 195–199
 transplantation, Coelenterates 36
 transvesicular percutaneous cystolithotomy (TPC),
 rabbit 249–250
 trauma
 reptiles 76–77
 small mammals 229
 trichobezoar
 rodents 307–309
 primate 394–395
 triclosan 14
 tubal ligations, nonhuman primates 397
 tube cystostomy, rabbits 250
 tube nephrostomy, rabbits 247–248
 tusks, elephants 370–371
 extraction 374–375

u

ulna fractures, small mammals 236–237
 upper respiratory tract surgery, avian
 choanal atresia 190–191
 devocalization 198
 hyperinflation of cervicocephalic air sac 200
 sinusotomy 190–195
 tracheal resection and anastomosis 196, 198–199
 tracheotomy 195–199
 ureteral stenting, rabbits 248
 ureterolithiasis, rabbits 247–248
 ureteroneocystostomy, rabbits 248–249
 ureterotomy
 rabbits 248
 rodents 312
 urethrolithiasis, rabbits
 tube cystostomy 250
 urethrotomy 250–251
 urethrotomy, rodents 313
 urinary tract surgery
 fish 56–57
 rabbits
 anatomy 245–246
 cystotomy 249
 inguinal urinary bladder herniation 252
 nephrectomy 246–247, 251
 nephrotomy 247
 partial cystectomy 251
 postoperative complications 252
 pyelotomy 247–248
 renal biopsy 251
 subcutaneous ureteral bypass device 248

- transurethral urinary bladder eversion and prolapse 252
 - transvesicular percutaneous cystolithotomy 249–250
 - tube cystostomy 250
 - tube nephrostomy 247–248
 - tumors 436
 - ureteral stenting 248
 - ureteroneocystostomy 248–249
 - ureterotomy 248
 - urethrotomy 250–251
 - urolithiasis 246–247
 - urochordates 42
 - urodeum
 - avian 175
 - reptiles 101
 - urogenital surgery
 - amphibians 68–69
 - ferrets 278–279, 288–289
 - tumors, reptiles 441
 - urolithiasis
 - chinchillas 311
 - ferrets 289–290
 - guinea pigs 311
 - rabbits 246
 - uterine adenocarcinoma, rabbits 435
 - uterine diseases, rodents 302
 - uterine prolapse, rodents 305
 - uterus
 - avian 170
 - ferrets 279
 - fish 56
 - hedgehog 322–324
 - mammals 96
 - primates 396, 397
 - rabbits 252, 254, 435
 - rhinoceros 422
 - rodents 215, 302, 303
 - sugar glider 334
- V**
- vascular access port (VAP), ferrets 292–293
 - vasectomy
 - avian 164, 168, 217, 219
 - megavertebrates 421
 - non-human primates 398–399
 - zoo animals 428–429
 - ventplasty, avian 185
 - ventral bulla osteotomy (VBO), rabbits 269–270
 - ventriculotomy, avian 181
 - ventriculus, avian 175
 - vessel sealing devices 7–8, 206
 - avian 206
 - veterinary cuttable plates (VCPs)
 - avian 122
 - reptiles 80
 - small mammals 233
 - Veterinary Orthopedics Implants (VOI) 123
 - veterinary standard precautions (VSP) 383
- W**
- wound healing 3, 11, 15, 18
 - amphibians 64–65
 - avian 118
 - fish 48–49
 - pinnipeds 409
 - small mammal dental 345
- X**
- Xenopus laevis* 17, 64
- Z**
- zebrafish 48
 - zoo animal surgery
 - contraception
 - castration 430
 - deslorelin implants 428
 - epididymectomy 429–430
 - melengestrol acetate implants 428
 - ovariectomy and ovariectomy 430
 - vasectomy 428–429
 - defanging primates and carnivores 431
 - diversity, extrapolation of techniques 426
 - onychectomy, large cats 430–431
 - partnerships with specialty surgeons 426
 - postoperative management 428
 - tail removal, primates 431
 - Zymbal's glands, rodents 306