

Jameel M. Al-Khayri
Lina M. Alnaddaf
S. Mohan Jain *Editors*

Nanomaterial Interactions with Plant Cellular Mechanisms and Macromolecules and Agricultural Implications

 Springer

Nanomaterial Interactions with Plant Cellular Mechanisms and Macromolecules and Agricultural Implications


Jameel M. Al-Khayri · Lina M. Alnaddaf ·
S. Mohan Jain
Editors

Nanomaterial Interactions with Plant Cellular Mechanisms and Macromolecules and Agricultural Implications

 Springer

Editors

Jameel M. Al-Khayri 
Department of Agricultural Biotechnology
King Faisal University
Al-Ahsa, Saudi Arabia

Lina M. Alnaddaf 
Department of Biotechnology
and Molecular
Albaath University
Homs, Syria

S. Mohan Jain 
Department Agricultural Sciences
University of Helsinki
Helsinki, Finland

ISBN 978-3-031-20877-5 ISBN 978-3-031-20878-2 (eBook)
<https://doi.org/10.1007/978-3-031-20878-2>

© The Editor(s) (if applicable) and The Author(s), under exclusive license to Springer Nature Switzerland AG 2023

Chapter 6 is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>). For further details see license information in the chapter. This work is subject to copyright. All rights are solely and exclusively licensed by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors, and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

Preface

Nanotechnology has the potential to revolutionize agricultural and food production processes. It is imperative to harness the advantages of nanotechnology to enhance plant nutrient availability, improve soil fertility and minimize nutrient losses, develop efficient nanomaterial-packaging contaminants and utilization as a biochemical delivery system in plants. Nanomaterials, owing to their high surface to volume area ratio, are more efficient than most traditional materials. Their nature allows the slow release and promotes efficient nutrient uptake by the crops. Consequently, a reduced number of active chemicals are incorporated in agricultural soils and are absorbed by the plants, thus curtailing the negative impact on the environment. Nanobiotechnology is a promising tool to manipulate plants for the benefit of humankind in terms of sustainable agriculture and food security. Realized agricultural benefits so far are enhanced seed germination, increased plant growth and higher crop yield as well as increased secondary metabolites accumulation for an improved defense system against stress factors including climate change. With increased beneficial prospects and reduced adverse environmental effects, nanomaterials are viewed as instrumental in curbing world hunger.

To fully understand the behavior of nanoparticles in the plant, it is essential to study the mechanism of absorbance and translocation of nanoparticles and their interaction with the plant cellular biochemical compounds and organelles. The movement and accumulation of nanoparticles within the plant are determined by the mode of entry and uptake process as well as biochemical and physical barriers before entering the cytoplasm. Therefore, the current challenge is a systematic understanding of the absorption, translocation, bioaccumulation, barriers and pathways of nanomaterials within the plants. Research concerned with the relationship between nanomaterials and plants has risen dramatically in recent years which have contributed to the current perspectives of nanomaterials influence on cellular processes including photosynthesis, photorespiration and pigment synthesis and accumulation. Other research focused on understanding the impact of nanomaterials on cellular macromolecules including carbohydrates, lipids, nucleic acids, proteins, hormones and antioxidant defense activities. Collectively these processes and biochemical compounds have implications on crop yield. Nonetheless, research is considered in its infancy and

more work is needed to fully comprehend the nature, mode of action and utilization of this incredible material in agriculture. The interaction between nanoparticles and plants depends on several aspects such as nanoparticle shape, size, surface characteristics, crystal chemistry and dose. Additionally, it depends on plant genotype, plant age, soil or medium associated plant growth and the application method. Numerous nanoparticles have growth regulating properties causing substantial increases in biomass and even improvement in nutritional quality, whereas, others have toxic effects on plants resulting in morphological, physiochemical and anatomical alterations. This book examines recent progress of the above-mentioned aspects with emphasis on the interaction between nanoparticles and plants on the cellular level. The book starts with an introductory chapter providing an overview of the impact of nanotechnology on plant cell biology and outline the presented topics within the context of the book theme. This is followed by 18 chapters grouped into 3 parts: Part I Cellular mechanisms, Part II Cellular macromolecules and Part III Agricultural implications.

Chapters are contributed by 63 globally recognized scientists from 14 countries and subjected to a rigorous review process to ensure quality presentation and scientific precision. Chapters begin with an introduction providing background and rationale followed by a detailed discussion of the topic accompanied by 71 high-quality color figures and 32 tables. Each chapter concludes with recommendations for future study directions and a detailed list of appropriate sources to encourage further reading. The book is an excellent reference source for plant scientists engaged in research related to cultivation and breeding, biotechnology and nanotechnology. It is suitable for both advanced undergraduate and postgraduate students specialized in biotechnology and agriculture. We extend our appreciation and gratitude to the contributing authors for their assiduous efforts and diligence and to Springer for the opportunity to publish this work.

Al-Ahsa, Saudi Arabia
Homs, Syria
Helsinki, Finland

Jameel M. Al-Khayri
Lina M. Alnaddaf
S. Mohan Jain

Contents

1	Introduction: Impact of Nanotechnology on Plant Cell Biology	1
	Lina M. Alnaddaf, Jameel M. Al-Khayri, and S. Mohan Jain	
Part I Cellular Mechanisms		
2	Effect of Nanomaterials on Water and Solutes Translocation in Plants	19
	Khaled F. M. Salem, Sawsan Abd-Ellatif, El-Sayed S. Abdel Razik, Mayada S. Fadel, Ahmed E. M. Elkhawas, Ezzat R. Marzouk, Mohamed A. M. Bassouny, and Amira A. Ibrahim	
3	Response of Plant Photosynthesis to Nanomaterials	49
	Sashi Sonkar, Prakash Kumar Sarangi, Brijesh Pandey, Anand Prakash, and Akhilesh Kumar Singh	
4	Impact of Nanomaterials on Chlorophyll Content in Plants	69
	Luis Páramo, Humberto Aguirre Becerra, José Emilio Ramírez Piña, José Antonio Cervantes Chávez, Ana A. Feregrino-Pérez, and Karen Esquivel	
5	Interactions of Nanomaterials with Plant Pigments	93
	Montcharles S. Pontes, Jaqueline S. Santos, Simone Y. Fernandes, Ivan P. Oliveira, Thaiz B. A. R. Miguel, Emílio C. Miguel, Gilberto J. Arruda, Renato Grillo, Anderson R. L. Caires, and Etenaldo F. Santiago	
6	Impact of Nanomaterials on Plant Secondary Metabolism	133
	Rajendran K. Selvakesavan, Dariusz Kruska, Preeti Shakya, Dibyendu Mondal, and Gregory Franklin	

7	Toxic Effects of Nanomaterials on Plant Cellular Mechanisms	171
	Ana A. Feregrino-Pérez, Susana Meraz Dávila, Claudia Elena Pérez García, Alejandro Escobar Ortiz, Daniel Mendoza Jiménez, José Emilio Piña Ramírez, José Antonio Cervantes Chávez, and Karen Esquivel	
Part II Cellular Macromolecules		
8	Interaction of Nanoparticles with Plant Macromolecules: Carbohydrates and Lipids	213
	Lei Qiao and Chunlan Xu	
9	Interaction of Nanomaterials with Plant Macromolecules: Nucleic Acid, Proteins and Hormones	231
	Roseanne Mascarenhas, Tanvi Mathur, Jaya Maheshwari, and Praveen Nagella	
10	Influence of Nanomaterials on Non-enzymatic Antioxidant Defense Activities in Plants	273
	Antonio Juárez-Maldonado	
11	2D-Nanosheets Based Hybrid Nanomaterials Interaction with Plants	299
	Divya Chauhan, Mohammad Ashfaq, R. V. Mangalaraja, and Neetu Talreja	
Part III Agricultural Implications		
12	Nanomaterial Impact on Plant Morphology, Physiology and Productivity	319
	Mahroos A. Bahwirth, Salim F. Bamsaoud, and Lina M. Alnaddaf	
13	Role of Nanomaterials in Improving Crop Productivity	341
	Lina M. Alnaddaf, Jameel M. Al-Khayri, and S. Mohan Jain	
14	Role of Nanomaterials in Plant Cell and Tissue Culture	359
	Sanaz Feizi	
15	Role of Nanomaterials in Improving the Nutritional Value of Crops	399
	Mansoureh Nazari, Mojtaba Kordrostami, Ali Akbar Ghasemi-Soloklui, and Jameel M. Al-Khayri	
16	Role of Nanomaterials in Improving Crop Tolerance to Abiotic Stress	423
	Farhat Yasmeen	

17 Plant Mediation to Tolerate Cadmium Stress with Selenium and Nano-Selenium	455
Ali Akbar Ghasemi-Soloklui, Fardad Didaran, Mojtaba Kordrostami, and Jameel M. Al-Khayri	
18 Synthesis and Applications of Cellulose Nanomaterials Derived from Agricultural Waste and Byproducts	471
Amira A. Ibrahim, Sawsan Abd-Ellatif, El-Sayed S. Abdel Razik, Mayada S. Fadel, Ahmed E. M. Elkhawas, Mahmoud Shaban, Khaled F. M. Salem, and Mohamed F. M. Salem	
Index	501

Contributors

Sawsan Abd-Ellatif Bioprocess Development Department, Genetic Engineering and Biotechnology Research Institute, City of Scientific Research and Technology Applications, Borg EL-Arab, Alexandria, Egypt

Humberto Aguirre Becerra Graduate and Research Division, Engineering Faculty, Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

Jameel M. Al-Khayri Department of Agricultural Biotechnology, College of Agriculture and Food Sciences, King Faisal University, Al-Ahsa, Saudi Arabia

Lina M. Alnaddaf Department of Field Crops, Albaath University, Homs, Syria; Biotechnology and Molecular Biology, Faculty of Agriculture, Al-Baath University, Homs, Syria

Gilberto J. Arruda Plant Resources Study Group, CERNA, Mato Grosso do Sul State University (UEMS), Dourados, MS, Brazil

Mohammad Ashfaq Advanced Ceramics and Nanotechnology Laboratory, Department of Materials Engineering, Faculty of Engineering, University of Concepción, Concepción, Chile;
Department of Biotechnology & University Centre for Research & Development (UCRD), Chandigarh University, Mohali, Punjab, India

Mahroos A. Bahwirth Biology Department, Faculty of Science, Center for Natural and Applied Science Hadhramout Foundation for Invention and Advancement of Sciences, Hadhramout University, Mukalla, Yemen

Salim F. Bamsaoud Physics Department, Faculty of Science, Center for Natural and Applied Science, Hadhramout Foundation for Invention and Advancement of Sciences, Hadhramout University, Mukalla, Yemen

Mohamed A. M. Bassouny Soil and Water Department, Faculty of Agriculture, Benha University, Benha, Egypt

Anderson R. L. Caires Optics and Photonic Group, Institute of Physics, Federal University of Mato Grosso Do Sul (UFMS), Campo Grande, MS, Brazil

José Antonio Cervantes Chávez Natural Sciences Faculty, Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

Divya Chauhan Department of Chemical and Biomedical Engineering, University of South Florida, Tampa, FL, USA

Fardad Didaran Department of Horticulture, University of Tehran, Tehran, Iran

Ahmed E. M. Elkhawas Nucleic Acid Research Department, Genetic Engineering and Biotechnology Research Institute, City of Scientific Research and Technological Applications, Borg EL-Arab, Alexandria, Egypt

Alejandro Escobar Ortiz Graduate and Research Division, Engineering Faculty, Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

Karen Esquivel Graduate and Research Division, Engineering Faculty, Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

Mayada S. Fadel Bioprocess Development Department, Genetic Engineering and Biotechnology Research Institute, City of Scientific Research and Technology Applications, Borg EL-Arab, Alexandria, Egypt

Sanaz Feizi Agrobiotech for Health, Plant Biology and Soil Science Department, Faculty of Biology, University of Vigo, Vigo, Spain;
Department of Plant, Cell and Molecular Biology, Faculty of Natural Sciences, University of Tabriz, Tabriz, Iran

Ana A. Feregrino-Pérez Graduate and Research Division, Engineering Faculty, Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

Simone Y. Fernandes Plant Resources Study Group, CERNA, Mato Grosso do Sul State University (UEMS), Dourados, MS, Brazil

Gregory Franklin Institute of Plant Genetics of the Polish Academy of Sciences, Poznan, Poland

Ali Akbar Ghasemi-Soloklui Nuclear Agriculture Research School, Nuclear Science and Technology Research Institute (NSTRI), Karaj, Iran

Renato Grillo Environmental Nanochemistry Group, Department of Physics and Chemistry, São Paulo State University (UNESP), Ilha Solteira, SP, Brazil

Amira A. Ibrahim Botany and Microbiology Department, Faculty of Science, Arish University, Al-Arish, Egypt

S. Mohan Jain Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland

Antonio Juárez-Maldonado Department of Botany, Agrarian Autonomous University Antonio Narro, Saltillo, Mexico

Mojtaba Kordrostami Nuclear Agriculture Research School, Nuclear Science and Technology Research Institute (NSTRI), Karaj, Iran

Dariusz Kruszka Institute of Plant Genetics of the Polish Academy of Sciences, Poznan, Poland

Jaya Maheshwari Molecular Biology and Human Genetics, Manipal School of Life Sciences, Manipal Academy of Higher Education, Bangalore, Karnataka, India

R. V. Mangalaraja Advanced Ceramics and Nanotechnology Laboratory, Department of Materials Engineering, Faculty of Engineering, University of Concepción, Concepción, Chile;
Faculty of Engineering and Science, Universidad Adolfo Ibanez, Penalolen, Santiago, Chile

Ezzat R. Marzouk Division of Soil and Water, Faculty of Environmental Agricultural Sciences, Arish University, Arish, North Sinai, Egypt

Roseanne Mascarenhas Department of Life Sciences, CHRIST (Deemed to be University), Bangalore, Karnataka, India

Tanvi Mathur Department of Life Sciences, CHRIST (Deemed to be University), Bangalore, Karnataka, India

Daniel Mendoza Jiménez Natural Sciences Faculty, Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

Susana Meraz Dávila Graduate and Research Division, Engineering Faculty, Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

Emilio C. Miguel Laboratory of Biomaterials, Department of Metallurgical and Materials Engineering, Federal University of Ceará (UFC), Fortaleza, CE, Brazil

Thaiz B. A. R. Miguel Laboratory of Biotechnology, Department of Food Engineering (DEAL), Federal University of Ceará (UFC), Fortaleza, CE, Brazil

Dibyendu Mondal Institute of Plant Genetics of the Polish Academy of Sciences, Poznan, Poland

Praveen Nagella Department of Life Sciences, CHRIST (Deemed to be University), Bangalore, Karnataka, India

Mansoureh Nazari Department of Horticultural Science, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran

Ivan P. Oliveira Institute of Agricultural Sciences, Federal University of Minas Gerais (UFMG), Montes Claros, MG, Brazil

Brijesh Pandey Department of Biotechnology, School of Life Sciences, Mahatma Gandhi Central University, East Champaran, Bihar, India

Luis Páramo Graduate and Research Division, Engineering Faculty, Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

Claudia Elena Pérez García Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

José Emilio Piña Ramírez Natural Sciences Faculty, Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

Montcharles S. Pontes Plant Resources Study Group, CERNA, Mato Grosso do Sul State University (UEMS), Dourados, MS, Brazil

Anand Prakash Department of Biotechnology, School of Life Sciences, Mahatma Gandhi Central University, East Champaran, Bihar, India

Lei Qiao The Key Laboratory for Space Bioscience and Biotechnology, School of Life Sciences, Northwestern Polytechnical University, Xi'an, Shaanxi, China

José Emilio Ramírez Piña Natural Sciences Faculty, Universidad Autónoma de Querétaro, Santiago de Querétaro, Qro, México

El-Sayed S. Abdel Razik Plant Protection and Biomolecular Diagnosis Department, Arid Lands Cultivation Research Institute, City of Scientific Research and Technology Applications, Borg EL-Arab, Alexandria, Egypt

Khaled F. M. Salem Department of Plant Biotechnology, Genetic Engineering and Biotechnology Research Institute (GEBRI), University of Sadat City, Sadat, Egypt; Department of Biology, College of Science and Humanitarian Studies, Shaqra University, Qwaieah, Saudi Arabia

Mohamed F. M. Salem Department of Plant Biotechnology, Genetic Engineering and Biotechnology Research Institute (GEBRI), University of Sadat City, Sadat, Egypt; Department of Environmental Biotechnology, Genetic Engineering and Biotechnology Research Institute (GEBRI), University of Sadat City, Sadat, Egypt

Etenaldo F. Santiago Plant Resources Study Group, CERNA, Mato Grosso do Sul State University (UEMS), Dourados, MS, Brazil

Jaqueline S. Santos Plant Resources Study Group, CERNA, Mato Grosso do Sul State University (UEMS), Dourados, MS, Brazil

Prakash Kumar Sarangi College of Agriculture, Central Agricultural University, Imphal, India

Rajendran K. Selvakesavan Institute of Plant Genetics of the Polish Academy of Sciences, Poznan, Poland

Mahmoud Shaban Department of Chemistry, Faculty of Science, University of Alexandria, Alexandria, Egypt

Preeti Shakya Institute of Plant Genetics of the Polish Academy of Sciences, Poznan, Poland

Akhilesh Kumar Singh Department of Biotechnology, School of Life Sciences, Mahatma Gandhi Central University, East Champaran, Bihar, India

Sashi Sonkar Department of Botany, Bankim Sardar College, South 24 Parganas, West Bengal, India

Neetu Talreja Advanced Ceramics and Nanotechnology Laboratory, Department of Materials Engineering, Faculty of Engineering, University of Concepción, Concepción, Chile;

Department of Science, Faculty of Science and Technology, Alliance University, Anekal, Bengaluru, Karnataka, India

Chunlan Xu The Key Laboratory for Space Bioscience and Biotechnology, School of Life Sciences, Northwestern Polytechnical University, Xi'an, Shaanxi, China; School of Life Sciences, Northwestern Polytechnical University, 127 West Youyi Road, Xi'an Shaanxi, China

Farhat Yasmeen Department of Biosciences, University of Wah, Wah Cantt., Pakistan

Chapter 1

Introduction: Impact of Nanotechnology on Plant Cell Biology



Lina M. Alnaddaf, Jameel M. Al-Khayri, and S. Mohan Jain

Abstract This book focuses on the recent progress of nanotechnology, emphasizing the interaction between nanoparticles (NPs) and plants at the cellular level. Furthermore, it covers understanding of pathways of nanomaterials (NMs) entry into plant cells, their influence on cellular organelle processes, and their influence on total crop yield. It includes 17 chapters, grouped in three sections: (1) Cellular mechanisms, (2) Cellular macromolecules, and (3) Implications of NMs. These chapters provide details on plant response to NMs applications including morphological, physico-chemical, and anatomical changes and their effect on plant growth and productivity. The mechanisms of absorbance and translocation of NPs and their interaction with the plant cellular biochemical compounds and organelles are also covered. This book describes the current perspective of NMs' influence on cellular processes including photosynthesis and pigment synthesis and accumulation. Also highlights the current understanding of the impact of NMs on cellular macromolecules, these processes and biochemical compounds have implications for crop yield.

Keywords Adsorption · Cellular macromolecules · Cellular mechanisms · Cell biology · Plant nanotechnology

L. M. Alnaddaf (✉)

Biotechnology and Molecular Biology, Al-Baath University, Homs, Syria
e-mail: lalnaddaf@albaath-univ.edu.sy

J. M. Al-Khayri

Department of Agricultural Biotechnology, College of Agriculture and Food Sciences, King Faisal University, Al-Ahsa, Saudi Arabia
e-mail: jkhayri@kfu.edu.sa

S. M. Jain

Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland
e-mail: mohan.jain@helsinki.fi

1.1 Introduction

Nanotechnology contributes novel tools to impact and enhance crop production and provide an alternative approach for crop improvement. Nanomaterials (NMs) interact with various cellular macromolecules leading to both negative and positive effects, especially the enhancement of plant growth and resistance to different stresses. NMs optimize plant water and nutrient conditions, improving production quality and quantity. On the other hand, NPs are toxic to plant growth and human health via the food chain (Minkina et al. 2020).

The NPs gather in cell walls, tissues, and sub-cellular organelles such as chloroplasts and vacuoles, resulting in decreasing biological activity such as photosynthesis and metabolism in plant cells, reducing the germination rate and decreasing the length of root and shoot. They promote oxidative stress, antioxidant, nutritional imbalance for edible crops and quality of productivity. NPs in plant tissues have an impact on physiological processes (as promote or inhibit) and on the safety of macromolecules and cell organelles (Fedorenko et al. 2020).

There are essential factors related to the NPs and their interaction with plant tissues such as the plant species involved, which have physiological and anatomical differences, chemical characteristics of the plant's cellular surface, also environmental conditions surrounding it (Minkina et al. 2020).

1.2 Cellular Mechanisms

This section describes the interaction of NPs in different parts of plants (roots, shoots and leaves) at the cell organelles level (Fig. 1.1).

1.2.1 Adsorption

The first step is to absorb NPs via the roots and distribute it to plant tissues through modifications such as crystalline dissolution, biomodification and bioaccumulation. Plant roots and plant growth tissues are therefore the primary hosts for receiving NPs (Rao and Shekhawat 2016).

A variety of factors influence the absorption of NMs, including nanomaterial features, plant physiology, application method and environmental conditions.

NMs can interact with microorganisms and soil compounds that can positively or negatively modify absorption efficiency according to the type of NMs. Furthermore, multiple tissues (epidermis, endodermis) and barriers (Casparian strip, cuticle) need to be crossed before reaching vascular tissues, according to the entry point (roots or leaves) (Rajput et al. 2018).




Fig. 1.1 The impact of nanotechnology on plant cellular mechanisms (Constructed by L.M. Alnaddaf)

1.2.2 Penetration

The pathway of NMs to penetrate and uptake inside the cell can be via many ways, such as pore formation: Some NMs may disturb the plasma membrane, causing pore formation to pass via the cell; ion channels: These channels are about 1 nm in size, making it highly unlikely that NMs will penetrate them effectively without significant modifications; carrier proteins: NMs may bind to cell membrane proteins which act as carriers for entry into the cell; endocytosis: The NMs are embedded in the cell as a vesicle which may move in various compartments of the cell; and plasmodesmata (Schwab et al. 2016).

1.2.3 Cell Wall

NPs mainly modify the chemical composition and physical parameters of the cell wall affecting its structure. This modification is related to some factors such as the rate of pectins, structural proteins and phenolic compounds, cellulose, and hemicellulose. Also, pH, cell-wall existing enzymes, and acid cell wall properties regulate its porosity (Milewska-Hendel et al. 2017).

There are some characteristics NPs possessed that allow them to pass through cell walls such as size, shape, dimensions, The surface charge of NPs, composition, concentrations, the amount of fertilizer used, plant species, and conditions of the environment. The interaction between NPs and plants, whether stimulating or inhibiting, varies according to the previous characteristics (Milewska-Hendel et al. 2021).

1.2.4 Translocation

NMs move via plant tissue in the apoplast and the symplast. Also, the importance of the way NMs move inside plants gives indications about where they reach, end and accumulate (Milewska-Hendel et al. 2021).

This book presents several examples of translocation and accumulation of NPs in plant tissues, whether applied as soaked grains, fertilizers and foliar applications. For instance, NMs of carbon-coated iron are transferred to the aerial parts in pea and wheat faster than in sunflower and tomato (Perez-de-Luque 2017).

Also, this process is related to the size of the material and the zeta potential when carbon dots as a model is used in the translocation via the leaves in corn (*Zea mays* L.) and cotton (*Gossypium hirsutum* L.). Moreover, FeONPs penetrate the roots in (*Cucurbita pepo* L.) and accumulate in the roots' cells without transferring to leaves or flowers due to NPs magnetic properties (Tombuloglu et al. 2020).

1.2.5 Photosynthetic

A nanotechnology is a feasible tool for optimizing photosynthesis. It is related to primary metabolism and is responsible for generating plant biomass. Also induces plant growth and development (Sáez et al. 2017).

The effectiveness of NPs on photosynthetic (enhancing or impeding) and the functionalities of photosynthetic elements vary according to different plant species. This effect appears via an effect on regulatory proteins of the thylakoids, photosynthetic pigment (chlorophyll a and b), the activity of ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCo), carbon dioxide aggregation, adequate grana development and structural stability of mesophyll cells as well as chloroplasts (Kataria et al. 2019).

The enhancing or impeding influences of NPs on photosynthesis and plant growth are varied depending on nonmetallic NPs or metallic NPs. The stimulated reactions of nonmetallic NPs of photosynthesis are generated by biocompatible and reducing the oxidation processes. Also, NPs can be exploited for harvesting more electrons, which in turn, enhancement of plant photosynthesis and increases biomass and crop productivity (Swift et al. 2019).

Conversely, metallic NPs had impeding photosynthesis, which harmfully affects the different photosynthetic mechanisms. Impeding effects can be summed up for

NPs on the photosynthetic as toxicity via production of ROS and reduction each of the following: the net photosynthesis, chlorophyll content, photosystem II activities, the number of thylakoids per grana, the transpiration rate, leaf stomatal conductance, intercellular carbon dioxide concentration rate as well as inhibiting the expression of genes each of photosystem structure and chlorophyll biosynthesis (Poddar et al. 2020). These effects appeared via an application of different types of NPs, such as walled carbon nanotubes (SWCNTs), iron oxide, silver NPs, TiO₂ NPs, CuO NPs, ZnO NPs, carbon dots, Al₂O₃ NPs, Si NPs, Se NPs, Superparamagnetic iron oxide nanoparticles (SPION) for a variety of plant species (Kataria et al. 2019).

Carbon-based NMs interact with accessory pigments in the chloroplasts and chlorophyll-a, and chlorophyll-b and promote the ability of plants to harvest light energy. Whereas, NPs used as artificial antennae permit chloroplasts to absorb wavelengths that aren't essential for photosynthesis. This enhances the ability of plants to interact with light and optimize its capture, which increases the productivity of crops and enables the plants to adapt to different environments, where there is extremely solar radiation or limited resource of light (Aguirre-Becerra et al. 2020).

1.2.6 Pigments

NPs interact with plant pigments (Chlorophyll and derivatives, Carotenoids, Anthocyanins, and Betalains). Once it penetrates the plant chloroplast then the attachment occurs between NMs and pigments plant, which acts as cell protection agents, light-harvesting complexes, transfers absorbed light energy to chlorophyll molecules, dissipates excess energy to the environment and as antioxidant molecules when stress occurs (Nguyen et al. 2021).

Comprehension of the interaction mechanisms of NMs with plant pigments is critical to know its possible side effects on the biochemistry, metabolism, and physiology of plant organisms when NMs are absorbed. The plant produces more than 200,000 different chemical compound types including colored (pigment) compounds (Tripathi et al. 2020).

Engineered NMs interact differently with plant pigments depending on pigment type and their physical-chemical properties resulting in two various basic responses: the first is the pigment content (increase/decrease) by (promoting/inhibiting) the pigment synthesis. The second change is the pigment activities, especially energy dissipation processes and light absorption. Most of the literature indicated that plants respond to different types of NPs by reducing chlorophyll content followed by increasing the accessory pigment's contents. Thus, any functional or structural modifications in plant pigments, especially photosynthetic ones, affect the photosynthetic performance and biomass productivity (Nguyen et al. 2021).

NPs interact with plants to know about the implications of NMs on the plant morphological, physiological, production responses, and biological phenomena, to promote absorption of light wavelengths and optimize its capture and enhance photosynthesis (Santiago et al. 2020).

There are many factors that affect the content of the pigment in plant cells. For instance, biotic and abiotic stress induce the generation of reactive oxygen species (ROS) which increase the expression of genes involved in the pigment biosynthesis pathway, the synthesis route of NMs (green or chemically synthesized) and the usage dose (Tripathi et al. 2020).

1.2.7 Secondary Metabolites

NMs play an essential vital Role in plant cell and tissue culture. NPs have stimulated the biosynthesized secondary metabolites and gene expression for it (Wang et al. 2021).

NPs promote secondary metabolites (for instance, flavonoids, phenolic acids, glucosinolates, terpenoids, and alkaloid compounds). These play an antioxidant activity to enhance plant growth under biotic and abiotic stresses. As well as it decreases drought-induced damage such as *Z. mays* L., improves the quality of fruits under drought stress for *P. granatum* and increases yield. Also, the pharmacological properties of several medicinal plants are imputed to secondary metabolism compounds. Therefore, any change in bioactive compounds by NPs could affect their pharmacological properties and market importance (Ma et al. 2020).

Once the cells absorb the NMs, the metal-base of the NMs is turned into reactive metal ions, which react with functional groups present in a cell leading to a change in their biochemical activity. These interactions are various depending on plant species and tissues, size, solubility, concentration, shape, thermodynamic properties, composition, and surface coating. The main indicators resulting in NMs toxicity in the plant are the reduction of photosynthetic processes and the generation of ROS (Zahedi et al. 2021).

1.2.8 Phytotoxicity

The phytotoxicity caused by NPs is usually triggered by the release of free radicals such as hydrogen peroxide and hydroxyl radical. The excessive production of ROS interacts with various biological molecules and causes different cellular damage. It can also increase the level of oxidative stress, the fragmentation of peptide chains and alter the electric charge. Furthermore, it increases the susceptibility of various proteins to proteolysis, DNA and cell membrane damage, the toxicity of carbon-based NMs, the toxicity of metal and metal oxide NMs and anatomical and morphological changes (Kolenčik et al. 2021).

Exposure to NPs lead to adverse effects on several biochemical and physiological processes in different plant species which affects the growth stages of plants. Also, accumulated NMs in different plant parts can affect human health. In the contrast,

some studies have indicated that not all plants treated with NMs demonstrated toxic effects (Tombuloglu et al. 2020).

1.2.9 Gene Expression

In plants, numerous alterations resulting treated with NMs observed in genetic (DNA mutations) and epigenetic (DNA methylation pattern, histone modifications, and RNA interference). They were reflected in many aspects of the plant growth and development, such as cell divisions, chromosome behavior, mitotic aberrations, DNA alterations, and gene regulation related to forming the cell wall, roots, leaves, stress, and water channels (Khan et al. 2019).

SWCNT inhibited histone H3K9 acetylation in Maize seed's response to drought stress. Applying carbon nanotubes (CNT) in *Solanum lycopersicum* L. leads to the downregulation of genes related to roots and leaves. In contrast, increasing the upregulation of genes is related to water channels and stress (LeAqp2). In addition, Multi-walled carbon nanotube (MWCNT) had an impact on the upregulation of marker genes NNtPIP1, NtLRX1, and CycB, related to the formation of cell divisions and the cell wall (Khan 2020).

1.3 Cellular Macromolecules

Carbohydrates, nucleic acid, hormones, proteins, enzymes, and lipids are the main components of plants. These compounds contributed to NPs synthesizing as a safe and eco-friendly method. Also, they are closely associated with stress tolerance, growth and development of plants via increasing or decreasing their percentage when applying several NPs to plants (Khan et al. 2021).

1.3.1 Carbohydrates and Lipids

The chapter on the interaction of NPs with plant macromolecules: carbohydrates and lipids summarizes recent advances in the different effects of NPs to promote or inhibit plant growth, such as seed germination, plant root, and above-ground growth and improve various stress tolerance, which is closely associated with plant carbohydrates and lipids (Shang et al. 2019).

This stimulating or inhibiting is occurring using various types of NPs, for example, silver, selenium, zinc oxide, copper oxide, magnesium oxide, and silicon dioxide in different stages of growth and development of a plant. In addition, this chapter




Fig. 1.2 The impact of nanotechnology on cellular macromolecules (Constructed by L.M. Alnaddaf)

discusses the different effects of NPs added as a spray or fertilizer on morphological, biochemical, and productivity indicators of various plants (Khan et al. 2021) (Fig. 1.2).

1.3.2 Nucleic Acid

All genetic information for every living entity exists in nucleic acids (DNA and RNA) (Tan et al. 2009). This book presents a chapter to introduce the NPs interaction with nucleic acids. This interaction relates to type of NPs and concentrations.

In addition, it can use as delivery system DNA and RNA by binding interactions such as carbon nanotubes and bioclay (Hossain et al. 2016). Also, explains in more details the different effects either positive (activate the genes related to stresses) or negative (chromatin condensation and cells shrinkage, damaged DNA, fragmentation of chromosome arms and suppressing transcriptional genes) (Chen et al. 2018).

1.3.3 Hormones

There are five types of hormones that exist naturally in plants: auxin, cytokinin, gibberellins, abscisic acid and ethylene (Gaspar et al. 1996). NPs enhance the various physiological activities of hormones in plants (Weyers and Paterson, 2001). These activities relate to growth (elongation of roots or shoots), maturation, plant tissue culture and biotic or abiotic stress responses in a plant (Yang et al. 2017). While, other side effects are linked to senescence or phytotoxicity (El-Shetehy et al. 2020).

1.3.4 Proteins

Proteins are necessary for all living cells and have several functions such as regulation, cell signaling, catalysis, support, membrane fusion, intra- and intercellular movement of nutrients and other molecules, and structural protection. When NMs penetrate into plant cells, it reacts with carboxyl and sulfhydryl groups and alters protein activity (Anjum et al. 2015).

The NMs reaction depends on the physical and chemical conditions of the cell environment, which is affected by reactive molecules, and temperatures. Also, its sizes and concentrations affect the folding process of a newly synthesized protein (Hossain et al. 2016).

The plant responds to NMs treatment via several indicators such as an increase or decrease in proteins, accumulation or synthesis of new types of proteins that are involved in primary metabolism and production of enzymes that help the metabolic adaptation of the plant. In addition, Increasing proteins -are related to photosynthesis, metabolism, cellular organization, and hormone metabolism (Fig. 1.2) (Hasan et al. 2017).

1.3.5 Enzymatic and Non-Enzymatic Antioxidant

The influence of NMs on enzymatic antioxidant defense activities in plants differs according to the nature and concentration of NPs.

There is a regulatory suitable balance between ROS and antioxidants (enzymatic and non-enzymatic) in natural conditions. However, the plant responds with an increased amount of ROS concentration in conjunction with antioxidants when NPs stress occurs in the plant (Shang et al. 2019).

Antioxidants break down the ROS and scavenge it due to their peculiar structures and detoxify cells, such as oxygen free radicals and lipid peroxidation radicals.

In various researches, the impact of NPs on plant growth was different between promoting and inhibiting (Fig. 1.2) (Khalil et al. 2020).

1.4 Preparation and Features of NM

This book presents the methods to synthesize NM, such as physical, chemical and biological manufacture. Each method has a different feature that is related to its application (Alnaddaf et al. 2021). In addition, emphasizes the advantages of biosynthesis of monometallic NPs and includes some examples of silver, gold, copper, palladium and oxide NPs. Also, explains the factors that affect the NMs traits and behavior which make them able to penetrate plant cells (Shekhawat et al. 2021).

1.4.1 Nanocellulose

Cellulose is a natural polymer derived from agricultural waste and by-products used for the synthesis of various kinds of NMs. This chapter converses the synthesis of nanocellulose. Then, explains in more detail the source, structure, and types of nanocellulose. Also, it highlights the preparation, characterization, and properties of nanocellulose. In addition, it discusses the application of nanocellulose in many sectors (Zhang et al. 2022).

1.4.2 2D-Nanosheets

A chapter presents the 2D-nanosheets based hybrid NMs interaction with the plants. In addition, explains the different methods of synthesis of 2D-nanosheets. Then, emphasizes the interaction of 2D-nanosheets with plants. Moreover, it also highlights to penetration of 2D-nanosheets the seed coats, translocation in the plants and effects on plant growth and development (Lee et al. 2021).

1.5 Implications of Nanomaterials on Crop

The plant has different responses to treatment with NPs, whether morphology physiology and productivity (Hossain et al. 2020) (Fig. 1.3).

This book highlights the mechanism of NPs interaction in seed and various effects on seed germination and root growth, shoots, leaves, flowers and fruits. In addition, this book includes examples of many types of NPs. Also, their role to improve crop productivity by mentioning features of NPs in crop quality and quantity improvement (Rivero-Montejo et al. 2021).




Fig. 1.3 The impact of AgNPs on wheat seedling morphology during 35 days. A: treatment with AgNPs, B: control, D: day (Constructed by L.M. Alnaddaf)

1.5.1 Nutritional Value

The future goals of nanotechnology include high bioactive compound content of secondary metabolites in foods which advantages in improving the nutritional value of industrial crops and stress tolerance (Neme et al. 2021). Plants consider an essential component of the human diet via the supply to our body of vitamins, proteins, minerals, fiber, carbohydrates, lipids, and water (Rivero-Montejo et al. 2021).

The use of NMs has improved the nutritional value of fruits and vegetables without requiring increased consumption by affecting the biochemical and physiological properties of plants (Gomez et al. 2021).

1.5.2 Abiotic Stresses

The soil salinity, water's decrease, and heavy metal increase are the most dangerous to a plant's life cycle which knows as abiotic stresses (Gong et al. 2020).

Stresses adversely affect the plant growth and development via changes in the structural and chemical composition of the plant which lead decreasing in quality and quantity of production. Plants have developed various mechanisms to tolerate these challenges via transferring the stress signals within cells and between cells and tissues, sorting suitable chemical compounds for survival and reproduction, and continuing their growth and development (Rivero-Montejo et al. 2021).

Many reports indicated that when plants are treated with NPs their responses to stresses will be various via influence on biological and metabolism pathways. Also, their role to improve crop tolerance to abiotic stress for instance drought, salinity, and heavy metal stresses (Zohra et al. 2021).

1.5.3 Tissue Culture

A chapter is included to discuss the role of NMs in plant cell and tissue culture. It explains the impact of NMs on in vitro responses. Different NPs in tissue culture media could improve the callus induction, biomass and morphogenetic potential in explants (Barbasz et al. 2016).

NPs play a vital role in various biochemicals, physiological and anatomical routes of tissue culture. NPs improve regeneration, organogenesis, decontamination and secondary metabolite production to protect plant cells and tissues from biotic and abiotic stress. NPs have affected nutrient and protein levels and modulate the expression of certain genes encoded in certain proteins. In addition, it highlights the mechanism affecting callus, and quantitative and qualitative features of calli (Dallavalle et al. 2015).

In addition, it presents an overview of some important applications of NMs in plant tissue cultures such as somaclonal variation, organogenesis, somatic embryogenesis, disinfection, genetic fidelity, and regeneration (Devasia et al. 2020).

1.6 Conclusions and Prospects

This book provides an update on research and development in plant nanotechnology. It covers comprehensively various methods to synthesize NM and its characterization and applications. Moreover, explains the interaction of the NPs with plant cellular mechanisms and macromolecules. The initial phase is interacting NPs with the plant surface lead to adsorption it from the root and penetrate cell wall to move in plants. The second phase begins from the series of different effects at various levels, such as

the molecular, biochemical, physiological, morphological and productivity, which it reflects stimulating or inhibiting on the growth and development of the plant.

In addition, the book highlights the implications of NPs in different stages of plant growth and their effect on decreasing or increasing the quality and quantity of production as well as application in tolerating various stresses. Moreover, the book presents the role of NPs in tissue culture and their impacts on the callus physiology, biomass of explants and secondary metabolites according to the type of NPs and their concentrations.

Hence, future research needs to understand the mechanical complexity of interactions NPs with the plant (uptake, translocate, and accumulation) in different parts of a plant. Also discussed in details their effect of different NPs on growth stages in various plant species at the cellular levels.

References

- Aguirre-Becerra H, García-Trejo JF, Vázquez-Hernández C, Alvarado AM et al (2020) Effect of extended photoperiod with a fixed mixture of light wavelengths on tomato seedlings. *HortScience* 1(aop):1–8
- Alnaddaf LM, Almuhamady AK, Salem KFM, Alloosh MT et al (2021) Green synthesis of nanoparticles using different plant extracts and their characterizations. In: Al-Khayri JM, Ansari MI, Singh Ak (eds) *Nanobiotechnology: mitigation of abiotic stress in plants*. Springer, Cham, pp 165–199. https://doi.org/10.1007/978-3-030-73606-4_8
- Anjum NA, Sofo A, Scopa A, Roychoudhury A et al (2015) Lipids and proteins—major targets of oxidative modifications in abiotic stressed plants. *Environ Sci Pollut Res Int* 22(6):4099–4121. <https://doi.org/10.1007/s11356-014-3917-1>
- Barbasz A, Kreczmer B, Oćwieja M (2016) Effects of exposure of callus cells of two wheat varieties to silver nanoparticles and silver salt (AgNO₃). *Acta Physiol Plant* 38(3):76. <https://doi.org/10.1007/s11738-016-2092-z>
- Chen B, Liu M, Zhan L, Li C, Huang C (2018) Terbium(III) modified fluorescent carbon dots for highly selective and sensitive ratiometry of stringent. *Anal Chem* 90(6):4003–4009. <https://doi.org/10.1021/acs.analchem.7b05149>
- Dallavalle M, Calvaresi M, Bottoni A et al (2015) Graphene can wreak havoc with cell membranes. *ACS Appl Mater Interfaces* 7(7):4406–4414. <https://doi.org/10.1021/am508938u>
- Devasia J, Muniswamy B, Mishra MK (2020) Investigation of ZnO nanoparticles on in vitro cultures of Coffee (*Coffea Arabica* L.). *Int J Nanosci Nanotechnol* 16(4):271–277. http://www.ijnnonline.net/article_47981_75f4eb2f559670d0c8bc469675302389.pdf
- El-Shetehy M, Moradi A, Maceroni M, Reinhardt D, Petri-Fink A, Rothen-Rutishauser B et al (2020) Silica NPs enhance disease resistance in *Arabidopsis* plants. *Nat Nanotechnol* 16(3):344–353. <https://doi.org/10.1038/s41565-020-00812-0>
- Fedorenko AG, Minkina TM, Chernikova NP, Fedorenko GM (2020) The toxic effect of CuO of different dispersion degrees on the structure and ultrastructure of spring barley cells (*Hordeum sativum distichum*). *Environ Geochem Health*. <https://doi.org/10.1007/s10653-020-00530-5>
- Gaspar T, Kevers C, Penel C et al (1996) Plant hormones and plant growth regulators in plant tissue culture. *In Vitro Cell Dev Biol- Plant* 32(4):272–289
- Gomez A, Narayan M, Zhao L, Jia X (2021) Effects of nano-enabled agricultural strategies on food quality: current knowledge and future research needs. *J Hazard Mater* 401:123385. <https://doi.org/10.1016/j.jhazmat.2020.123385>

- Gong Z, Xiong L, Shi H, Yang S (2020) Plant abiotic stress response and nutrient use efficiency. *Sci China Life Sci.* 63:635–674. <https://doi.org/10.1007/s11427-020-1683-x>
- Hasan MK, Cheng Y, Kanwar MK, Chu XY (2017) Responses of plant proteins to heavy metal stress—a review. *Front Plant Sci* 1–16. <https://doi.org/10.3389/fpls.2017.01492>
- Hossain Z, Mustafa G, Sakata K, Komatsu S (2016) Insights into the proteomic response of soybean towards Al₂O₃, ZnO, and Ag nanoparticles stress. *J Hazard Mater* 304:291–305. <https://doi.org/10.1016/j.jhazmat.2015.10.071>
- Hossain Z, Yasmeen F, Komatsu S (2020) Nanoparticles: synthesis, morphophysiological effects, and proteomic responses of crop plants. *Int J Mol Sci* pp 21: 3056–3064 doi:<https://doi.org/10.3390/ijms21093056>
- Kataria S, Jain M, Rastogi A, Živčák M et al (2019) Role of nanoparticles on photosynthesis: avenues and applications. In: *Nanomaterials in plants, algae and microorganisms: concepts and controversies*, vol 2. Academic Press, pp 103–127. ISBN 978-0-12-811488-9. <https://doi.org/10.1016/C2016-0-00175-4>
- Khalil AT, Ovais M, Ullah I, Ali M et al (2020) Physical properties, biological applications and biocompatibility studies on biosynthesized single phase cobalt oxide (Co₃O₄) nanoparticles via *Sageretia thea* (Osbeck.). *Arab J Chem* 13(1):606–619
- Khan I, Raza MA, Khalid MH, Awan SA et al (2019). Physiological and biochemical responses of Pearl Millet (*Pennisetum glaucum* L.) seedlings exposed to silver nitrate (AgNO₃) and silver nanoparticles (AgNPs). *Int J Environ Res Publ Health* 16(13). <https://doi.org/10.3390/ijerph16132261>
- Khan S, Mansoor S, Rafi Z, Kumari B, Shoaib A, Saeed M et al (2021) A review on nanotechnology: properties, applications, and mechanistic insights of cellular uptake mechanisms. *J Mol Liq* 118008
- Khan SA (2020). Metal nanoparticles toxicity: role of physicochemical aspects. In Shah MR, Imran M, Ullah S (eds) *Metal nanoparticles for drug delivery and diagnostic applications*. Elsevier, pp 1–11. <https://www.sciencedirect.com/science/article/pii/B978012816960500001X>
- Kolenčík M, Lucia NemčekMartin Šebesta et al (2021) Effect of TiO₂ as plant growth-stimulating nanomaterial on crop production. In: Singh VP, Singh S, Tripathi DK, Prasad SM, Chauhan DK (eds) *Plant responses to nanomaterials, Nanotechnology in the life sciences*. Springer, Cham, pp 129–144. https://doi.org/10.1007/978-3-030-36740-4_5
- Lee JY, Kim MJ, Chung H (2021) Effects of graphene oxide on germination and early growth of plants. *J Nanosci Nanotechnol* 21(10):5282–5288. <https://doi.org/10.7717/Fpeerj.8387>
- Ma YJ, Xia J, Wang Y, Wang JW (2020) Stimulation of tanshinone production in *Salvia miltiorrhiza* hairy roots by β-cyclodextrin-coated silver nanoparticles. *Sustain Chem Pharm* 100271. <https://doi.org/10.1016/j.scp.2020.100271>
- Milewska-Hendel A, Chmura D, Wyrwał K, Kurczyn´ska EU, (2021) Cell wall epitopes in grasses of different novel ecosystem habitats on post-industrial sites. *Land Degrad Dev* 32:1680–1694
- Milewska-Hendel A, Zubko M, Karcz J, Stró'z D, et al (2017) Fate of neutral charged gold nanoparticles in the roots of the *Hordeum vulgare* L. cultivar Karat. *Sci Rep* 7:1–13
- Minkina T, Rajput V, Fedorenko G, Fedorenko A et al (2020) Anatomical and ultrastructural responses of *Hordeum sativum* to the soil spiked by copper. *Environ Geochem Health* 42:45–58
- Neme K, Nafady A, Uddin S, Tola YB (2021) Application of nanotechnology in agriculture, postharvest loss reduction and food processing: food security implication and challenges. *Heliyon* 7(12):e08539. <https://doi.org/10.1016/j.heliyon.2021.e08539>
- Nguyen DV, Nguyen HM, Le NT, Nguyen KH et al (2021) Copper nanoparticle application enhances plant growth and grain yield in maize under drought stress conditions. *J Plant Growth Regulation*. <https://doi.org/10.1007/s00344-021-10301-w>
- Pérez-de-Luque A (2017) Interaction of nanomaterials with plants: what do we need for real applications in agriculture? *Front Environ Sci* 5:12. <https://doi.org/10.3389/fenvs.2017.00012>
- Poddar K, Sarkar D, Sarkar A (2020) Nanoparticles on photosynthesis of plants: effects and role. In: Patra J, Fraceto L, Das G, Campos E (eds) *Green nanoparticles. Nanotechnology in the life sciences*. Springer, Cham. https://doi.org/10.1007/978-3-030-39246-8_13

- Rajput V, Minkina T, Fedorenko A, Sushkova S et al (2018) Toxicity of copper oxide nanoparticles on spring barley (*Hordeum sativum* distichum). *Sci Total Environ* 645:1103–1113
- Rao S, Shekhawat GS (2016) Phytotoxicity and oxidative stress perspective of two selected nanoparticles in *Brassica juncea*. *Biotech* 3(6):244. <https://doi.org/10.1007/s13205-016-0550-3>
- Rivero-Montejo SdJ, Vargas-Hernandez M, TorresPacheco I (2021). Nanoparticles as novel elicitors to improve bioactive compounds in plants. *Agriculture* 11:134. <https://doi.org/10.3390/agriculture11020134>
- Sáez PL, Bravo LA, Cavieres LA, Vallejos V (2017) Photosynthetic limitations in two Antarctic vascular plants: importance of leaf anatomical traits and Rubisco kinetic parameters. *J Exp Bot* 68:2871–2883. <https://doi.org/10.1093/jxb/erx148>
- Santiago EF, Pontes MS, Arruda GJ et al (2020) understanding the interaction of nanopesticides with plants. In: Fraceto LF, de Castro VL, Grillo R, Avila D, Caixeta Oliveira H, Lima R (eds) *Nanopesticides—from research and development to mechanisms of action and sustainable use in agriculture*. Springer, Cham, Switzerland, pp 69–109
- Schwab F, Zhai G, Kern M et al (2016) Barriers, pathways and processes for uptake, translocation and accumulation of nanomaterials in plants—critical review. *Nanotoxicol* 10(3):257–278. <https://doi.org/10.3109/17435390.2015.1048326>
- Shang Y, Hasan K, Ahammed GJ (2019) Applications of nanotechnology in plant growth and crop protection: a review. *Molecules* 24:2558. <https://doi.org/10.3390/molecules24142558>
- Shekhawat GS, Mahawar L, Rajput P et al (2021) Role of engineered carbon nanoparticles (CNPs) in promoting growth and metabolism of *Vigna radiata* (L.) Wilczek: Insights into the biochemical and physiological responses. *Plants* 10(7):1317. <https://doi.org/10.3390/plants10071317>
- Swift TA, Oliver TAA, Galan MC, Whitney HM (2019) Functional nanomaterials to augment photosynthesis: evidence and considerations for their responsible use in agricultural applications. *J R Soc Interface Focus* 9(1):20180048
- Tan X, Lin C, Fugetsu B (2009) Studies on toxicity of multi-walled carbon nanotubes on suspension rice cells. *Carbon* 47(15):3479–3487. <https://doi.org/10.1016/j.carbon.2009.08.018>
- Tombuloglu H, Slimani Y, AlShammari TM et al (2020) Uptake, translocation, and physiological effects of hematite (α -Fe₂O₃) nanoparticles in barley (*Hordeum vulgare* L.). *Environ Pollut* 266:115391. <https://doi.org/10.1016/j.envpol.2020.115391>
- Tripathi D, Rai KK, Rai SP (2020) Impact of green synthesized WcAgNPs on in-vitro plant regeneration and withanolides production by inducing key biosynthetic genes in *Withania coagulans*. *Plant Cell Rep* 40:283–299. <https://doi.org/10.1007/s00299-020-02630-z>
- Wang B, Guan C, Fu Q (2021) The traditional uses, secondary metabolites, and pharmacology of *Lycopodium* species. *Phytochem Rev*. <https://doi.org/10.1007/s11101-021-09746-4>
- Weyers J, Paterson N (2001) Plant hormones and the control of physiological processes. *New Phytol* 152(3):375–407. <https://doi.org/10.1046/j.0028-646x.2001.00281.x>
- Yang J, Cao W, Rui Y (2017) Interactions between NPs and plants: phytotoxicity and defense mechanisms. *J Plant Interact* 12(1):158–169. <https://doi.org/10.1080/17429145.2017.1310944>
- Zahedi SM, Hosseini MS, Daneshvar Hakimi Meybodi N, Peijnenburg W (2021) Mitigation of the effect of drought on growth and yield of pomegranates by foliar spraying of different sizes of selenium nanoparticles. *J Sci Food Agric* 101(12):5202–5213. <https://doi.org/10.1002/jsfa.11167>
- Zhang C, Cha R, Zhang P et al (2022) Cellulosic substrate materials with multi-scale building blocks: fabrications, properties and applications in bioelectronic devices. *Chem Eng J* 430:132562. <https://doi.org/10.1016/j.cej.2021.132562>
- Zohra E, Ikram M, Omar AA, Hussain M et al (2021) Potential applications of biogenic selenium nanoparticles in alleviating biotic and abiotic stresses in plants: a comprehensive insight on the mechanistic approach and future perspectives. *Green Process and Synth* 10:456–475. <https://doi.org/10.1515/gps-0047>

Part I
Cellular Mechanisms