

Breastfeeding and Metabolic Programming

Özlem Naciye Şahin
Despina D. Briana
Gian Carlo Di Renzo
Editors

Breastfeeding and Metabolic Programming

Özlem Naciye Şahin • Despina D. Briana
Gian Carlo Di Renzo
Editors

Breastfeeding and Metabolic Programming

 Springer

Editors

Özlem Naciye Şahin
Department of Child Health and Diseases
Arel University, Medical Faculty
Istanbul, Türkiye

Despina D. Briana
Pediatrics Department
National and Kapodistrian University of
Athens, Greece

Gian Carlo Di Renzo
Department of Obstetrics and Gynecology
University of Perugia
Perugia, Perugia, Italy

ISBN 978-3-031-33277-7 ISBN 978-3-031-33278-4 (eBook)
<https://doi.org/10.1007/978-3-031-33278-4>

© The Editor(s) (if applicable) and The Author(s), under exclusive license to Springer Nature Switzerland AG 2023

This work is subject to copyright. All rights are solely and exclusively licensed by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors, and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

Preface

It is very well known that breastmilk is the healthiest option for newborns as the source of nutrients, improves immunity, and for many other known and unknown mechanisms.

Metabolic programming stands at a not well-known site. Long-term health is affected by prenatal and early postnatal environments, the developmental origins of health and illness theory suggests. Growing data from human studies show that insufficient and excessive nutrition during pregnancy can alter gene expression and have lasting effects on child phenotypes through epigenetic alterations. In light of the possibility of its application in the programming of both health and illness, epigenetics has garnered considerable interest. Little is understood, however, about the processes through which early postnatal diet can influence long-term health.

Recent research shows that several common prenatal exposures, such as intra-uterine growth restriction, extremes of birth weight, maternal obesity, and maternal diabetes, are linked to increased fat mass, decreased muscle mass, and decreased bone density, with effects reported throughout infancy, childhood, and middle age. Genetic and epigenetic factors, stem cell commitment and function, mitochondrial metabolism, and maternal food are mechanisms and mediators. Changes in body composition are a typical phenotype after exposure to adverse conditions during pregnancy, and they may play a role in the prenatal programming of obesity and diabetes risk.

This book synthesizes the findings of several fields of studies into a single, accessible volume, focusing on the central idea of breastmilk and metabolic programming.

Initially, I would like to express my most heartfelt thanks to my father, Orhan Rami Atan, a well-known geology professor, who made me love science. My father told me about the importance of biological differentiation, that is, evolutionary processes, in the adaptation process of living things in the name of survival, by witnessing the changes of the same species over millennia and the fossils of geological periods he showed me in childhood. My father discovered and analyzed many fault lines, including the northern Anatolian fault line, sometimes on foot in the Taurus

Mountains, sometimes on horseback, and showed them to us by transferring them to geology education books.

I offer my sincerest appreciation to Gian Carlo Di Renzo and Despina D. Briana, who were accepted to be co-authors with me, to Cemal Cingi and Nuray Bayar Muluk, who planned and edited the “Breastfeeding: Special Conditions and Issues” section.

Last but not least, my appreciation is to Ayse Engin Arisoy, a leader or even a hero with her intelligence, wisdom, and knowledge in the newborn era of our country, and Emin Sami Arisoy, who is a hard worker but always stays calm. They guided me in planning the contents of the book.

This book will be helpful for readers and set the first step for other papers and books in this field.

Istanbul, Türkiye
Athens, Greece
Perugia, Perugia, Italy
March 2, 2023

Özlem Naciye Şahin
Despina D. Briana
Gian Carlo Di Renzo

Contents

Part I Overview

Human Milk Composition: Nutrients and Bioactive Factors	3
Senem Alkan Özdemir, Özlem Naciye Şahin, and Despina D. Briana	
Perinatal Maternal Nutrition and Breast Milk Composition	17
Vesim Bekraki, Gian Carlo Di Renzo, and Ayşe Engin Arısoy	
Immunological Aspects of Breast Milk	29
Hilal Güngör and Sinan Tüfekçi	
Immunological and Anti-Infectious Benefits of Breastfeeding	39
Mustafa Törehan Aslan, Ayşe Engin Arısoy, and Armando G. Correa	
Physiological Aspects of Lactation	51
Nevin Cambaz Kurt and Gian Carlo Di Renzo	
Breast Milk Oligosaccharides	61
Özlem Naciye Şahin, Aysel Özpınar, and Despina D. Briana	
Breast Milk as a Biological System	73
Ayten Guner Atayoglu	
Microbiota Composition of Breast Milk	83
Şeyhan Erişir Oygucu and Özlem Bekem	
Renal Physiology of Pregnancy	99
Fatih Palit	

Part II Metabolic Programming

The Evolution and Genomic Aspects of Milk	111
Özlem Naciye Şahin and Gian Carlo Di Renzo	
Programming Molecules in Early Life	121
Özlem Naciye Şahin and Despina D. Briana	

The Part Breast Milk Plays in Epigenetic Programming	131
Özlem Naciye Şahin	
The Metabolome of Breast Milk and Its Potential Long-Term Effects on the Child	149
Özlem Naciye Şahin, Despina D. Briana, and Gian Carlo Di Renzo	
The Infant–Mother Molecular Conversation Involving Breast Milk mRNA	159
Özlem Naciye Şahin and Despina D. Briana	
Genetic Causes of Obesity and Bioactive Substances	173
Özlem Naciye Şahin, Samim Ozen, and Despina D. Briana	
Epigenetic Causes of Obesity	191
Özlem Naciye Şahin, Samim Ozen, and Despina D. Briana	
Gestational Diabetes and Variety in the Composition of Breast Milk	201
Özlem Naciye Şahin and Gian Carlo Di Renzo	
Maternal PUFA Supplementation and Epigenetic Influences on Fat Tissue	213
Özlem Naciye Şahin and Aysel Özpınar	
Breast Milk Proteases	219
Özlem Naciye Şahin and Despina D. Briana	
Breast Milk and Leptin Resistance	227
Özlem Naciye Şahin and Muhittin Abdülkadir Serdar	
The Effects of Metabolic Alteration on Embryonic Stem Cells	235
Özlem Naciye Şahin and Muhittin Abdülkadir Serdar	
Part III Breastfeeding and Neuro-developmental Programming	
Neural Maturation of Breastfed Infants	243
Özge Serçe Pehlevan, Bülent Kara, and Despina D. Briana	
Breastfeeding and Motor Development in Preterm and Term Infants	267
Gülten Öztürk, Bülent Kara, and Gian Carlo Di Renzo	
Breastfeeding, Intelligence, and Social-Language Development	281
İpek Dokurel Çetin and Bülent Kara	
Breastfeeding in Immune-Mediated Demyelinating Disorders of the Central Nervous System	299
Tuğçe Damla Dilek, Sema Saltık, and Bülent Kara	

Part IV Breastfeeding and Infections

Role of Breastfeeding in the Prevention of Infectious Diseases 317

Funda Çipe, Ayşe Engin Arısoy, Emin Sami Arısoy,
and Sheldon L. Kaplan

Vaccination and Breastfeeding. 341

Gonca Keskindemirci and Gülbin Gökçay

Breastfeeding and Maternal Bacterial Infections 355

Özge Kaba, Ayşe Engin Arısoy, Emin Sami Arısoy,
and Sheldon L. Kaplan

Breastfeeding and Maternal Viral Infections 377

Murat Sütçü, Funda Yıldız, and Fatma Levent

Breastfeeding and Maternal Fungal Infections 389

Derin Oygur and Despina D. Briana

Breastfeeding and Maternal Parasitic Infections. 399

Yıldız Ekemen Keleş, Ayşe Engin Arısoy, Emin Sami Arısoy,
and Armando G. Correa

Maternal Infections: Who Can and Who Cannot Breastfeed? 415

Gonca Keskindemirci and Gülbin Gökçay

Part V Breastfeeding: Special Conditions and Issues

Programming the Gustatory and Olfactory Systems 433

Fatih Yucedag, Cemal Cingi, and Sergei Karpischenko

Abnormalities of Smell and Taste in Eating Behaviour Through Life 441

Muhammet Pamukcu, Cemal Cingi, and Glenis Scadding

Olfactory Influences on Human Feeding Behaviour, from Sense to Satiety. 455

Soner Taşar, Cemal Cingi, and Dilyana Vicheva

Choanal Atresia 465

Rezzan Budak, Cemal Cingi, and Dennis Chua

Nasal Toilet in Infants. 475

Ayşe Karaoğullarından, Nuray Bayar Muluk, and Luisa Maria Bellussi

Newborn Nasal Congestion 483

Tarik Yagci, Riza Dundar, and Nuray Bayar Muluk

Newborn Epistaxis	491
Aykut Ceyhan, Nuray Bayar Muluk, and Desiderio Passali	
Could Breastfeeding be a Protective Factor for Sinusitis?	499
Nagehan Dilsad Erdogdu Küçükcan, Cemal Cingi, and Codrut Sarafoleanu	
Could Breastfeeding Be a Protective Factor for Sleep Apnea?	509
Saltuk Bugra Kilinc, Nuray Bayar Muluk, and Suela Sallavaci	
The Role of Breastfeeding on the Development and Prevention of Allergic Diseases	519
Özlem Sancaklı, Demet Can, and Hesham Negm	
Breastfeeding and Atopic Dermatitis	531
Sait Karaman and Demet Can	
Congenital Abnormalities of the Airway	539
Elad Azizli, Cemal Cingi, and Jeffrey C. Bedrosian	
Upper Airway Lesions in the Neonate	551
Mustafa Caner Kesimli	
Newborn Hearing Screening	561
Taylan Bilici, Nuray Bayar Muluk, and Yusuf Dundar	
Hearing Loss in Neonates and Infants	575
Bilal Sizer, Nuray Bayar Muluk, and Nitin R. Ankle	
Programming the Auricle	587
Ergun Sevil, Cemal Cingi, and Mario Milkov	
The Effect of Breastfeeding on Childhood Otitis Media	595
Muhammet Pamukcu, Nuray Bayar Muluk, and Peter Catalano	
Candidiasis During Breastfeeding	603
Mehtap Koparal, Nuray Bayar Muluk, and Gabriela Kopacheva Barsova	
Frenotomy for Tongue-Tie in Neonates	611
Selis Gülseven Güven and Ahmet Köder	
Dietary Modifications and Supplementation in Breastfeeding Mothers	621
İlke Özer Aslan and Keziban Doğan	
Human Milk Banking: A Brief Overview	631
Güniz Yaşöz, Esra Polat, and Despina D. Briana	
Emotional Communication During Breastfeeding	639
Can Cemal Cingi and Dilek Eroğlu	
Tobacco Use in Breastfeeding Mothers	649
Ali Timucin Atayoglu and Ayten Guner Atayoglu	

Contributors

Nitin R. Ankle, MD Department of ENT & Head-Neck Surgery, KLE Academy of Higher Education & Research (KAHER), J. N. Medical College, Belagavi, Karnataka, India

Ayşe Engin Arısoy, MD Faculty of Medicine, Division of Neonatology, Department of Pediatrics, Kocaeli University (Emeritus), Kocaeli, Türkiye

Emin Sami Arısoy, MD Faculty of Medicine, Division of Pediatric Infectious Diseases, Department of Pediatrics, Kocaeli University (Emeritus), Kocaeli, Türkiye

İlke Özer Aslan, MD Faculty of Medicine, Department of Obstetrics and Gynecology, Tekirdag Namık Kemal University, Tekirdag, Türkiye

Mustafa Törehan Aslan, MD Division of Neonatology, Department of Pediatrics, Faculty of Medicine, Tekirdağ Namık Kemal University, Tekirdağ, Türkiye

Ali Timucin Atayoglu, MD Department of Family Medicine, Istanbul Medipol University, International School of Medicine, Istanbul, Türkiye

Ayten Guner Atayoglu, MD Department of Family Medicine, Beylikdüzü State Hospital, Istanbul, Türkiye

Elad Azizli, MD Otorhinolaryngology Specialist, Private Practice, Istanbul, Türkiye

Gabriela Kopacheva Barsova, MD, PhD Faculty of Medicine, Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

Jeffrey C. Bedrosian, MD Rhinology and Skull Base Surgery, Specialty Physician Associates, St. Luke's Medical Center, Bethlehem, PA, USA

Özlem Bekem, MD Division of Pediatric Gastroenterology, Hepatology and Nutrition, Department of Pediatrics, Health Sciences University, Dr Behcet Uz Children's Hospital, İzmir, Türkiye

Vesim Bekraki, MD Department of Pediatrics, Türkiye Hospital, İstanbul, Türkiye

Luisa Maria Bellussi, MD University of Siena- ENT Clinic, Siena, Italy

Taylan Bilici, MD Department of Otorhinolaryngology, Adana Seyhan State Hospital, Adana, Türkiye

Despina D. Briana, MD, PhD NICU, Third Department of Pediatrics, National and Kapodistrian University of Athens, University General Hospital “ATTIKON”, Athens, Greece

Rezzan Budak, MD Department of Otorhinolaryngology, Şehit Sait Ertürk State Hospital, Ankara, Türkiye

Demet Can, MD Division of Allergy and Immunology, Department of Pediatrics, Izmir Dr. Behcet Uz Children’s Hospital, Izmir, Türkiye

Peter Catalano, MD, FACS, FARS Department of Otolaryngology, Tufts University, School of Medicine, St. Elizabeth’s Medical Center, Boston, MA, USA

İpek Dokurel Çetin, MD Faculty of Medicine, Division of Pediatric Neurology, Department of Pediatrics, Balikesir University, Balikesir, Türkiye

Aykut Ceyhan, MD Department of Otorhinolaryngology, Kutahya Evliya Çelebi Training and Research Hospital, Kutahya, Türkiye

Dennis Chua, MD Department of Otorhinolaryngology, ENT Surgeons Medical Centre, Mount Elizabeth Hospital, Singapore, Singapore

Can Cemal Cingi, PhD Faculty of Communication Sciences, Department of Communication Design and Management, Anadolu University, Eskişehir, Türkiye

Cemal Cingi, MD Medical Faculty, Department of Otorhinolaryngology, Eskisehir Osmangazi University, Eskisehir, Türkiye

Funda Çipe, MD Department of Pediatric Allergy-Immunology, Al Qassimi Women's and Children’s Hospital, Sharjah, United Arab Emirates

Armando G. Correa, MD Division of Academic General Pediatrics, Department of Pediatrics, Baylor College of Medicine, and Section of International and Destination Medicine, Texas Children’s Hospital, Houston, TX, USA

Tuğçe Damla Dilek, MD Cerrahpasa Medical Faculty, Division of Pediatric Neurology, Department of Pediatrics, Istanbul University, Istanbul, Türkiye

Gian Carlo Di Renzo, MD, PhD Centre of Perinatal Medicine of the University of Perugia/Sechenov University, Moscow, Russia

Keziban Doğan, MD Department of Obstetrics and Gynecology, University of Health Sciences, Bakırköy Dr. Sadi Konuk Training and Research Hospital, İstanbul, Türkiye

Riza Dundar, MD Medical Faculty, Department of Otorhinolaryngology, Bilecik Seyb Edebali University, Bilecik, Türkiye

Yusuf Dundar, MD Department of Otolaryngology-Head and Neck Surgery, Texas Tech University, Health Sciences Center, Lubbock, TX, USA

Dilek Eroğlu, PhD School of Foreign Languages, Department of Foreign Languages, Anadolu University, Eskişehir, Türkiye

Gülbin Gökçay, MD Division of Social Pediatrics, Department of Pediatrics, İstanbul Faculty of Medicine, İstanbul University, İstanbul, Türkiye

Department of Social Pediatrics, Institute of Child Health, İstanbul University, İstanbul, Türkiye

Hilal Güngör, MD Department of Internal Medicine, Eskisehir City Hospital, Eskisehir, Türkiye

Selis Gülseven Güven, MD Medical Faculty, Department of Otorhinolaryngology, Trakya University, Edirne, Türkiye

Özge Kaba, MD İstanbul Faculty of Medicine, Division of Pediatric Infectious Diseases, Department of Pediatrics, İstanbul University, İstanbul, Türkiye

İstanbul Çam and Sakura Hospital, Section of Pediatric Infectious Diseases, University of Health Science, İstanbul, Türkiye

Sheldon L. Kaplan, MD Division of Infectious Diseases, Department of Pediatrics, Baylor College of Medicine, and Infectious Disease Service, Texas Children's Hospital, Houston, TX, USA

Bülent Kara, MD Faculty of Medicine, Division of Pediatric Neurology, Department of Pediatrics, Kocaeli University, Kocaeli, Türkiye

Sait Karaman, MD Division of Allergy and Immunology, Department of Pediatrics, Manisa City Hospital, Manisa, Türkiye

Ayşe Karaoğullarından Department of Otorhinolaryngology, Adana City Hospital, Adana, Türkiye

Sergei Karpischenko, MD Department of Otorhinolaryngology, The First Pavlov State Medical University of Saint Petersburg, St. Petersburg, Russia

Yıldız Ekemen Keleş, MD University of Health Sciences, Tepecik Training and Research Hospital, Section of Pediatric Infectious Diseases, İzmir, Türkiye

Mustafa Caner Kesimli, MD Caner Kesimli Private ENT Clinic, İstanbul, Türkiye

Gonca Keskindemirci, MD Division of Social Pediatrics, Department of Pediatrics, İstanbul Faculty of Medicine, İstanbul University, İstanbul, Türkiye

Saltuk Bugra Kilinc, MD Department of Otorhinolaryngology, Yozgat Sarıkaya State Hospital, Yozgat, Türkiye

Ahmet Köder, MD Medical Faculty, Department of Otorhinolaryngology, Trakya University, Edirne, Türkiye

Mehtap Koparal, MD Department of Otorhinolaryngology, Health Sciences University, Adıyaman Training and Research Hospital, Adıyaman, Türkiye

Nagehan Dilsad Erdogdu Küçükcan, MD Department of Otorhinolaryngology, Çukurova State Hospital, Adana, Türkiye

Nevin Cambaz Kurt, MD Department of Pediatrics, Başakşehir Çam and Sakura City Hospital, İstanbul, Türkiye

Fatma Levent, MD Division of Pediatric Infectious Diseases, Department of Pediatrics, School of Medicine, Texas Tech University, Lubbock, TX, USA

Mario Milkov, MD Faculty of Medicine, Department of Otorhinolaryngology, Varna University, Varna, Bulgaria

Nuray Bayar Muluk, MD Medical Faculty, Department of Otorhinolaryngology, Kirikkale University, Kirikkale, Türkiye

Hesham Negm, MD Faculty of Medicine, Department of Otorhinolaryngology, Cairo University, Cairo, Egypt

Derin Oygur, MD Department of Pediatric Infectious Diseases, Neurology Department, Dr. Burhan Nalbantoğlu State Hospital, Nicosia, Northern Cyprus

Seyhan Erişir Oygucu, MD Division of Neonatology, Department of Pediatrics, University of Kyrenia, Kyrenia, Cyprus

Senem Alkan Özdemir, MD Department of Neonatology, Behçet Uz Children's Hospital, Health Sciences University, İzmir, Türkiye

Samim Ozen, MD Medical Faculty, Pediatric Endocrinology Division, Department of Pediatrics, Ege University, İzmir, Türkiye

Aysel Özpınar, MD Medical Faculty, Department of Basic Sciences, Medical Biochemistry, Acibadem Mehmet Ali Aydınlar University, İstanbul, Türkiye

Gülten Öztürk, MD Division of Pediatric Neurology, Department of Pediatrics, Marmara University Pendik Education and Research Hospital, İstanbul, Türkiye

Fatih Palit, MD Department of Nephrology, Basakşehir Çam and Sakura City Training and Research Hospital, İstanbul, Türkiye

Muhammet Pamukcu, MD Medical Faculty, Department of Otorhinolaryngology, İstanbul Altınbas University, İstanbul, Türkiye

Desiderio Passali, MD, PhD Department of Medical, Surgical and Neuroscience Sciences, University of Siena, Siena, Italy

Department of Otorhinolaryngology, University of Siena, Siena, Italy

Özge Serçe Pehlevan, MD Faculty of Medicine, Division of Neonatology, Department of Pediatrics, Kocaeli University, Kocaeli, Türkiye

Esra Polat, MD Division of Pediatric Gastroenterology, Department of Pediatrics, Health Sciences University, Umraniye Education and Research Hospital, Istanbul, Türkiye

Özlem Naciye Şahin, MD, PhD Department of Child Health and Diseases, Arel University, Medical Faculty, Istanbul, Türkiye

Suela Sallavaci, MD, MSc, PhD Department of Otorhinolaryngology, University Hospital Centre “Mother Teresa”, Tirana, Albania

Sema Saltık, MD Cerrahpasa Medical Faculty, Division of Pediatric Neurology, Department of Pediatrics, Istanbul University, Istanbul, Türkiye

Özlem Sancaklı, MD Division of Allergy and Immunology, Department of Pediatrics, Izmir Dr. Behcet Uz Children’s Hospital, Izmir, Türkiye

Codrut Sarafoleanu, MD Sfanta Maria Clinical Hospital ENT&HNS Department, University of Medicine and Pharmacy “Carol Davila”, Bucharest, Romania

Glenis Scadding, MD, FRCP University College Hospital London, RNENT Hospital, London, UK

Muhittin Abdülkadir Serdar, MD Medical Faculty, Department of Basic Sciences, Medical Biochemistry, Acibadem Mehmet Ali Aydınlar University, Istanbul, Türkiye

Ergun Sevil, MD Department of Otorhinolaryngology, Alaaddin Keykubat University, Alanya Training and Research Hospital, Alanya, Antalya, Türkiye

Bilal Sizer, MD Department of Otorhinolaryngology, Diyarbakir Memorial Hospital, Diyarbakir, Türkiye

Murat Sütçü, MD Division of Pediatric Infectious Diseases, Department of Pediatrics, Faculty of Medicine, Istinye University İstanbul, İstanbul, Türkiye

Soner Taşar, MD Department of Otorhinolaryngology, Afyonkarahisar State Hospital, Afyonkarahisar, Türkiye

Sinan Tüfekçi, MD Medical Faculty, Neonatology Section, Department of Pediatrics, Tekirdağ Namık Kemal University, Tekirdağ, Türkiye

Dilyana Vicheva, MD, PhD Department of Otorhinolaryngology, Medical University of Plovdiv, Plovdiv, Bulgaria

Tarık Yagci, MD Medical Faculty, Department of Otorhinolaryngology, Bilecik Seyb Edebali University, Bilecik, Türkiye

Güniz Yaşöz, MD Department of Pediatrics, University of Kyrenia, Kyrenia, Northern Cyprus

Funda Yıldız, MD Faculty of Medicine, Department of Pediatrics, Istinye University İstanbul, Zeytinburnu/İstanbul, Türkiye

Fatih Yucedag, MD Faculty of Medicine, Department of Otorhinolaryngology, Karamanoglu Mehmetbey University, Karaman, Türkiye

Part I

Overview

Human Milk Composition: Nutrients and Bioactive Factors



Senem Alkan Özdemir, Özlem Naciye Şahin, and Despina D. Briana

1 Introduction

It is widely accepted that the ideal way for a human infant to be nourished up to the age of 6 months is exclusive breastfeeding. Thereafter, breastfeeding with additional sources of nutrition is suitable up to the first or second birthday, or for even more prolonged periods [1, 2]. Breast milk is the sole food which precisely addresses the developmental requirements of the infant, providing both nutritional elements and bioactive compounds to ensure survival as well as healthy growth [3]. In this chapter, the nutritional and bioactive elements of breast milk are summarised. The latter category includes cells, molecules which target infectious pathogens and dampen down inflammation, growth factors and prebiotics. Breast milk differs from artificial substitutes insofar as the latter is of essentially fixed composition, whereas breast milk varies considerably in its constituents, whether in a single nursing session, over the course of the day or over the whole period of lactation. Human milk composition differs from mother to mother, as it does between populations. There are a number of reasons for this variable composition, including factors related to both the mother and the surrounding environment, as well as to the way milk is expressed and the feeding pattern. Knowledge of the constituents of breast milk is vital, in order to best utilise the opportunities breastfeeding offers, especially for those children who are most vulnerable to adverse outcomes, and to inform

S. A. Özdemir (✉)

Department of Neonatology, Behçet Uz Children's Hospital, Health Sciences University, İzmir, Türkiye

Ö. N. Şahin

Department of Child Health and Diseases, Arel University, Medical Faculty, Istanbul, Türkiye

D. D. Briana

NICU, Third Department of Pediatrics, National and Kapodistrian University of Athens, University General Hospital "ATTIKON", Athens, Greece

discussion about how breast milk can potentially be stored and whether it should be pasteurised. The chapter will also discuss bioactive factors which have been identified as potential prophylactic or therapeutic agents and may be suitable to undergo pharmaceutical development and testing [3–5].

2 Stages of Lactation

The initial milk expressed by the mother postpartum is colostrum, which differs from mature milk in the amount expressed, its appearance and constituents. This early milk is expressed in low volumes over the initial days after delivery and contains abundant immune-related cells and humoral factors, notably secretory IgA, lactoferrin and white blood cells. It also contains factors that direct development, such as epidermal growth factor (EGF) [4–6]. The level of lactose in colostrum is low and it appears that this fluid is optimised for immune support and directing growth more than for supplying nutrition. The electrolyte composition of colostrum also differs from transitional or mature milk, being richer in sodium, chloride and magnesium ions, but lower in potassium and calcium [5, 6]. The rising of milk lactose concentration heralds the onset of transitional milk. Within the breast, this reflects a structural change, with the epithelium developing tight junctions between cells. The exact point at which this change occurs varies from mother to mother, however the expression of transitional milk, known as the second stage of lactogenesis, generally begins a few days after delivery. The second stage is considered delayed if it has not started by 72 h postpartum. This delay seems to be associated with neonatal prematurity and maternal obesity. It also correlates with specific metabolic biomarkers [7, 8]. The changes in biochemical composition that confirm the onset of the second stage include sodium concentration, sodium to potassium ratio, as well as citrate and lactose levels [9].

Transitional milk somewhat resembles colostrum, although it involves a considerable step up in milk production as called for by the infant's rapid development and growth, which impose greater metabolic demands. The usual period during which transitional milk is expressed is from 5 days to 2 weeks after delivery. It is followed by the stage of mature milk. Mature milk contains all the expected elements at around 4–6 weeks after birth. Thereafter, there are only relatively minor adjustments to the constituents contained in breast milk, although compositional variety is always a feature of human milk [10, 11].

3 Studies of Breast Milk Composition

Studies concerning the composition of breast milk, as indexed in Medline, have steadily increased in number over the last half-century. There are still announcements of newly discovered constituents in human milk, and an active global research

effort is underway to determine the functions of each constituent. There have been multiple studies examining the composition of breast milk in different groups of mothers. These studies employed a range of methods to collect, store and analyse human milk. The ideal method of sampling is to obtain multiple samples of milk from the same mother at different points during a 24-h period [10, 11]. This sampling technique suffers from the disadvantage that it is costly and may discourage mothers from enrolling in a study. A second technique that allows for standardisation is to take all samples at a particular point during the day, such as in the morning, and to collect the entire milk content of whichever breast has not been used to feed the infant in the preceding 2–3 h. This procedure needs to be repeated several times in the same study participant on different days [12]. The majority of studies in the literature, nonetheless, are based on milk obtained in a non-standardised way from milk banks, where the milk arrives at different times and different points during nursing and is expressed from mothers at different stages of lactation. These studies on the composition of breast milk also lack standardisation in how and for how long milk was stored, how frequently was frozen and defrosted and whether it was pasteurised or not. All these factors may partly explain the heterogeneity in the results obtained.

4 Nutritional Constituents within Breast Milk

There are three sources for the nutritional components of breast milk. The lactocytes synthesise a number of compounds for secretion into milk. The maternal diet supplies other components, whilst others come from the body stores of the mother. In general, breast milk retains its nutritional value despite maternal dietary deficiencies, but there are specific vitamins and fatty acids which must be present in the maternal diet for these compounds to be present in breast milk. This issue is discussed elsewhere in this volume [12–14].

4.1 *Macronutrients*

The level of macronutrients in breast milk does exhibit a degree of variation between individuals and over the whole course of breastfeeding, but there is a notable consistency in macronutrient content in different groups, despite differences in maternal diets. On average, the concentration of protein in mature milk is calculated to lie between 0.9 and 1.2 g/dL, of lipids from 3.2 to 3.6 g/dL and of lactose from 6.7 to 7.8 g/dL. The calorific value of breast milk is estimated at between 65 and 70 kcal/dL and is mostly related to the lipid content. There are differences between milk expressed following a term pregnancy and that seen from mothers of preterm infants. Preterm milk generally contains a higher concentration of protein and lipids. A study undertaken in the Californian city of Davis examined mature milk

expressed 4 months after delivery and related its macronutrient composition to specific characteristics of the mother. There were a number of maternal characteristics found to be of significance, in particular body mass index (BMI), dietary protein, parity, whether menses had recommenced and how often the infant was fed. Individual mothers whose milk production was greater generally produced milk that was less rich in lipids and protein but richer in lactose.

Breast milk proteins can be separated into those that aggregate in the whey and those that are in the casein fraction. Each fraction contains a surprisingly varied range of different proteins and peptides [14, 15]. The proteins present at the highest concentration are casein, α -lactalbumin, lactoferrin, secretory immunoglobulin IgA, lysozyme and serum albumin [16, 17]. Around 25% of the nitrogen contained in breast milk is not in protein form, being found as urea, urate, creatine, creatinine, amino acids or nucleotides. There is a significantly higher level of protein in breast milk from mothers of preterm infants. Whatever the stage of pregnancy when delivery occurred, protein content falls in breast milk between the first and fourth to sixth weeks. If a preterm infant is fed by donor milk, the concentration of protein and amino acids is insufficient unless artificially supplemented. This is because most donor milk is of mature type and provided by mothers who delivered a term infant. The level of protein in breast milk does not depend on what the mother eats, but it does rise if the BMI is greater and falls when the volume of milk expressed is very high [10, 18].

The typical lipid composition of breast milk involves a high volume of palmitic acid and oleic acid residues attached to triacylglycerides. Palmitic acid is generally attached mid-chain to the glycerol moiety, whereas oleic acid is attached to either of the terminal carbons. The macronutrient category subject to the most variation in breast milk is lipid. The lipid content of hindmilk, the end portion of a feed, potentially contains three-fold more lipid than the initial foremilk [19]. One study which evaluated milk sampled from 71 women over the course of a full day reported that the lipid concentration was highest during the afternoon or evening and this finding was of statistical significance [20]. A different study correlated a quarter of the variation in fat contained in breast milk with variation in maternal protein consumption [10].

The aliphatic acid content of breast milk is affected by maternal dietary consumption, especially consumption of long chain polyunsaturated fatty acids (LCPUFAs). In Western countries, most fatty acids of this type in the diet are omega-6 aliphatic acids, with less consumption of omega-3 acids than is ideal. In North American mothers, breast milk is markedly deficient in docosahexaenoic acid; therefore lactating mothers whose diet lacks this nutrient should be advised to consume supplements to remedy the defect [18, 21, 22].

The main type of sugar within breast milk is lactose, a disaccharide. Lactose varies the least amongst the macronutrients found in breast milk, although it has been noted that women with a high milk volume do secrete greater concentrations of lactose into milk [10]. Oligosaccharides are also a major carbohydrate component in breast milk. They are typically present at a concentration of around 1 g/dL,

although this varies according to the stage in breastfeeding and the genetic composition of the mother [23–25]. Oligosaccharides in breast milk are not nutritive, but they perform other biological functions, as described later.

Figures representing the mean concentration of macronutrients in breast milk are helpful, but should not be allowed to give the misleading impression that individual samples of milk vary little in their composition. The opposite, in fact, applies, and this variation is especially apparent in lipid and protein concentrations. The total protein concentration in breast milk from mothers who delivered at term ranges from 0.6 to 1.4 g/dL. The range for lipids is from 1.8 to 8.9 g/dL, for lactose from 6.4 to 7.6 g/dL and the calorific value is between 50 and 115 kcal/dL [26]. There are significant differences between milk from mothers who delivered at term and those whose delivery was premature. Accordingly, when a nutritional plan for an infant at high risk is being formulated, the milk available needs to be analysed and fortified as necessary, whilst also monitoring effects on the infant's growth [27].

4.2 *Micronutrients*

Breast milk provides the standard to which artificial formula needs to approach to achieve optimal nutritional value. However, there is significant variation in the micronutrient content of breast milk as a result of variation in maternal diet and body stores. This especially applies to vitamins A, B₁, B₂, B₆, B₁₂, D, and iodine. Given the fact that mothers frequently consume a suboptimal diet, mothers can be advised to keep taking multivitamins whilst breastfeeding continues [28, 29]. The level of vitamin K in breast milk is always low, whatever the mother's dietary intake. This is the reasoning behind the American Academy of Paediatrics' recommendation that vitamin K should be administered intramuscularly to all neonates to prevent haemorrhagic disease of the newborn [28]. The level of vitamin D in breast milk is also low, especially if the mother has inadequate exposure to sunlight. Due to lifestyle changes, this is now frequently the case all across the globe [30]. Although benefit from supplementing the diet of breastfeeding mothers with vitamin D has not yet been proven experimentally, current guidelines advise supplying additional vitamin D to all new breastfeeding mothers and their offspring. The inability to provide a thorough, detailed review of every micronutrient contained in breast milk is a limitation of this chapter. Interested readers can find several detailed reviews available elsewhere in the literature [28, 29, 31].

5 **Bioactive Constituents and Their Sources**

One definition of bioactive constituents in the diet considers them as substances which 'affect biological processes or substrates which hence have an impact on body function or condition and ultimately health' [32, 33]. The bioactive molecules

found in breast milk originate from several different sources. The epithelial cells of breast produce and secrete certain such compounds, whilst cellular components of milk are the source of others. Furthermore, the breast epithelium recognises and actively transports some molecules from the plasma to the milk, by means of specific receptor interactions. The epithelial cells secrete milk fat globules (MFGs), which contain within them various proteins attached to membranes as well as specific lipid components [34]. These different sources all contribute to the abundance of bioactive molecules within breast milk. In women who are breastfeeding, B-lymphocytes are drawn towards the breast, and secretory immunoglobulin produced by these cells is then transferred into the milk ducts via polymeric immunoglobulin receptors (pIgR) [35]. In contrast, vascular endothelial growth factor (VEGF) concentrations are significantly higher in breast milk than in maternal circulation, which suggests that VEGF is manufactured and secreted by the breast itself [36, 37]. A knowledge of where the constituents of breast milk that possess bioactivity originate is helpful in understanding the varying levels of secreted maternal medications in milk.

There are a number of ways in which research into the bioactive breast milk compounds can impact clinical practice. A key concept that both patients and the wider public need to grasp is that the research so far clearly shows that breast milk is more sophisticated than just another type of food. Indeed, breast milk provides infants with a number of compounds which enhance both survival and healthy growth. Accordingly, for infants at risk and where the mother cannot supply her own milk, donor breast milk is vital, provided it is safe to use. Proteomic studies reveal that the types of proteins present differ according to the stage of breastfeeding, in addition to differences associated with whether delivery occurred at term or prematurely [14, 15]. Thus, wherever the possibility exists, donor milk should be from a donor whose own infant is at the same stage of development as the patient. In actual clinical practice, this precise matching may not be achievable. Recognition of the role of bioactive compounds in breast milk has led to a rethink in how donor milk is collected, stored and pasteurised. Retaining bioactivity has become the goal. Furthermore, the mechanisms by which bioactive milk constituents enhance infant health offer new targets for future pharmaceutical or other therapeutic interventions [36, 37].

This chapter does not aim to comprehensively review the entire spectrum of bioactivity in breast milk. Instead, the objective is to examine a subset of bioactive factors in breast milk and relate their changing levels to the developmental requirements of the infant. The patterns of secretion associated with term and premature delivery are described, as well as the variations associated with the stage of lactation. In many cases, the bioactive factors function in a synergistic manner, which means that replacement or supplementation of individual bioactive molecules or sets thereof is generally less efficacious than supplying complete breast milk [38].

6 Growth Factors

Multiple different growth factors have been identified in breast milk. These molecules act on various infant organs, such as the gut, circulatory, nervous and endocrine systems [36–38].

6.1 *Epidermal Growth Factor (EGF): Development and Repair of the Intestinal System*

EGF plays a vital role in healthy development and repair of the gut lining. It is detectable in breast milk as well as amniotic fluid [38–40]. EGF remains intact even in the highly acidic gastric interior, and is not degraded by digestive enzymes. Within the intestines it has a stimulatory effect upon the gut lining, promoting the synthesis of DNA and cellular replication. It also increases water and glucose uptake by the gut and promotes protein manufacture [41, 42]. EGF also performs several protective roles within the infant gut, such as preventing excessive apoptosis and opposing the action of tumour necrosis factor- α (TNF- α). The latter promotes inflammation and alters the tight junctions between different enterocytes and between hepatocytes [43]. Heparin-binding growth factor also belongs to the same family of related proteins. This is the principal signalling molecule involved in resolving hypoxic injury, ischaemic-reperfusion damage, shock secondary to haemorrhage, resuscitation injury and necrotising enterocolitis [44]. The highest concentration of EGF occurs at the beginning of the lactation period, decreasing as lactation continues [45, 46]. The mean concentration of EGF in colostrum exceeds the mother's circulating level by 2000 times. In mature milk, EGF is 100 times more abundant than in maternal circulation [41]. Additionally, the level in breast milk of mothers who delivered prematurely is higher than in cases where delivery occurred at term [45, 46].

6.2 *Neuronal Growth Factors: Normal Maturation*

The intestinal system of infants is immature in several different ways, including the development of the enteric nervous system, which depends for proper maturation on the presence of brain-derived neurotrophic factor (BDNF) and glial cell-line-derived neurotrophic factor (GDNF). Peristaltic intestinal motion is often dysfunctional in premature infants. This function improves in the presence of BDNF. Rodent models exhibit severe deficiency of neurones in the gut of animals without GDNF. Ciliary neurotrophic factor (CNF) is a protein with similarities to BDNF and GDNF. All three molecules are present in breast milk for a period lasting 90 days after delivery. When neurones were exposed to GDNF isolated from human milk, cellular survival improved and the cells developed new projections [30, 31].

6.3 The Insulin-like Growth Factor (IGF) Superfamily and Its Effects on Tissues

Breast milk contains several IGF-related proteins, notably IGF-I and II, as well as IGF-binding proteins and proteases with specificity for IGF. These molecules are most abundant in colostrum, but their levels steadily fall as lactation proceeds. Preterm and term milk do not importantly differ in their level of IGF apart from IGF-binding protein-2 being more abundant in preterm samples. Rodents exposed to surgical stress and fed by total parenteral nutrition exhibited higher levels of tissue growth and lower levels of gut atrophy when administered IGF-I. This effect has not been demonstrated in human subjects so far [1, 2]. IGF-I potentially also increases cellular survival when the gut is exposed to oxidative injury. Breastfeeding increases the concentration of IGF-I in the circulation. IGF absorbed through the gut retains its bioactivity and can enter the vascular circulation. Exactly how IGF taken up by this route functions so far remains unknown, however it has been shown that IGF-I administered by an enteral route at physiological concentrations increases red cell synthesis and raises the haematocrit [32, 33].

6.4 Vascular Endothelial Growth Factor (VEGF): Regulating Vascular Development

The development of new blood vessels is mostly regulated by the balance between VEGF and molecules which antagonise its effect. Colostrum contains the greatest level of VEGF amongst different types of breast milk. Term milk has a greater concentration of VEGF than preterm milk [34, 35]. The pathogenesis of retinopathy of prematurity is believed to arise from a combination of immaturity of the lungs, therapeutic oxygen administration and down-regulation of VEGF. This then causes the retina to grow blood vessels in a disorganised fashion [36, 37]. It is possible that breast milk, which contains VEGF, may be beneficial in mitigating the severity of this condition.

6.5 Erythropoietin: Promoting Development of the Gut and Preventing Anaemia

Erythropoietin is the principal hormone which stimulates erythrocyte synthesis. Haemorrhage, disease of the gut and under-developed haematopoiesis are factors contributing to the anaemia which may affect premature infants. This condition then severely harms normal growth and development [36–41]. It has been proposed that

exogenous administration of erythropoietin may be beneficial in the prevention of anaemia of prematurity, however results of this practice have produced unclear benefit [38]. Nonetheless, it has been demonstrated that where erythropoietin and iron are co-administered, a potential increase in haemoglobin and haematocrit occurs [39]. A trial involving low numbers of premature infants who were administered erythropoietin enterally found increased numbers of reticulocytes in the circulation. Furthermore, erythropoietin has significant trophic effects and leads to tighter integration of the enterocytic junctions. It appears that exogenous erythropoietin may partially prevent vertical transmission of HIV and may lower the incidence of necrotising enterocolitis [36–41].

6.6 Calcitonin and Somatostatin: Regulation of Growth

There are high concentrations of both calcitonin and the precursor molecule, procalcitonin, in breast milk [40]. The calcitonin receptor is present on cells of the intestines from the latter stages of pregnancy into the first year of life, as demonstrated by immunochemical methods. In normal circumstances, somatostatin undergoes swift degradation within the jejunum. It is not absorbed via the gut lining. When this hormone is delivered in breast milk, however, degradation does not occur and it remains bioactive in the intestines. The usual physiological role of somatostatin is inhibition of the action of growth factors. Its exact functions in breast milk is, however, still not fully elucidated [41, 42].

6.7 Adiponectin and Related Hormones: The Regulation of Metabolism and How the Body Is Composed

Adiponectin has a high molecular weight and possesses multiple endocrine functions, in particular the regulation of metabolism and inhibition of inflammatory responses. It is present at high concentration in breast milk and traverses the gut lining, after which it has an apparent regulatory function in metabolism in the infant. There is an inverse correlation between the concentration of breast milk adiponectin and both the infant body mass and BMI in exclusively breastfed infants. It has been hypothesised that adiponectin in breast milk may help to prevent individuals becoming obese or overweight at a later stage in life, but this has not yet been proven to occur. There are several other hormones with a regulatory role in metabolism that can be detected in breast milk. This group includes leptin, resistin and ghrelin, all of which significantly influence energy production and the relative proportions of fat and muscle, as well as orexigenesis [1, 42, 43].

7 Oligosaccharides: Selection of Optimal Gut Microbial Flora

Human milk oligosaccharides (HMO) are unique to humans and consist of chains of sugars between 3 and 32 residues in length [23, 24]. Despite their non-nutritional function, these sugars form a strikingly large component within breast milk, at a level similar to that of proteins overall. The synthesis of HMO depends on the action of glycosyltransferases, a group of enzymes which participate in the synthesis of other carbohydrates. HMO function as prebiotics, i.e. they assist the growth of a beneficial (probiotic) intestinal flora. Many pathogens remain in the gut by attachment to oligosaccharides attached to the enterocytic membrane. The HMO and conjugated proteins within the lumen act as alternative binding sites for pathogens, thus acting as a decoy and preventing their adhering to the intestinal lining. There is a variety in the particular HMO secreted by different mothers, the so-called lactotype, which bears comparison with blood groups [23–25]. However, there is no aspect of the lactotype corresponding to donor–recipient incompatibility, as occurs with blood transfusions. Thus, an infant may consume milk from any non-related mother. The different lactotypes have probably evolved because of varying binding affinities amongst pathogens. Certain patterns of HMO secretion are able to prevent diarrhoea secondary to infection by specific pathogens, and to inhibit HIV transmission. There is a need for more detailed research focusing on the secretion patterns for HMO and lactose in preterm milk [44, 45].

The traditional medical belief that breast milk is normally sterile has been overturned, and it now appears that breast milk has its own associated microbiota. There are various maternal factors that alter the composition of this microbiota. It also varies as breastfeeding progresses [46, 47]. The specific HMO in milk affect which microbial species colonise the gut and may partly determine which bacteria are present in breast milk.

8 Conclusion

Breast milk is a fluid which is dynamic in composition and acts in numerous different ways to promote the healthy development of the infant. It has both a nutritive and bioactive function. The constituents differ at different stages of breastfeeding and between term and preterm milk samples. Despite the numerous investigations into how breast milk is composed, research continues to identify previously unrecognised constituents. There is a pressing requirement for a study examining in a standard way breast milk samples from multiple populations. This will then allow a thorough, detailed and complete reference for the nutritive and bioactive factors in human milk to be compiled. At present, we are still at the discovery stage in understanding how breast milk contributes to the healthy development of human infants [1, 2, 48].

References

1. Ballard O, Morrow AL. Human Milk composition: nutrients and bioactive factors. *Pediatr Clin N Am*. 2013;60(1):49–74. <https://doi.org/10.1016/j.pcl.2012.10.002>.
2. Breastfeeding and the use of human milk. *Pediatrics*. 2012;129(3):e827–41.
3. Oftedal OT. The evolution of milk secretion and its ancient origins. *Animal*. 2012;6(3):355–68.
4. Castellote C, Casillas R, Ramirez-Santana C, Perez-Cano FJ, Castell M, Moretones MG, Lopez- Sabater MC, Franch A. Premature delivery influences the immunological composition of colostrum and transitional and mature human milk. *J Nutr*. 2011;141(6):1181–7.
5. Pang WW, Hartmann PE. Initiation of human lactation: secretory differentiation and secretory activation. *J Mammary Gland Biol Neoplasia*. 2007;12(4):211–21.
6. Kulski JK, Hartmann PE. Changes in human milk composition during the initiation of lactation. *Aust J Exp Biol Med Sci*. 1981;59(1):101–14.
7. Henderson JJ, Hartmann PE, Newnham JP, Simmer K. Effect of preterm birth and antenatal corticosteroid treatment on lactogenesis II in women. *Pediatrics*. 2008;121(1):e92–100.
8. Nommsen-Rivers LA, Dolan LM, Huang B. Timing of stage II lactogenesis is predicted by antenatal metabolic health in a cohort of primiparas. *Breastfeed Med*. 2012;7(1):43–9.
9. Cregan MD, De Mello TR, Kershaw D, McDougall K, Hartmann PE. Initiation of lactation in women after preterm delivery. *Acta Obstet Gynecol Scand*. 2002;81(9):870–7.
10. Nommsen LA, Lovelady CA, Heinig MJ, Lonnerdal B, Dewey KG. Determinants of energy, protein, lipid, and lactose concentrations in human milk during the first 12 mo of lactation: the DARLING study. *Am J Clin Nutr*. 1991;53(2):457–65.
11. Bauer J, Gerss J. Longitudinal analysis of macronutrients and minerals in human milk produced by mothers of preterm infants. *Clin Nutr (Edinburgh, Scotland)*. 2011;30(2):215–20.
12. Geraghty SR, Davidson BS, Warner BB, Sapsford AL, Ballard JL, List BA, Akers R, Morrow AL. The development of a research human milk bank. *J Hum Lact*. 2005;21(1):59–66.
13. Prentice A. Regional variations in the composition of human milk. In: Jensen RG, editor. *Handbook of milk composition*. San Diego, CA: Academic Press, Inc.; 1995. p. 919.
14. Liao Y, Alvarado R, Phinney B, Lonnerdal B. Proteomic characterization of human milk whey proteins during a twelve-month lactation period. *J Proteome Res*. 2011;10(4):1746–54.
15. Gao X, McMahon RJ, Woo JG, Davidson BS, Morrow AL, Zhang Q. Temporal changes in milk proteomes reveal developing milk functions. *J Proteome Res*. 2012;11(7):3897–907.
16. Lonnerdal B. Human milk proteins: key components for the biological activity of human milk. *Adv Exp Med Biol*. 2004;554:11–25.
17. Jensen RG. *Handbook of milk composition*. San Diego, CA: Academic Press, Inc.; 1995. *Pediatr Clin North Am*. Author manuscript; available in PMC 2014 February 01.
18. Valentine CJ, Morrow G, Fernandez S, Gulati P, Bartholomew D, Long D, Welty SE, Morrow AL, Rogers LK. Docosahexaenoic acid and amino acid contents in pasteurized donor Milk are low for preterm infants. *J Pediatr*. 2010;157(6):906–10.
19. Saarela T, Kokkonen J, Koivisto M. Macronutrient and energy contents of human milk fractions during the first six months of lactation. *Acta Paediatr*. 2005;94(9):1176–81.
20. Kent JC, Mitoulas LR, Cregan MD, Ramsay DT, Doherty DA, Hartmann PE. Volume and frequency of breastfeedings and fat content of breast milk throughout the day. *Pediatrics*. 2006;117(3):e387–95.
21. Valentine CJ, Morrow G, Pennell M, Morrow AL, Hodge A, Haban-Bartz A, Collins K, Rogers LK. Randomized controlled trial of docosahexaenoic acid supplementation in midwestern U.S. human milk donors. *Breastfeed Med*. 2012; <https://doi.org/10.1089/bfm.2011.0126>.
22. Martin MA, Lassek WD, Gaulin SJ, Evans RW, Woo JG, Geraghty SR, Davidson BS, Morrow AL, Kaplan HS, Gurven MD. Fatty acid composition in the mature milk of Bolivian forager-horticulturalists: controlled comparisons with a US sample. *Matern Child Nutr*. 2012;8(3):404–18.
23. Newburg DS, Ruiz-Palacios GM, Morrow AL. Human milk glycans protect infants against enteric pathogens. *Annu Rev Nutr*. 2005;25:37–58.

24. Morrow AL, Ruiz-Palacios GM, Jiang X, Newburg DS. Human-milk glycans that inhibit pathogen binding protect breast-feeding infants against infectious diarrhea. *J Nutr.* 2005;135(5):1304–7.
25. Gabrielli O, Zampini L, Galeazzi T, Padella L, Santoro L, Peila C, Giuliani F, Bertino E, Fabris C, Coppa GV. Preterm milk oligosaccharides during the first month of lactation. *Pediatrics.* 2011;128(6):e1520–31.
26. Michaelsen KF, Skafte L, Badsberg JH, Jorgensen M. Variation in macronutrients in human bank milk: influencing factors and implications for human milk banking. *J Pediatr Gastroenterol Nutr.* 1990;11(2):229–39.
27. Arslanoglu S, Moro GE, Ziegler EE. The Wapm Working Group on nutrition. Optimization of human milk fortification for preterm infants: new concepts and recommendations. *J Perinat Med.* 2010;38(3):233–8.
28. Greer FR. Do breastfed infants need supplemental vitamins? *Pediatr Clin N Am.* 2001;48(2):415–23.
29. Allen LH. B vitamins in breast milk: relative importance of maternal status and intake, and effects on infant status and function. *Adv Nutr.* 2012;3(3):362–9.
30. Dawodu A, Zalla L, Woo JG, Herbers PM, Davidson BS, Heubi JE, Morrow AL. Heightened attention to supplementation is needed to improve the vitamin D status of breastfeeding mothers and infants when sunshine exposure is restricted. *Matern Child Nutr.* 2012;10:383.
31. Pediatrics, AAo. Pediatric nutrition handbook. 6th ed. Elk Gove Village, IL: American Academy of Pediatrics; 2009.
32. Korhonen H, Marnila P, Gill HS. Milk immunoglobulins and complement factors. *Br J Nutr.* 2000 Nov;84 Suppl 1:S75–80.
33. Garofalo R. Cytokines in human milk. *J Pediatr.* 2010;156(2 Suppl):S36–40.
34. Cavaletto M, Giuffrida MG, Conti A. The proteomic approach to analysis of human milk fat globule membrane. *Clin Chim Acta.* 2004;347(1–2):41–8.
35. Van de Perre P. Transfer of antibody via mother's milk. *Vaccine.* 2003;21(24):3374–6.
36. Kobata R, Tsukahara H, Ohshima Y, Ohta N, Tokuriki S, Tamura S, Mayumi M. High levels of growth factors in human breast milk. *Early Hum Dev.* 2008;84(1):67–9.
37. Patki S, Patki U, Patil R, Indumathi S, Kaingade P, Bulbule A, Nikam A, Pishte A. Comparison of the levels of the growth factors in umbilical cord serum and human milk and its clinical significance. *Cytokine.* 2012;59(2):305–8.
38. Hirai C, Ichiba H, Saito M, Shintaku H, Yamano T, Kusuda S. Trophic effect of multiple growth factors in amniotic fluid or human milk on cultured human fetal small intestinal cells. *JPGN.* 2002;34:524–8.
39. Chailier P, Menard D. Ontogeny of EGF receptors in the human gut. *Front Biosci.* 1999;4:87–101.
40. Wagner CL, Taylor SN, Johnson D. Host factors in amniotic fluid and breast milk that contribute to gut maturation. *Clinic Rev Allerg Immunol.* 2008;34:191–204.
41. Read LC, Upton FM, Francis GL, Wallace JC, Dahlenberg GW, Ballard FJ. Changes in the growth-promoting activity of human milk during lactation. *Pediatr Res.* 1984;18(2):133–9.
42. Chang C-Y, Chao JC-J. Effect of human milk and epidermal growth factor on growth of human intestinal caco-2 cells. *JPGN.* 2002;34:394–401.
43. Khailova L, Dvorak K, Arganbright KM, Williams CS, Halpern MD, Dvorak B. Changes in hepatic cell junctions structure during experimental necrotizing enterocolitis: effect of EGF treatment. *Pediatr Res.* 2009;66(2):140–4.
44. Radulescu A, Zhang H-Y, Chen C-L, Chen Y, Zhou Y, Yu X, Otabor I, Olson JK, Besner GE. Heparin-binding EGF-like growth factor promotes intestinal anastomotic healing. *J Surg Res.* 2011;171:540–50.
45. Dvorak B, Fitch CC, Williams CS, Hurst NM, Schanler RJ. Increased epidermal growth factor levels in human milk of mothers with extremely premature infants. *Pediatr Res.* 2003;54(1):15–9.

46. Dvorak B, Fituch CC, Williams CS, Hurst NM, Schanler RJ. Concentrations of epidermal growth factor and transforming growth factor-alpha in preterm milk. In: Pickering LK, et al., editors. *Protecting infants through human milk*. Kluwer Academic/Plenum Publishers; 2004. p. 407–9.
47. Rodrigues D, Li A, Nair D, Blennerhassett M. Glial cell line-derived neurotrophic factor is a key neurotrophin in the postnatal enteric nervous system. *Neurogastroenterol Motil*. 2011;23:e44–56.
48. Boesmans W, Gomes P, Janssens J, Tack J, Berghe PV. Brain-derived neurotrophic factor amplifies neurotransmitter responses and promotes synaptic communication in the enteric nervous system. *Gut*. 2008;57:314–22.