

CURRENT CLINICAL IMAGING • RICHARD C. SEMELKA, SERIES EDITOR

WOMEN'S IMAGING

MRI with Multimodality Correlation

EDITED BY

MICHELE A. BROWN

HAYDEE OJEDA-FOURNIER

DRAGANA DJILAS

MOHAMED EL-AZZAZI

RICHARD C. SEMELKA

WILEY Blackwell






Table of Contents

[Title page](#)

[Copyright page](#)

[Contributors](#)

[Preface](#)

[Chapter 1: Pelvis MRI: introduction and technique](#)

[Imaging evaluation of the female pelvis](#)

[Indications for MRI](#)

[Patient preparation for MRI](#)

[Sequence protocols](#)

[Image optimization at 3T](#)

[Image interpretation](#)

[Chapter 2: Imaging the vagina and urethra](#)

[Vagina](#)

[Female urethra](#)

[Chapter 3: Pelvic floor imaging](#)

[Anatomy](#)

[Physiology](#)

[MR imaging of the pelvic floor](#)

[Pathology](#)

Chapter 4: Imaging the uterus

Normal anatomy

Congenital uterine anomalies

Benign disease of the uterine corpus

Benign disease of the cervix

Malignant disease of the uterine corpus and cervix

Chapter 5: Imaging the adnexa

Normal anatomy

Benign disease of the adnexa

Malignant disease of the adnexa

Malignant disease of the fallopian tube

Adnexal mass characterization algorithm

Chapter 6: Imaging maternal conditions in pregnancy

Safety

Pain in pregnancy

Obstetrical complications

Postpartum uterus

Placental imaging

Chapter 7: Fetal imaging

Indications

Safety

Normal anatomy

Fetal anomalies

Chapter 8: Breast MRI: introduction and technique

Breast cancer

Rationale for the use of breast MR

Advancements in breast MRI

ACR appropriateness guidelines for the use of breast MRI

Screening

Adjunctive evaluation for extent of disease

Additional evaluation of clinical or physical finding

Technique

American College of Radiology accreditation [42]

Patient preparation and positioning

Safety (contraindications)

CAD for breast MRI

BI-RADS assessment

Difficulties in breast MR image interpretation

Chapter 9: ACR breast MRI lexicon and interpretation

Breast MR lexicon

Breast MR interpretation

Reporting breast MR

Breast MR lexicon lesion type

Associated findings

Extramammary findings

Kinetic assessment

BI-RADS Assessment categories

Common benign breast lesions

Chapter 10: Preoperative breast cancer evaluation and advanced breast cancer imaging

Preoperative breast cancer MR evaluation

Invasive ductal carcinoma (IDC)

Ductal carcinoma *in situ* (DCIS)

Invasive lobular carcinoma (ILC)

Medullary carcinoma

Colloid carcinoma

Tubular carcinoma

Papillary carcinoma

Paget's disease

Locally advanced breast cancer

Chapter 11: Postsurgical breast and implant imaging

Introduction

Breast conservation surgery

Mastectomy

Reconstruction

Radiation therapy

Axillary node sampling/dissection

Benign procedural changes

Chapter 12: MR-guided breast interventions

Introduction

[Percutaneous guided core biopsies](#)

[Imaging protocols](#)

[Equipment](#)

[Patient selection and preparation](#)

[Patient positioning](#)

[Lesion localization](#)

[Procedure](#)

[Complications](#)

[Pearls and pitfalls](#)

[What to expect](#)

[3T biopsy experience at UCSD](#)

[Index](#)

Women's Imaging MRI with Multimodality Correlation

Edited by

Michele A. Brown, MD

Professor of Clinical Radiology
UC San Diego Health System
San Diego, CA, USA

Haydee Ojeda-Fournier, MD

Associate Professor of Clinical Radiology
Medical Director, Breast Imaging Section
Director Medical Student Education in Radiology
Moore's Cancer Center
UC San Diego Health System
La Jolla, CA, USA

Dragana Djilas, MD, PhD

Associate Professor of Radiology
Center for Diagnostic Imaging
Oncology Institute of Vojvodina
Novi Sad, Serbia

Mohamed El-Azzazi, MD, PhD

Clinical Research Scholar, MRI Section, Department of Radiology
University of North Carolina, Chapel Hill, NC, USA;
Professor of Radiology, Al-Azhar University
Cairo, Egypt;
Associate Professor of Radiology
Dammam University, Saudi Arabia; and
Consultant in Radiology
King Fahad University Hospital, Saudi Arabia

Richard C. Semelka, MD

Director, Magnetic Resonance Services;
Professor, Vice Chairman of Clinical Research; and
Vice Chairman of Quality and Safety
Department of Radiology
University of North Carolina at Chapel Hill
Chapel Hill, NC, USA

WILEY Blackwell

Copyright © 2014 by John Wiley & Sons, Inc. All rights reserved

Published by John Wiley & Sons, Inc., Hoboken, New Jersey

Published simultaneously in Canada

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4470, or on the web at www.copyright.com. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008, or online at <http://www.wiley.com/go/permissions>.

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting a specific method, diagnosis, or treatment by health science practitioners for any particular patient. The publisher and the author make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of fitness for a particular purpose. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other

things, any changes in the instructions or indication of usage and for added warnings and precautions. Readers should consult with a specialist where appropriate. The fact that an organization or Website is referred to in this work as a citation and/or a potential source of further information does not mean that the author or the publisher endorses the information the organization or Website may provide or recommendations it may make. Further, readers should be aware that Internet Websites listed in this work may have changed or disappeared between when this work was written and when it is read. No warranty may be created or extended by any promotional statements for this work. Neither the publisher nor the author shall be liable for any damages arising herefrom.

For general information on our other products and services or for technical support, please contact our Customer Care Department within the United States at (800) 762-2974, outside the United States at (317) 572-3993 or fax (317) 572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic formats. For more information about Wiley products, visit our web site at www.wiley.com.

Library of Congress Cataloging-in-Publication Data:

Women's imaging : MRI with multimodality correlation / edited by Michele A. Brown, Haydee Ojeda-Fournier, Dragana Djilas, Mohamed El-Azzazi, Richard C. Semelka.

p. ; cm.

Includes bibliographical references and index.

ISBN 978-1-118-48284-1 (cloth)

I. Brown, Michele A., editor of compilation. II. Ojeda-Fournier, Haydee, editor of compilation. III. Djilas, Dragana, editor of

compilation. IV. El-Azzazi, Mohamed, editor of compilation. V. Semelka, Richard C., editor of compilation.

[DNLM: 1. Magnetic Resonance Imaging-methods. 2. Women's Health. WN 185]

RC386.6.M34

616.07'548-dc23

2013042717

Cover design by Wiley

Contributors

Chayanin Anghong, MD

Department of Radiology
University of North Carolina at Chapel Hill
Chapel Hill, NC, USA

Jasmina Boban, MD

Center for Diagnostic Imaging
Oncology Institute of Vojvodina
Novi Sad, Serbia

Dragana Bogdanovic-Stojanovic, MD, PhD

Assistant Professor of Radiology
Center for Diagnostic Imaging
Oncology Institute of Vojvodina
Novi Sad, Serbia

Michele A. Brown, MD

Professor of Clinical Radiology
UC San Diego Health System
San Diego, CA, USA

Julie Bykowski, MD

Assistant Professor of Clinical Radiology
Moore's Cancer Center
UC San Diego Health System
La Jolla, CA, USA

Jade de Guzman, MD

Assistant Professor of Clinical Radiology

Moore's Cancer Center
UC San Diego Health System
La Jolla, CA, USA

Dragana Djilas, MD, PhD

Associate Professor of Radiology
Center for Diagnostic Imaging
Oncology Institute of Vojvodina
Novi Sad, Serbia

Mohamed El-Azzazi MD, PhD

Clinical Research Scholar, MRI Section, Department of
Radiology
University of North Carolina, Chapel Hill, NC, USA;
Professor of Radiology, Al-Azhar University
Cairo, Egypt;
Associate Professor of Radiology
Dammam University, Saudi Arabia;
Consultant in Radiology
King Fahad University Hospital, Saudi Arabia

Randy Fanous, MD, BHSc

Department of Radiology
UC San Diego Health System
San Diego, CA, USA

Michael J. Gabe, MD

Department of Radiology
UC San Diego Health System
San Diego, CA, USA

Vladimir Ivanovic, BSCEE

Center for Diagnostic Imaging

Oncology Institute of Vojvodina
Novi Sad, Serbia

Reena Malhotra

Department of Radiology
North Shore-LIJ Health System
New Hyde Park, NY, USA

Mary K. O'Boyle, MD

Clinical Professor of Radiology
Chief of Ultrasound
UC San Diego Health System
San Diego, CA, USA

Haydee Ojeda-Fournier, MD

Associate Professor of Clinical Radiology
Medical Director, Breast Imaging Section
Director Medical Student Education in Radiology
Moores Cancer Center
UC San Diego Health System
La Jolla, CA, USA

Dag Pavic, MD

Associate Professor of Radiology
Department of Radiology
Medical University of South Carolina
Charleston, SC, USA

Natasa Prvulovic Bunovic, MD

Consultant in Radiology
Center for Diagnostic Imaging
Oncology Institute of Vojvodina
Novi Sad, Serbia

Steven S. Raman, MD

Professor of Radiology, Surgery and Urology
David Geffen School of Medicine at UCLA
Los Angeles, CA, USA

Katherine M. Richman, MD

Clinical Professor of Radiology
UC San Diego Health System
La Jolla, CA, USA

Ryan C. Rockhill, MD

Body and Breast Imaging
Naval Medical Center San Diego
San Diego, CA, USA

Lorene E. Romine, MD

Assistant Clinical Professor of Radiology
UC San Diego Health System
San Diego, CA, USA

Laura E. Rueff, MD, MPH

Department of Radiology
University of California Los Angeles
Los Angeles, CA, USA

Richard C. Semelka, MD

Director, Magnetic Resonance Services
Professor, Vice Chairman of Clinical Research and
Vice Chairman of Quality and Safety
Department of Radiology
University of North Carolina at Chapel Hill
Chapel Hill, NC, USA

Shannon St Clair, MD

Department of Radiology
UC San Diego Health System
San Diego, CA, USA

Preface

Women's health issues consume a large portion of medical resources and healthcare dollars. Proper management requires a team of physicians from various specialties. Within the field of Radiology, there has been a trend toward developing a subspecialty dedicated to comprehensive imaging of women's healthcare needs, including gynecological, obstetric, genitourinary, and breast conditions. The term "Women's Imaging" is used differently in different contexts; for the purpose of this textbook, the term is used to describe imaging of the female reproductive system, including the pelvis and breast. An effective women's imager must work closely with clinical colleagues of various specialties and maintain a current understanding of diagnostic strategies, clinical implications of imaging findings, and the appropriate use of imaging tests to detect and monitor treatment.

The use of magnetic resonance imaging (MRI) for evaluation of gynecological, obstetric, and breast conditions has increased in recent years. MRI provides excellent tissue contrast resolution in the female pelvis and breast without ionizing radiation. Used together with complementary modalities, such as ultrasound and mammography, MRI has been shown to add important information to help guide patient care. The current text aims to provide the essentials of MRI in Women's Imaging, including indications, technique, and interpretation. For a number of entities, we illustrate the companion imaging studies of computed tomography, ultrasound, or mammography. Hopefully this text serves to redress the considerable underutilization of MRI in these settings. Used appropriately, MRI is cost-effective and singularly informative. There are other textbooks on the

separate topics of pelvic and breast MRI; the goal of this text is to combine and update the essentials of Women's Imaging MRI into a comprehensive and succinct overview.

The present volume is separated into two main sections: female pelvis ([chapters 1-7](#)) and breast ([chapters 8-12](#)). The first chapter presents current common indications and sample protocols for female pelvis MRI. [Chapters 2-5](#) address pathology and respective imaging findings of the vagina and female urethra, pelvic floor, uterus, and adnexa. [Chapters 6](#) and [7](#) focus on issues specific to pregnancy. [Chapter 8](#) discusses rationale and technique for MRI of the breast. [Chapters 9-12](#) are dedicated to the imaging features of breast disease and the role of MRI-guided intervention in the care of women with abnormal breast imaging findings.

This text is the collective effort of many individuals. I would like to thank the co-editors and contributors for their hard work. In addition, I am indebted to my radiology colleagues at the University of California San Diego for their help and support, with special thanks to every member of the body imaging and breast imaging divisions.

Michele A. Brown, MD

Chapter 1

Pelvis MRI: introduction and technique

Michele A. Brown & Richard C. Semelka

Imaging evaluation of the female pelvis

- Imaging plays an important role in the management of gynecological disease
- Ultrasound is often the initial imaging test
- Poor tissue contrast of CT limits gynecologic applications
- MRI benefits from excellent tissue contrast and lack of ionizing radiation
- Increased experience and availability have led to increased role of MRI
- MRI deemed appropriate by American College of Radiology for gynecological conditions, especially pre-treatment assessment of endometrial and cervical cancer, work-up of suspected adnexal mass, and evaluation of acute pelvic pain in reproductive-aged women in the setting of indeterminate ultrasound [1-4]
- Numerous gynecological and obstetric conditions are depicted by MRI, which may provide initial imaging (e.g., suspected urethral diverticulum) or problem-solving after ultrasound

Indications for MRI

([Table 1.1](#))

Table 1.1. Indications for MRI of the female pelvis

Indication	Protocol	Notes
Pelvic pain	General	FS T1WI for endometriosis
Urethral diverticulum	Urethra	Contrast if known/visualized mass
Vaginal mass	Urethra	Contrast if known/visualized mass
Pelvic floor symptoms	Pelvic floor	Sagittal images with Valsalva
Uterine anomaly	Uterine anomaly	True coronal to uterine fundus
Adenomyosis	General	Bright myometrial foci on T2WI
Fibroids	General	Add contrast if pre-embolization
Fibroid versus adnexal mass	General	Vessels extending from uterus to mass suggest uterine origin
Endometrial cancer	Uterine malignancy	High resolution T2WI and T1WI + contrast oblique to endometrium for tumor invasion
Cervical cancer	Uterine malignancy	High resolution T2WI oblique to cervix for parametrial invasion
Adnexal mass characterization	General	FS T1WI for dermoid, endometrioma
Abdominal pain in pregnancy	Maternal abdominal pain	SS-ETSE (+ FS), and steady-state GE for appendix, monitor if possible
Fetal anomaly	Fetal	SS-ETSE oriented to region of interest, monitor if possible

FS = fat saturated; T1WI = T1-weighted images; T2WI = T2-weighted images; SS = single shot; ETSE = echo-train spin-echo; GE = gradient echo

- **Benign uterine conditions**

- Anomalies
 - MRI considered imaging modality of choice
 - Informs management decisions (e.g., septate versus bicornuate uterus)
- Acquired disease
 - Problem solving for indeterminate ultrasound

- MRI allows definitive diagnosis for conditions such as urethral diverticulum, leiomyoma, adenomyosis, endometriosis, and dermoid
- **Uterine malignancy**
 - Endometrial cancer
 - Preoperative staging: deep myometrial invasion correlated with lymph node invasion [5, 6]
 - MRI shown to aid management for advanced and high grade cancer [7]
 - Cervical carcinoma
 - Depth of stromal and parametrial invasion [8, 9]
 - MRI particularly aids management for
 - Tumors larger than 2 cm
 - Endocervical tumors [10]
 - Biopsy-proved adenocarcinoma (cervical versus endometrial origin)
 - Coexistent pelvic mass(es)
 - New diagnosis of cervical cancer during pregnancy
 - Prior radiation therapy [11-15]
- **Adnexal mass**
 - Determine origin of mass
 - Tissue characterization aids specific diagnosis (e.g., endometrioma, dermoid)
 - MRI helps predict likelihood of malignancy to direct proper management and limit surgical intervention for benign disease [16, 17]
 - For known ovarian cancer, CT typically used for staging; MRI if CT contraindicated
 - MRI may yield definitive diagnosis for adnexal disease that is indeterminate on ultrasound, obviating need for follow-up imaging
- **Abdominal pain in pregnancy**

- Accurate evaluation for appendicitis (and other acute diseases) without ionizing radiation [18, 19]
- Increasing availability of MRI in acute setting
- **Fetal anomalies**
 - Problem solving for indeterminate ultrasound
 - Usefulness of MRI has increased with ultrafast sequences


Patient preparation for MRI

- Empty bladder
- Fasting 4 hours
- Optional
 - Antispasmodic (e.g., glucagon 1 mg)
 - Intra-vaginal gel [20]
- Supine position, or decubitus in late pregnancy
- Phased-array coil positioned over pelvis
- To reduce artifact, may utilize
 - Saturation band over anterior abdominal wall for non-fat-saturated sagittal
 - Supplemental anteroposterior frequency-encoding direction for axial images
- Intrauterine contraceptive devices are safely imaged [21]

Sequence protocols

- Many protocol options
- Appropriate choice depends on
 - Specific clinical question
 - Available equipment and expertise
- For known or suspected uterine disease/anomalies, T2-weighted sequences are obtained in an oblique plane oriented to uterus ([Figure 1.1](#))

Figure 1.1. Imaging planes oriented to the uterus. Multiple T2-weighted images in a patient with septate uterus. Large field-of-view single-shot sequence **(a)** is obtained first and is used to plan an oblique sagittal T2-weighted sequence **(b)** obtained parallel to the endometrium (line, **a**). The oblique sagittal is used to plan an oblique axial **(c)** obtained perpendicular to the endometrium (line, **b**). The oblique axial may then be used to plan a true coronal of the uterus **(d)** obtained parallel to the endometrium (line, **c**). In the absence of 3D T2-weighted imaging, this process assures appropriate imaging planes regardless of angle/tilt of the uterus.



- Individual sequence parameters may vary based on manufacturer, etc.
- Sequences may include
 - Single-shot (SS) echo-train spin echo (ETSE)
 - For example, HASTE or SSFSE

- Sensitive to fluid, resistant to motion and susceptibility
- Large field of view
 - Localization, evaluation of coil position
 - Coronal: evaluation of renal anomalies/obstruction
 - Axial: prescribe true sagittal view of uterus
- T2-weighted
 - Breathhold may be sufficient for benign disease
 - Non-breathhold (high-resolution) for uterine malignancy
 - With or without fat saturation
 - May be done as 3D ETSE
 - Best sequence for uterine zonal anatomy
- T1-weighted
 - Breathhold in- and out-of-phase dual echo
 - Differentiates fat- and blood-containing lesions
 - Sensitive to small foci of fat within adnexal mass
 - Non-breathhold (high-resolution) for uterine cancer
 - Chemically selective fat saturation for endometriosis
- T2/T1-weighted steady-state free precession gradient echo (GE)
 - For example, TruFISP or FIESTA
 - Rapid, resistant to motion
 - Differentiates vessels from bowel (e.g., appendix)
 - Useful for fetal and maternal imaging
- T1-weighted 3D GE pre- and post-contrast
 - Fat-suppressed GE, repeated for dynamic imaging
 - Provides enhancement information
 - May use MRA parameters (e.g., vascular malformation)
- Diffusion-weighted imaging (DWI) (optional)

- B values of 0 and at least one other value up to 1000
- Apparent diffusion coefficient (ADC) map created
- DWI sequence and ADC map interpreted together
- Aids detection of tumor, inflammation
- Additional functional techniques may have increasing role [7]
- Oblique planes oriented to the endometrium or cervix important for cancer [22]
- Protocol tailored to clinical question ([Table 1.2](#), [Table 1.3](#), [Table 1.4](#), [Table 1.5](#), [Table 1.6](#), [Table 1.7](#), [Table 1.8](#))

Table 1.2. General female pelvis

Sequence	Plane	FOV (cm)	Slice thickness (mm)
SS-ETSE	Coronal	32	8
SS-ETSE	Axial	32	8
T2 ETSE	Sagittal	24	5
T2 ETSE	Axial	24	5
T1 GE In/out- of-phase	Axial	24	5
T1 GE FS	Axial	24	5
DWI (optional)	Axial	28	6
T1 3D GE FS (pre)	Axial or sagittal	24	3
<i>Contrast</i>			
T1 3D GE FS (post × 3)	Axial or sagittal	24	3
T1 GE FS (delayed)	Axial	24	5

SS = single shot; ETSE = echo-train spin-echo; GE = gradient echo; FS = fat saturated; DWI = diffusion-weighted imaging

Table 1.3. Urethra

Sequence	Plane	FOV (cm)	Slice thickness (mm)
SS-ETSE	Coronal	32	8
T2 ETSE	Coronal	16	4
T2 ETSE	Axial	16	4
T1 GE	Axial	16	4
T1 3D GE FS (pre)	Axial	24	3
Contrast (if known or visualized lesion)			
T1 GE FS (delayed)	Axial	24	3

SS = single shot; ETSE = echo-train spin-echo; GE = gradient echo; FS = fat saturated

Table 1.4. Pelvic floor

Sequence	Plane	FOV (cm)	Slice thickness (mm)
SS-ETSE	Coronal	32	8
SS-ETSE	Axial	32	8
SS-ETSE	Sagittal	32	5
SS-ETSE (Valsalva, repeat × 3)	Sagittal	32	5 (midline slice)

SS = single shot; ETSE = echo-train spin-echo

Table 1.5. Uterine anomaly

Sequence	Plane	FOV (cm)	Slice thickness (mm)
SS-ETSE	Coronal	32	8
SS-ETSE	Axial	32	8
T2 ETSE	Sagittal (to uterus)	24	5
T2 ETSE	Axial (to uterus)	24	5
T2 ETSE	Coronal (to uterus)	24	5
T1 GE In/ out-of-phase	Coronal (to uterus)	24	5
T1 GE FS	Axial	24	5

SS = single shot; ETSE = echo-train spin-echo; GE = gradient echo; FS = fat saturated

Table 1.6. Uterine malignancy

Sequence	Plane	FOV (cm)	Slice thickness (mm)
SS-ETSE	Coronal	32	8
SS-ETSE	Axial	32	8
T2 ETSE	Sagittal	24	5
T2 ETSE	Axial	24	5
T1 GE In/out-of-phase	Axial	24	5
T1 GE FS	Axial	24	5
DWI (optional)	Axial	28	6
T1 3D GE FS (pre)	Axial or sagittal	24	3
<i>Contrast</i>			
T1 3D GE FS (post × 3)	Axial or sagittal	24	3
T1 GE FS (delayed)	Axial	24	5

SS = single shot; ETSE = echo-train spin-echo; GE = gradient echo; FS = fat saturated

Table 1.7. Maternal abdominal pain

Sequence	Plane	FOV (cm)	Slice thickness (mm)
SS-ETSE	Coronal	32–40	8
SS-ETSE	Axial	32	5
SS-ETSE FS	Axial	32	5
Steady-state GE	Coronal	32	5
Steady-state GE	Axial	32	5
T1 GE In/out-of-phase	Axial	32	5

SS = single shot; ETSE = echo-train spin-echo; GE = gradient echo; FS = fat saturated

Table 1.8. Fetal

Sequence	Plane	FOV (cm)	Slice thickness (mm)
SS-ETSE	Coronal	40	8
SS-ETSE	Axial	40	8
SS-ETSE	Sagittal	40	8
SS-ETSE (repeat as needed)	Directed	24–32	4–6
Steady-state GE (optional)	Directed	24–32	4–6
T1 GE In/out-of-phase (optional)	Directed	24–32	4–6

SS = single shot; ETSE = echo-train spin-echo; GE = gradient echo;
FS = fat saturated

Image optimization at 3T

- Potential advantages
 - Increase in signal-to-noise ratio (SNR), or
 - Similar SNR at a faster speed
- Challenges
 - Signal shading magnified by dielectric effects
 - Increased specific absorption rates (SARs)
 - Changes in optimal TR and TE
 - Increased signal inhomogeneities
 - Greater shimming challenge for extrinsic magnetic field
 - Intrinsic field distortion due to increased susceptibility/chemical shift
- Solutions [23–28]
 - Dielectric effect: dielectric pad (= radiofrequency cushion) placed between patient and surface coil
 - Susceptibility: use shorter TE/higher receiver bandwidth, higher spatial resolution

- 3D GE and ETSE sequences may benefit from higher field strength
- Consider individual patient
 - Pregnant patients less suitable for 3T due to standing wave effects from amniotic fluid and safety concerns [26]
 - Non-pregnant patients may be imaged safely and effectively at 3T using optimized parameters [28]

Image interpretation

- Large volume data acquisition
- May be useful to employ a systematic checklist ([Table 1.9](#))
- Several gynecological conditions have MRI features that allow definitive diagnosis

Table 1.9. Diagnostic checklist for female pelvis MRI

Structure	MRI features evaluated
Gynecological	
Uterine corpus	Size and position Presence of myometrial mass Endometrium thickness Junctional zone thickness
Cervix	Presence of cystic mass Presence of solid tumor Size of lesion Parametrial involvement
Vagina	Presence of cystic mass Presence of wall thickening/solid tumor
Adnexa	Ovarian size Presence of ovarian mass Cystic or solid Fat containing Blood containing Enhancement features Unilateral or bilateral Paraovarian cystic or solid mass
Non-gynecological	

Bladder	Presence of solid mass Presence of cystocele
Urethra	Presence of diverticulum Size and configuration Solid/enhancing components Presence of hypermobility
Bowel	Caliber Presence of rectocele
Musculoskeletal	Bone marrow signal Degenerative changes Traumatic injury
Lymphatic	Enlarged lymph nodes

References

1. Lee, J.H., Dubinsky, T., Andreotti, R.F., et al. ACR Appropriateness Criteria(R) pretreatment evaluation and follow-up of endometrial cancer of the uterus. *Ultrasound Quarterly* 2011; 27(2):139-45.
2. Siegel, C.L., Andreotti, R.F., Cardenes, H.R., et al. ACR Appropriateness Criteria(R) pretreatment planning of invasive cancer of the cervix. *Journal of the American College of Radiology* 2012; 9(6):395-402.
3. Harris, R.D., Javitt, M.C., Glanc, P., et al. ACR Appropriateness Criteria(R) clinically suspected adnexal mass. *Ultrasound Quarterly* 2013; 29(1):79-86.
4. Andreotti, R.F., Lee, S.I., Dejesus Allison, S.O., et al. ACR Appropriateness Criteria(R) acute pelvic pain in the reproductive age group. *Ultrasound Quarterly* 2011; 27(3):205-10.
5. Kinkel, K., Kaji, Y., Yu, K.K., et al. Radiologic staging in patients with endometrial cancer: a meta-analysis. *Radiology* 1999; 212(3):711-18.
6. Wakefield, J.C., Downey, K., Kyriazi, S., deSouza, N.M. New MR techniques in gynecologic cancer. *AJR. American Journal*

of Roentgenology 2013; 200(2):249-60

7. Frei, K.A., Kinkel, K., Bonél, H.M., et al. Prediction of deep myometrial invasion in patients with endometrial cancer: clinical utility of contrast-enhanced MR imaging - a meta-analysis and Bayesian analysis. *Radiology* 2000; 216(2):444-9.

8. Sironi, S., De Cobelli, F., Scarfone, G., et al. Carcinoma of the cervix: value of plain and gadolinium-enhanced MR imaging in assessing degree of invasiveness. *Radiology* 1993; 188(3):780-97.

9. Subak, L.L., Hricak, H., Powell, C.B., Azizi, L., Stern, J.L. Cervical carcinoma: computed tomography and magnetic resonance imaging for preoperative staging. *Obstetrics and Gynecology* 1995; 86(1):43-50.

10. Hricak, H., Powell, C.B., Yu, K.K., et al. Invasive cervical carcinoma: role of MR imaging in pretreatment work-up - cost minimization and diagnostic efficacy analysis. *Radiology* 1996; 198(2):403-9.

11. Flueckiger, F., Ebner, F., Poschauko, H., et al. Cervical cancer: serial MR imaging before and after primary radiation therapy - a 2-year follow-up study. *Radiology* 1992; 184(1):89-93.

12. Hricak, H., Swift, P.S., Campos, Z., et al. Irradiation of the cervix uteri: value of unenhanced and contrast-enhanced MR imaging. *Radiology* 1993; 189(2):381-8.

13. Weber, T.M., Sostman, H.D., Spritzer, C.E., et al. Cervical carcinoma: determination of recurrent tumor extent versus radiation changes with MR imaging. *Radiology* 1995; 194(1):135-9.

14. Yamashita, Y., Harada, M., Torashima, M., et al. Dynamic MR imaging of recurrent postoperative cervical cancer. *Journal of Magnetic Resonance Imaging* 1996; 6(1):167-71.

15. Hertel, H., Köhler, C., Grund, D., et al. Radical vaginal trachelectomy (RVT) combined with laparoscopic pelvic