



Essentials

of Autism Spectrum Disorders Evaluation and Assessment

Second Edition

- A practical guide for conducting all stages of psychological assessment
- Provides up-to date evidence based practices for ASD evaluations
- Provides expert guidance for multidisciplinary assessments

Celine A. Saulnier
Pamela E. Ventola

Alan S. Kaufman & Nadeen L. Kaufman, *Series Editors*

WILEY

Essentials of Autism Spectrum Disorders Evaluation and Assessment

Essentials of Psychological Assessment Series

Series Editors, Alan S. Kaufman and Nadeen L. Kaufman

Essentials of 16 PF 1 Assessment

by Heather E.-P. Cattell and James M. Schuerger

Essentials of Assessment Report Writing

by Elizabeth O. Lichtenberger, Nancy Mather, Nadeen

L. Kaufman, and Alan S. Kaufman

Essentials of Assessment with Brief Intelligence Tests

by Susan R. Homack and Cecil R. Reynolds

Essentials of Autism Spectrum Disorders Evaluation and Assessment

by Celine A. Saulnier and Pamela E. Ventola

Essentials of Bayley Scales of Infant Development—II Assessment

by Maureen M. Black and Kathleen Matula

Essentials of Behavioral Assessment

by Michael C. Ramsay, Cecil R. Reynolds,

and R.W. Kamphaus

Essentials of Career Interest Assessment

by Jeffrey P. Prince and Lisa J. Heiser

Essentials of CAS Assessment

by Jack A. Naglieri

Essentials of Cognitive Assessment with KAIT and Other Kaufman Measures

by Elizabeth O. Lichtenberger, Debra Broadbooks, and Alan S. Kaufman

Essentials of Conners Behavior Assessments™

by Elizabeth P. Sparrow

Essentials of Creativity Assessment

by James C. Kaufman, Jonathan A. Plucker, and John Baer

Essentials of Cross-Battery Assessment, Second Edition

by Dawn P. Flanagan, Samuel O. Ortiz, and Vincent C. Alfonso

Essentials of DAS-II 1 Assessment

by Ron Dumont, John O. Willis, and Colin D. Elliot

Essentials of Dyslexia Assessment and Intervention

by Nancy Mather and Barbara J. Wendling

Essentials of Evidence-Based Academic Interventions

by Barbara J. Wendling and Nancy Mather

Essentials of Forensic Psychological Assessment, Second Edition

by Marc J. Ackerman

Essentials of IDEA for Assessment Professionals

by Guy McBride, Ron Dumont, and John O. Willis

Essentials of Individual Achievement Assessment

by Douglas K. Smith

Essentials of KABC-II Assessment

by Alan S. Kaufman, Elizabeth O. Lichtenberger, Elaine

Fletcher-Janzen, and Nadeen L. Kaufman

Essentials of Millon™ Inventories Assessment, Third Edition

by Stephen Strack

Essentials of MMPI-ATM Assessment

by Robert P. Archer and Radhika Krishnamurthy

Essentials of MMPI-21 Assessment, Second Edition

by David S. Nichols

Essentials of Myers-Briggs Type Indicator 1 Assessment, Second Edition

by Naomi Quenk

Essentials of NEPSY1-II Assessment

by Sally L. Kemp and Marit Korkman

Essentials of Neuropsychological Assessment, Second Edition

by Nancy Hebben and William Milberg

Essentials of Nonverbal Assessment

by Steve McCallum, Bruce Bracken, and John Wasserman

Essentials of PAI 1 Assessment

by Leslie C. Morey

Essentials of Processing Assessment

by Milton J. Dehn

Essentials of Response to Intervention

by Amanda M. VanDerHeyden and Matthew K. Burns

Essentials of Rorschach 1 Assessment

by Tara Rose, Nancy Kaser-Boyd, and Michael P. Maloney

Essentials of School Neuropsychological Assessment

by Daniel C. Miller

Essentials of Specific Learning Disability Identification

by Dawn Flanagan and Vincent C. Alfonso

Essentials of Stanford-Binet Intelligence Scales (SB5) Assessment

by Gale H. Roid and R. Andrew Barram

Essentials of TAT and Other Storytelling Assessments, Second Edition

by Hedwig Teglassi

Essentials of Temperament Assessment

by Diana Joyce

Essentials of WAIS 1-IV Assessment

by Elizabeth O. Lichtenberger and Alan S. Kaufman

Essentials of WIAT 1-III and KTEA-II Assessment

by Elizabeth O. Lichtenberger and Kristina C. Breaux

Essentials of WTSC 1-IV Assessment, Second Edition

by Dawn P. Flanagan and Alan S. Kaufman

Essentials of WJ III™ Cognitive Abilities Assessment, Second Edition

by Fredrick A. Schrank, Daniel C. Miller, Barbara J. Wendling,

and Richard W. Woodcock

Essentials of WJ III™ Tests of Achievement Assessment

by Nancy Mather, Barbara J. Wendling, and

Richard W. Woodcock

Essentials of WMS1-IV Assessment

by Lisa Whipple Drozdzick, James A. Holdnack,

and Robin C. Hilsabeck

Essentials of WNV™ Assessment

by Kimberly A. Brunnett, Jack A. Naglieri,

and Steven T. Hardy-Braz

Essentials of WPPSI™-III Assessment

by Elizabeth O. Lichtenberger and Alan S. Kaufman

Essentials of WRAML2 and TOMAL-2 Assessment

by Wayne Adams and Cecil R. Reynolds

Essentials of Autism Spectrum Disorders Evaluation and Assessment, Second Edition

by Celine A. Saulnier and Pamela E. Ventola

Essentials

of Autism Spectrum Disorders Evaluation and Assessment

Second Edition

Celine A. Saulnier

Pamela E. Ventola

WILEY

Copyright © 2024 by John Wiley & Sons, Inc. All rights reserved.

Published by John Wiley & Sons, Inc., Hoboken, New Jersey.

Published simultaneously in Canada.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4470, or on the web at www.copyright.com. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008, or online at <http://www.wiley.com/go/permission>.

Trademarks: Wiley and the Wiley logo are trademarks or registered trademarks of John Wiley & Sons, Inc. and/or its affiliates in the United States and other countries and may not be used without written permission. All other trademarks are the property of their respective owners. John Wiley & Sons, Inc. is not associated with any product or vendor mentioned in this book.

Limit of Liability/Disclaimer of Warranty: While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages. Further, readers should be aware that websites listed in this work may have changed or disappeared between when this work was written and when it is read. Neither the publisher nor authors shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services or for technical support, please contact our Customer Care Department within the United States at (800) 762-2974, outside the United States at (317) 572-3993 or fax (317) 572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic formats. For more information about Wiley products, visit our web site at www.wiley.com.

Library of Congress Cataloging-in-Publication Data for:

Hardback ISBN: 9781119982517

Cover Design: Wiley

Cover Image: © Robert/Adobe Stock Photos

Set in 10/12pt Adobe Garamond Pro by Straive, Pondicherry, India

*To Tony, Lucienne, & Vivi, thank you for your
unwavering love and support. ~ Celine*

*To Noah, Maddy, & Tucker, thank you for your unfaltering strength,
love, and endless encouragement. ~ Pam*

TABLE OF CONTENTS

	About the Authors	x
	Series Preface	xi
	Preface	xiii
	Acknowledgments	xiv
One	Overview	1
	Diagnostic Criteria	1
	References	7
Two	Assessment of Developmental and Cognitive Skills	9
	Selecting Instruments	10
	Developmental and Early Cognitive Measures	12
	Measures of Cognition and Intelligence in School-Age Individuals and Adults	16
	Measures of Nonverbal Intelligence	19
	Neuropsychological Assessments	20
	Qualitative Observations	22
	Summary	23
	References	24
	Test Yourself	26
	Answers	27
Three	Speech, Language, and Communication Assessment	28
	Receptive Language	31
	Expressive Language	32
	Pragmatic Language/Social Communication	34
	Stages of Language Development	36
	Formulation of Findings	38
	Summary	38
	References	42
	Test Yourself	43
	Answers	43

Four	Assessment of Adaptive Behavior, Executive Functioning, and Behavioral Regulation Skills	45
	Assessment of Adaptive Behavior	45
	Assessment of Executive Functioning	52
	Assessment of Behavioral Regulation Skills	53
	Summary	56
	References	56
	Test Yourself	58
	Answers	59
Five	Developmental History and Record Review	60
	Clinical Interview	61
	Methods of Collecting Information on Current and Historical Presentation	70
	Summary	74
	References	75
	Test Yourself	75
	Answers	76
Six	Direct Diagnostic Assessment	77
	Screening Measures for Risk of ASD	77
	Diagnostic Rating Scales	79
	Direct Diagnostic Assessment	81
	Considerations for First-Time Diagnostic Evaluations with Adolescents and Adults	94
	Direct Measures of Autism Symptomatology	96
	Options for Remote/Telehealth Diagnostic Assessments for ASD	99
	Summary	100
	References	101
	Test Yourself	103
	Answers	104
Seven	Diagnostic Differentials and Co-Occurring Conditions	105
	Intellectual Disability	106
	Communication Disorders	110
	Learning Disabilities	111
	Attention Deficit/Hyperactivity Disorder	114
	Anxiety Disorders	115
	Depression and Mood Disorders	116
	Obsessive-Compulsive Disorder (OCD)	118
	TIC Disorders and Tourette's Disorder	119
	Psychiatric Conditions in Adulthood	120
	Summary	122
	References	123
	Test Yourself	125
	Answers	127

Eight	Catching the Misdiagnosed, Late Diagnosed, and Missed for Diagnosis	128
	Factors Related to Racial, Ethnic, Socioeconomic, and Geographical Disparities	129
	Factors Related to Sex Differences in ASD Symptomatology	130
	Factors Related to Gender and Sexual Identity Development	133
	Factors Related to Subtle or Complex Symptom Expression	134
	References	136
	Test Yourself	137
	Answers	138
Nine	Preparing for Transition to Adulthood	139
	Idea and Transition Planning	139
	Assessments Supporting the Transition Plan Process	140
	Needs to Consider for Adulthood	146
	Job Coaching and Vocational Training	148
	Academic Accommodations and Modifications	149
	Life Skills and Life Coaching/Mentoring	151
	Transitional and College-Preparation Programs	153
	References	156
	Test Yourself	159
	Answers	160
Ten	Diagnostic Formulations and Case Conceptualizations	161
	Providing Feedback of Results	161
	The Written Report	162
	Case Samples	163
	Case Sample 1: Initial Diagnosis—Toddler	164
	Case Sample 2: Re-Evaluation—School-Aged Child with Complex Presentation	175
	Case Sample 3: Initial Diagnosis: Adult Female	191
	Chapter Summary	205
	Author Index	206
	Subject Index	211

ABOUT THE AUTHORS

Dr. Celine A. Saulnier is a licensed clinical psychologist and owner of Neurodevelopmental Assessment & Consulting Services (NACS) in Decatur, GA. She trained and worked at the Yale Child Study Center's Autism Program for nearly a decade before relocating to Emory University School of Medicine and the Marcus Autism Center, where she oversaw a large-scale clinical research program. In 2018, she opened NACS, where she specializes in diagnostic evaluations throughout the lifespan and conducts trainings worldwide. Dr. Saulnier has published numerous articles and is an author of the *Vineland Adaptive Behavior Scales*, Third Edition. She also wrote another book in the Essentials series, *Essentials of Adaptive Behavior Assessments for Neurodevelopmental Disorders* with Dr. Cheryl Klaiman.

Dr. Pamela E. Ventola is an Associate Professor at the Yale Child Study Center and Chief Science Officer at Cogstate. Dr. Ventola has vast experience conducting psychological and neuropsychological assessments with autistic individuals and individuals with other developmental disabilities. Her current research focuses on biological markers of treatment response in autistic children. Dr. Ventola also consults with pharmaceutical sponsors on industry-sponsored clinical trials in autism spectrum disorder (ASD) and rare neurodevelopmental disorders. Dr. Ventola has published extensively on ASD and neurodevelopmental disorders.

SERIES PREFACE

In the *Essentials of Psychological Assessment* series, we have attempted to provide the reader with books that will deliver key practical information in the most efficient and accessible style. The series features assessment instruments in a variety of domains, such as cognition, personality, education, and neuropsychology. For the experienced clinician, books in the series offer a concise yet thorough way to master use of the continuously evolving supply of new and revised instruments, as well as a convenient method for keeping up to date on the tried-and-true measures. The novice will find here a prioritized assembly of all the information and techniques that must be at one's fingertips to begin the complicated process of individual psychological diagnosis.

Wherever feasible, visual shortcuts to highlight key points are used alongside systematic, step-by-step guidelines. Chapters are focused and succinct. Topics are targeted for an easy understanding of the essentials of administration, scoring, interpretation, and clinical application. Theory and research are continually woven into the fabric of each book, but always to enhance clinical inference, never to sidetrack or overwhelm. We have long been advocates of “intelligent” testing—the notion that a profile of test scores is meaningless unless it is brought to life by the clinical observations and astute detective work of knowledgeable examiners. Test profiles must be used to make a difference in the child's or adult's life, or why bother to test? We want this series to help our readers become the best intelligent testers they can be.

In this volume, *Essentials of Autism Spectrum Disorder Assessments, Second Edition*, the widely renowned authors illustrate an up-to-date comprehensive developmental model for multidisciplinary diagnostic evaluations for ASD. This model is derived from years of experience in conducting multidisciplinary diagnostic evaluations for ASD following standards of best practice. Given the neurodevelopmental nature of ASD, symptoms unfold over the course of early development and subsequently impact multiple areas of functioning. For this reason, the developmental skills that

ultimately need to be assessed often require clinicians with different disciplinary expertise. Thus, in this Essentials series, the authors outline the components of best practices while highlighting the necessity of integrating findings from multiple sources so that one comprehensive and cohesive diagnostic formulation can be provided for an individual's optimal care. We also emphasize the need to take a strengths-based approach to capitalize upon the innumerable exceptional abilities and characteristics of ASD that should be honed to foster more vulnerable skills.

Alan S. Kaufman, PhD, and
Nadeen L. Kaufman, EdD, Series Editors,
Neag School of Education,
University of Connecticut

PREFACE

In the *Essentials of Autism Spectrum Disorder Assessments*, the authors illustrate a comprehensive developmental model for multi-disciplinary diagnostic evaluations for ASD. This model is derived from years of experience in conducting multidisciplinary diagnostic evaluations for ASD following standards of best practice. Given the neurodevelopmental nature of ASD, symptoms unfold over the course of early development and subsequently impact multiple areas of functioning. For this reason, the developmental skills that ultimately need to be assessed often require clinicians with different disciplinary expertise. Thus, in this Essentials series, the authors outline the components of best practices while highlighting the necessity of integrating findings from multiple sources so that one comprehensive and cohesive diagnostic formulation can be provided for an individual's optimal care. We also emphasize the need to take a strengths-based approach to capitalize upon the innumerable exceptional abilities and characteristics of ASD that should be honed to foster more vulnerable skills.

ACKNOWLEDGMENTS

The completion of this book would not have been possible without the help and support from so many people. This publication exemplifies the knowledge and expertise in autism spectrum and related disorders, diagnostic evaluation, and childhood development that was acquired over decades thanks to the teachings and collaborations of our incredible mentors, colleagues, and friends at the University of Connecticut, Yale Child Study Center/Yale University School of Medicine, and the Marcus Autism Center/Emory University School of Medicine including but certainly not limited to Drs. Ami Klin, Fred Volkmar, Sara Sparrow, Kasia Chawarska, Deborah Fein, Marianne Barton, Julie Wolf, and Cheryl Klaiman. Working alongside you has cultivated our knowledge, expertise, and passion for what we do!

We would also like to thank our fabulous contributors to this revised edition, including Leah Booth, Erin Brooker-Lozott, Anna Milgramm, and Sydney Terroso. In addition, we thank our editors at Wiley including Monica Rogers & Janaki Kumudam for your patience and assistance with this revision, as well as Alan and Nadeen Kaufman for recognizing the utility and necessity for including the topic of autism assessments within the *Essentials* series.

Finally, our overwhelming gratitude to the children, individuals, and families who have shared their lives and experiences with us through the years. Thank you!

OVERVIEW

DIAGNOSTIC CRITERIA

Autism spectrum disorder (ASD) is among the most common childhood disorders, with prevalence rates currently estimating 1 in 36 school-aged children (Maenner, Warren, Williams et al., 2023). ASD is four times more prevalent in males than females, but as will be highlighted later in this book, the diagnostic presentation of females assigned at birth is complex and, consequently, could suggest an underrepresentation of true female prevalence rates. Although the cause(s) of ASD is likely neurobiological/genetic in nature, the exact etiology remains unknown. As such, the diagnosis of ASD has historically been determined based on behavioral symptomatology that, through the lens of a medical model, are generally described as delays, deficits, impairments, or similar terminology that implies the need for remediation or a “cure” (Pellicano & den Houting, 2022). As the spectrum has broadened over the 21st century to include an increasing number of individuals with intact language and cognition, many autistic individuals identify as being neurodiverse or neurodivergent and do not seek to be cured or viewed as disordered, but rather wish to be accepted and supported for their unique differences (den Houting, 2019). Nevertheless, the label still implies that the individual is experiencing challenges, vulnerabilities, and in some cases delays/deficits that negatively affect multiple areas in life. The authors of this book have tremendous respect for neurodiversity, as we do for every individual across the extremely diverse and complex spectrum, including those more severely affected. Thus, in our aim to help educate and train clinicians on how to effectively identify ASD based on behavioral “symptoms,” we strive to focus on the diagnostic label serving to enhance the quality of life and self-determination rather than to imply curing or fixing anybody. We will use terms

such as symptoms, disorder, condition, and disability throughout the text given that this terminology is still part of the diagnostic criteria and medical nomenclature to which clinicians are exposed. We will also use identity-first language (autistic individual) opposed to person-first language (individual with autism) to respect that not all individuals view ASD as something separate from who they are.

The current diagnostic criteria for ASD put forth in the *Diagnostic and Statistical Manual, Fifth Edition, Text Revision* (DSM-5-TR; APA, 2022) require persistent symptoms in two broad areas: (1) *the presence of delays/deficits/challenges in social communication and interaction* in all three of the following areas either currently or by developmental history: (a) social-emotional reciprocity (e.g., shared interests, shared affect, and conversations), (b) nonverbal communication (e.g., integrated eye contact, and directed facial expressions), (c) use and understanding of gestures and body posture), and (d) in developing, maintaining, and understanding relationships (e.g., interactive imaginative play, interest in peers, and making friends); and (2) *the presence of restricted, repetitive, or stereotyped behaviors* that include at least two of the following four areas: (a) stereotypical motor movements or use of objects (e.g., hand flapping, spinning, lining up toys, and repetitive or stereotyped speech), (b) insistence on sameness or routine (e.g., difficulty with transitions, changes in routine, and rigidity in thinking), (c) highly restricted interests that become overly intense or all-consuming (e.g., intense interests on topics and fixation on objects), and (d) sensory processing impairments (e.g., hyper- or hypo-reactivity to sensory input, or seeking out sensory input). This combination of social vulnerabilities and stereotyped behaviors must be present in early development (even if symptoms go undetected until adolescence or adulthood) and clinically impairing across multiple contexts in life.

Rapid Reference 1.1

Social communication and interaction criteria (must include all three either currently or by history):

1. Challenges in social-emotional reciprocity
2. Challenges in nonverbal communication
3. Challenges in developing, maintaining, and understanding relationships

Restricted, repetitive, and stereotyped behaviors (must include at least two either currently or by history)

1. Stereotypical motor movements or use of objects
2. Insistence on sameness or routine
3. Highly restricted interests
4. Sensory processing impairments

In addition to listing the diagnostic criteria for the two categories of social communication/interaction symptoms and restricted, repetitive, and stereotyped behaviors, the DSM-5 has three severity levels that clinicians need to apply to both categories. Level 1 is specified for individuals who require support at the lowest level (e.g., symptoms interfere with functioning in one or more contexts, but they are mild); Level 2 is specified for those who require substantial support (e.g., there are marked impairments that interfere across multiple contexts); and Level 3 is specified for those who require very substantial support (e.g., delays or deficits are severe and entail comprehensive supports across contexts). Clinicians can specify different levels based on each category. For instance, an individual with intact language and cognition may only require Level 1 support for their social communication and interaction skills, but same individual's stereotypical behaviors might be so intrusive to life that Level 2 is required for this category.

The DSM-5 also offers clinical specifiers for ASD that allow for descriptions of common co-occurring conditions. These include specifying whether ASD is accompanied by co-occurring intellectual impairment or language impairment (e.g., nonverbal, minimally verbal, phrase speech, and sentence speech), whether ASD is associated with a known genetic or other medical condition (e.g., Fragile X syndrome and Down syndrome), environmental factor, other neurodevelopmental (e.g., ADHD), mental (e.g., anxiety and depression), or behavioral problem, or whether co-occurring catatonia is present. Catatonic behaviors can include posturing, limited response to external stimuli, stupor, or mutism. Catatonia can be present in as many as 10% of autistic individuals, with features

DON'T FORGET

The DSM-5 has three levels of severity that need to be applied to each category of symptoms (social communication/interaction and restricted, repetitive, and stereotyped behaviors) as follows:

- *Level 1:* the individual requires support
- *Level 2:* the individual requires substantial support
- *Level 3:* the individual requires very substantial support

DON'T FORGET

The DSM-5 has clinical specifiers that allow for indicating all co-occurring conditions. This helps inform levels of support needed as well as types of interventions.

- With or without accompanying intellectual impairment
- With or without accompanying language impairment
- Associated with a known genetic or medical condition, or environmental factor
- Associated with another neurodevelopmental, mental, or behavioral problem
- With catatonia

generally developing in late adolescence (Vaquerizo-Serrano, Salazar De Pablo, Singh, & Santosh, 2021).

There is vast heterogeneity in symptom expression across the spectrum and throughout development, with some individuals experiencing only mild challenges in certain areas and other individuals experiencing quite debilitating challenges. Symptoms that impact autistic individuals are not just limited to the criteria for ASD, but also include varying levels of speech and language, cognition, emotional and behavioral regulation, and attention and executive functioning. Thus, clinicians need to be knowledgeable of these broader areas. When autism was first defined in the 1900s, the vast majority of individuals had co-occurring language and cognitive deficits that significantly affected development. However, current epidemiological data show that only about 38% of autistic children in the United States have co-occurring intellectual disability (ID; Maenner et al., 2023). This shift in levels of cognitive and language functioning has sparked a great deal of controversy in the field. Where historically there used to be a clear delineation of impairment to meet diagnostic criteria; currently, the diagnostic line between what is considered neurotypical and neurodiverse is quite blurred (Rodgaard, Jensen, Vergnes, Soulieres, & Mottron, 2019). This raises concerns about whether autism is a single condition or perhaps multiple disabilities, and whether the spectrum has become too broad to be accurately defined. Subtypes of autism were developed in the DSM-IV and DSM-IV-TR (APA, 1994, 2000) with autistic disorder, Asperger's disorder, pervasive developmental disorder, not otherwise specified, Rett's disorder, and childhood disintegrative disorder. However, research failed to show that these subtypes indeed had distinct phenotypes. Rather, research indicated that the main factor differentiating subtypes was the geographical site or clinicians providing the diagnoses (Lord, Petkova, Hus et al., 2012). Even expert clinical teams using gold-standard measures were not reliable in determining the categorical subtypes, of autism which led to the transition to the more dimensional definition of ASD.

Certainly, individuals with significant cognitive and language impairments and affiliated behavioral challenges have drastically different needs than individuals with intact cognition and language. In 2021, the Lancet Commission published a report

that formally recognized the term “profound autism” to distinguish autistic individuals who are so severely affected that they will likely require 24-hour support throughout their lives (Lord, Charman, Havdahl et al., 2022). In contrast, individuals with intact language and cognition can often function quite independently across aspects of life. Many of these individuals identify as being neurodiverse and do not view themselves as having a disability but

DON'T FORGET

There is significant variability in symptom expression throughout the spectrum in autism symptomatology, speech and language skills, cognition, emotional and behavioral regulation, and attention and executive functioning. Thus, clinicians need to be knowledgeable of these broad areas.

rather advocate for acceptance of neurodivergence. Yet, it is important to highlight that even some higher-cognitive autistic individuals struggle tremendously in life despite their strong intellectual capacity. For this reason, the historically misused phrase “high functioning autism” has resulted in an underestimate of services and supports that autistic individuals require because of the erroneous assumption that intact language and cognition equate to functional independence (Alvares, Bebbington, Cleary et al., 2019). Similarly, the phrase “low functioning autism” misrepresents individuals with cognitive and language impairments who may be functioning quite well despite their delays. Clinicians are therefore advised to keep terminology descriptive of specific skill areas, and use the severity levels and clinical specifiers to advocate for appropriate supports.

ASD has a complex genetic etiology with symptoms unfolding over the first few years of life. Infants who will go on to develop ASD are not born exhibiting full autism symptom expression; rather, as symptoms emerge, developmental milestones are derailed resulting in a diverging trajectory away from neurotypical development. Thus, some infants appear to be developing within age expectations in the first few months or even the first year of life, but they then fail to keep pace with chronological age expectations during the second year. As developmental divergence occurs, typically developing milestones can fade, while the presence of atypical behaviors can emerge. It is therefore essential that clinicians be knowledgeable of normative development in addition to merely observing for the presence of ASD symptoms because it is often the case that infants at risk for ASD have words, eye contact, gestures, etc., but the limited frequency and range of these skills are what causes concern.

Research indicates that 80–90% of parents report their first concerns about their child’s development by the second birthday and often earlier. However, the mean age of diagnosis continues to be over the age of four despite these concerns (Maenner et al., 2023; Chawarska et al., 2007). When experienced clinicians make a diagnosis of ASD at 18 to 24 months, the stability of diagnosis is

CAUTION

Avoid using terminology such as “high functioning or low functioning autism,” as these labels make erroneous assumptions about an individual’s true profile of strengths and/or vulnerabilities. Preferred language describes specific profiles:

- Autism without cognitive impairment
- Autism without language impairment
- Minimally verbal or nonverbal
- Profound autism

CAUTION

Clinicians involved in early detection of ASD need to be knowledgeable of normative developmental milestones in addition to ASD symptoms because as ASD unfolds in infancy and toddlerhood, it is often the limited frequency and range of expected skills that causes concern rather than the emergence of atypical behaviors.

CAUTION

Most parents of children who develop ASD express concerns regarding their child's development prior to the second birthday, over two years before diagnostic evaluations take place, on average. Professionals need to be extra vigilant in not only validating concerns but also taking immediate action to assess and identify potential likelihood for ASD.

CAUTION

Social motivation and reciprocity should not be interpreted in isolation when distinguishing autism from non-autism conditions.

DON'T FORGET

Autistic individuals with intact language and cognition may have awareness of their challenges and therefore are at great risk for mood disorders such as anxiety and depression. These symptoms can emerge at any point in development and should be monitored and treated accordingly.

quite strong, also around 80–90% (Chawarska et al., 2009). This highlights the extremely concerning gap between when first concerns are raised and when something is actually done to help the child, often because of limited awareness of the early markers of ASD by professionals on the front line as well as barriers to access to services. These facts underscore the necessity for clinicians of all disciplines to learn about and be vigilant of the early signs of ASD so that children can be effectively evaluated and efficiently diagnosed. These include (but are not limited to) delays or deficits in joint attention, shared affect, social smiling, response to name, frequency and range of gestures, communicative intent, and functional speech.

In autistic toddlers and preschoolers who have intact cognition and language, social vulnerabilities can be masked by strengths in other areas, such as precocious language, fixation on numbers and letters to the point of self-reading, and burgeoning circumscribed interests. It is typically not until these children are immersed in social settings, where the social demands outweigh the capacity to engage, that social concerns are raised. As autistic children with intact language and cognition move through the school-age years,

they can be motivated to interact with their peers, but they might insert themselves into interactions inappropriately and/or lack the appropriate social awareness to effectively navigate interactions. Yet, they can have just enough awareness to understand the failed nature of their attempts, placing them at great risk for experiencing symptoms of anxiety, depression, and isolation. This contrasts with more profound forms of autism, where individuals can be socially passive, withdrawn, or less likely to initiate interactions with others. Nonetheless, social motivation and reciprocity should not be interpreted in isolation when distinguishing autism from non-autism conditions.

Based on our collective and extensive experience in conducting multidisciplinary diagnostic evaluations, we present a comprehensive developmental approach to the assessment and diagnosis of ASD, including how to assess for the common differential and co-occurring condition throughout the lifespan, how to interpret assessment results and convey the findings in a comprehensive written report, and how to use individual profiles of strengths and vulnerabilities to inform educational and intervention programming. Beginning with Chapter 2, we discuss the necessity for obtaining a baseline of developmental and cognitive functioning to identify individual profiles of strengths and vulnerabilities. Chapter 3 then outlines how the speech, language, and communication assessment informs the diagnostic process. Given the significance of adaptive behavior to functional outcome in ASD, Chapter 4 describes how assessing for adaptive behavior, executive functioning, and social-emotional skills helps inform both diagnosis and intervention.

Chapters 5 and 6 collectively outline the assessment of autism symptomatology, which entails the gathering of historic information, observing the individual in natural contexts, and directly assessing behaviors through interaction and play. Chapter 7 focuses on the common differentials and co-occurring conditions that arise through referrals for diagnostic evaluations in ASD throughout the lifespan. As the spectrum continues to broaden, Chapter 8 focuses on the many challenges in correctly identifying ASD when symptom expression is obscured by racial/ethnic/cultural disparities, gender and sexual identity, and other complexities that can result in misdiagnoses or missed diagnoses. Chapter 9 covers the extremely important transitional period from school-age to adulthood, where evaluations are often necessary to inform postsecondary education, college, vocations, and adult life. Chapter 10 ties the process together with three samples of integrated reports from model comprehensive diagnostic evaluations—one of a toddler, one of a school-aged child, and one of a female adult receiving an ASD diagnosis for the first time. Our hope is that this model will be useful in informing both burgeoning clinicians just starting out in the field and seasoned professionals who are experiencing an increased exposure to ASDs and, subsequently, are seeking knowledge of how to effectively identify, diagnose, and/or refer patients for evaluations.

REFERENCES

- Alvares, G. A., Bebbington, K., Cleary, D., Evans, K., Glasson, E. J., Maybery, M. T., Pillar, S., Uljarević, M., Varcin, K., Wray, J., & Whitehouse, A. J. (2019). The misnomer of 'high functioning autism': Intelligence is an imprecise predictor of functional abilities at diagnosis. *Autism*, 24(1), 221–232.
- American Psychiatric Association. (1994). *Diagnostic and Statistical Manual of Mental Disorders* (4th ed.). Washington, DC: Author.
- American Psychiatric Association. (2000). *Diagnostic and Statistical Manual of Mental Disorders* (4th ed., text rev.). Washington, DC: Author.

- American Psychiatric Association. (2022). *Diagnostic and Statistical Manual of Mental Disorders* (5th ed., text rev.). Washington, DC: Author.
- Chawarska, K., Klin, A., Paul, R., Macari, S., & Volkmar, F. (2009). A prospective study of toddlers with ASD: Short-term diagnostic and cognitive outcomes. *Journal of Child Psychology and Psychiatry*, 50(10), 1235–1245.
- Chawarska, K., Paul, R., Klin, A., Hannigen, S., Dichtel, L. E., & Volkmar, F. (2007). Parental recognition of developmental problems in toddlers with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 37(1), 62–73.
- den Houting, J. (2019). Neurodiversity: An insider's perspective. *Autism*, 23(2), 271–273.
- Lord, C., Charman, T., Havdahl, A., et al. (2022). The Lancet Commission on the future of care and clinical research in autism. *Lancet*, 399(10321), 271–334.
- Lord, C., Petkova, E., Hus, V., et al. (2012). A multisite study of the clinical diagnosis of different autism spectrum disorders. *Archives of General Psychiatry*, 69(3):306–313.
- Maenner, M. J., Warren, Z., Williams, A. R., et al. (2023). Prevalence and characteristics of autism spectrum disorder among children aged 8 years: Autism and developmental disabilities monitoring network, 11 sites, United States, 2020. *MMWR Surveillance Summaries*, 72(2):1–14.
- Pellicano, E., & den Houting, J. (2022). Annual research review: Shifting from 'normal science' to neurodiversity in autism science. *Journal of Child Psychology and Psychiatry*, 63(4), 381–396.
- Rodgaard, E., Jensen, K., Vergnes, J., Soulieres, I., & Mottron, L. (2019). Temporal changes in effect sizes of studies comparing individuals with and without autism. A meta-analysis. *JAMA*, 76(11):1124–1132.
- Vaquerizo-Serrano J., Salazar De Pablo G., Singh J., & Santosh P.(2021). Catatonia in autism spectrum disorders: A systematic review and meta-analysis. *Eur Psychiatry*, 65(1):e4.