

Edited by Paul T. Anastas

 WILEY-VCH

Green Processes



Volume 7: Green Synthesis

Volume Editor:
Chao-Jun Li



Handbook of Green Chemistry

Volume 7
Green Synthesis

Edited by
Chao-Jun Li

Related Titles

Jiménez-González, C., Constable, D. J. C.

Green Chemistry and Engineering

A Practical Design Approach

2010

ISBN: 978-0-470-17087-8

Dunn, P., Wells, A., Williams, M. T. (eds.)

Green Chemistry in the Pharmaceutical Industry

2010

ISBN: 978-3-527-32418-7

Loos, K. (ed.)

Biocatalysis in Polymer Chemistry

2010

ISBN: 978-3-527-32618-1

Reichardt, C., Welton, T.

Solvents and Solvent Effects in Organic Chemistry

Fourth, Updated and Enlarged Edition

2010

ISBN: 978-3-527-32473-6

Pignataro, B. (ed.)

Tomorrow's Chemistry Today

**Concepts in Nanoscience, Organic Materials and Environmental Chemistry
Second Edition**

2009

ISBN: 978-3-527-32623-5

Roesky, H. W., Kennepohl, D. (eds.)

Experiments in Green and Sustainable Chemistry

2009

ISBN: 978-3-527-32546-7

Handbook of Green Chemistry

Volume 7
Green Synthesis

Edited by
Chao-Jun Li



WILEY-VCH Verlag GmbH & Co. KGaA

The Editor

Prof. Dr. Paul T. Anastas

Yale University
Center for Green Chemistry & Green Engineering
225 Prospect Street
New Haven, CT 06520
USA

Volume Editor

Prof. Dr. Chao-Jun Li

McGill University
Department of Chemistry
801 Sherbrooke Street West
Montreal, QC H3A 2K6
Canada

Handbook of Green Chemistry – Green Processes

Vol. 7: Green Synthesis
ISBN: 978-3-527-32602-0
Vol. 8: Green Nanoscience
ISBN: 978-3-527-32628-0
Vol. 9: Designing Safer Chemicals
ISBN: 978-3-527-32639-6

Set III (3 volumes):
ISBN: 978-3-527-31576-5

Handbook of Green Chemistry

Set (12 volumes):
ISBN: 978-3-527-31404-1
oBook ISBN: 978-3-527-62869-8

The cover picture contains images from Corbis
Digital Stock (Dictionary) and PhotoDisc, Inc./Getty
Images (Flak containing a blue liquid).

All books published by **Wiley-VCH** are carefully produced. Nevertheless, authors, editors, and publisher do not warrant the information contained in these books, including this book, to be free of errors. Readers are advised to keep in mind that statements, data, illustrations, procedural details or other items may inadvertently be inaccurate.

Library of Congress Card No.: applied for

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library.

Bibliographic information published by the Deutsche Nationalbibliothek

The Deutsche Nationalbibliothek lists this publication in the Deutsche Nationalbibliografie; detailed bibliographic data are available on the Internet at <http://dnb.d-nb.de>.

© 2012 Wiley-VCH Verlag & Co. KGaA,
Boschstr. 12, 69469 Weinheim, Germany

All rights reserved (including those of translation into other languages). No part of this book may be reproduced in any form – by photoprinting, microfilm, or any other means – nor transmitted or translated into a machine language without written permission from the publishers. Registered names, trademarks, etc. used in this book, even when not specifically marked as such, are not to be considered unprotected by law.

Composition Thomson Digital, Noida, India
Printing and Binding betz-druck GmbH,
Darmstadt
Cover Design Adam Design, Weinheim

Printed in the Federal Republic of Germany
Printed on acid-free paper

ISBN: 978-3-527-32602-0

Contents

About the Editors XIII

List of Contributors XV

Preface XIX

1 Atom Economy: a Challenge for Enhanced Synthetic Efficiency 1

Barry M. Trost

1.1 Vinylidenes 2

1.1.1 Cycloisomerization of Hydroxyalkynes 2

1.1.2 Reconstitutive Condensation 6

1.2 Redox Isomerization 7

1.2.1 Allyl Alcohols 7

1.3 Ruthenacyclopentadiene Intermediates 9

1.4 Ruthenacyclopentene Intermediates 13

1.4.1 Intermolecular Alkene-Alkyne Coupling 13

1.4.2 Butenolide Formation and Related Reactions 16

1.4.3 Pyran Formation 17

1.4.4 Intramolecular Alkene-Alkyne Coupling 19

1.4.5 [5 + 2] Cycloaddition 19

1.4.6 Vinyl Ketones as Alkyne Partners 21

1.5 Allylic C–H Insertion 22

1.6 Reactions of Alkenes 24

1.6.1 Allene–Alkene Coupling 24

1.6.2 Heterocycles via Allene–Alkene Coupling 27

1.7 Conclusion 28

References 29

2 Evaluating the Greenness of Synthesis 35

David J.C. Constable and Concepción “Conchita” Jiménez-González

2.1 General Considerations About Green Chemistry and Green Engineering Metrics 35

2.2 Selected Metrics Used in the Past 37

2.2.1 Yield 37

2.2.1.1 Effective Mass Yield 37

2.2.2	E-Factor	38
2.2.3	Atom Economy	39
2.2.3.1	Key Assumptions About Atom Economy	40
2.2.3.2	How Atom Economy Is calculated	41
2.3	Reaction Mass Efficiency	42
2.4	Mass Intensity and Mass Productivity (Mass Efficiency)	43
2.5	Cost Implications and Green Chemistry Metrics	47
2.6	Life-Cycle Assessment Metrics	49
2.7	Process Metrics	50
2.7.1	Materials	52
2.7.1.1	Physical Form and Properties	52
2.7.1.2	Mass	52
2.7.1.3	Inherent Hazard	53
2.7.1.4	Cost	54
2.7.1.5	Renewability	54
2.7.1.6	Recyclability	55
2.7.2	Equipment and Operability Intertwined	55
2.7.2.1	Type and Number of Unit Operations	56
2.7.2.2	Size of Unit Operations	56
2.7.2.3	Scalability	56
2.7.2.4	Controllability	57
2.7.2.5	Robustness	58
2.7.2.6	Throughput/Cycle Time	58
2.7.2.7	Energy	59
2.7.2.8	Cleaning and Maintenance	60
2.7.3	EHS Hazards and Risk	61
2.7.3.1	Occupational Exposure Hazards and Risk	62
2.7.3.2	Process Safety Hazards Risk	63
2.7.3.3	Environmental Hazards and Risk	63
2.7.4	Quality	64
2.7.4.1	Purity	64
2.8	Conclusions	65
	References	65

3 Alternative Feedstocks for Synthesis 69

Arno Behr and Leif Johnen

3.1	Introduction	69
3.1.1	Renewable Resources as Natural Feedstock	69
3.1.2	Challenges of Using Renewable Resources	70
3.2	Carbohydrates	71
3.2.1	Polysaccharides	71
3.2.1.1	Cellulose	71
3.2.1.2	Starch	72
3.2.2	Disaccharides	73
3.2.2.1	Sucrose	74

3.2.2.2	Lactose	74
3.2.3	Monosaccharides	74
3.2.3.1	D-Glucose	75
3.2.3.2	D-Fructose	75
3.3	Lignin	76
3.4	Fats and Oils	77
3.4.1	Catalytic Derivatization of Unsaturated Fatty Compounds	78
3.4.1.1	Selective Catalytic Hydrogenation	79
3.4.1.2	Selective CC Linkage Reactions	80
3.4.1.3	CN Linkage Reactions	80
3.4.1.4	CO Linkage Reactions	81
3.4.2	Glycerol	82
3.4.2.1	Glycerol Esters	83
3.4.2.2	Etherification	83
3.4.2.3	Glycerol Oxidation and Dehydration	84
3.5	Terpenes	85
3.6	Carbon Dioxide	87
3.6.1	Reactions with Alkanes, Alkenes, and Dienes	88
3.6.2	Conversion to Formic Acid and Dimethylformamide	89
3.6.3	Plasma Activation of Carbon Dioxide	89
	References	90
4	Synthesis in Green Solvents	93
	<i>László Orha, Geoffrey R. Akien, and István T. Horváth</i>	
4.1	The Role of Solvents in Synthesis	93
4.2	Types of Solvent	94
4.2.1	Atomic Liquids	94
4.2.2	Molecular Liquids	94
4.2.3	Ionic Liquids	95
4.2.4	Solvent Polarity	95
4.2.5	Protic Solvents	96
4.3	Problems with Solvents	96
4.4	Application of Green Solvents	96
4.4.1	Water	96
4.4.2	Fluorous Solvents	102
4.4.3	Supercritical Carbon Dioxide	107
4.4.4	Ionic Liquids	112
4.5	Conclusion	117
	References	117
5	Development and Application of Isocyanide-based Multicomponent Reactions	121
	<i>Jieping Zhu, Qian Wang, and Mei-Xiang Wang</i>	
5.1	Introduction	121
5.2	Basic Principle of MCRs	124

5.3	Discovering Novel MCRs	125
5.3.1	Union Concept	127
5.3.2	Rational Substrate Design	127
5.3.3	Mechanism-Based Design	137
5.3.3.1	“Split-Ugi” Reaction	137
5.3.3.2	Ugi–Smiles 4CR	139
5.3.3.3	Activation of Imines by Other Electrophiles	140
5.3.4	Serendipity	142
5.4	MCRs Imitated by Addition of Isocyanides to Alkynes	144
5.5	Metal-Catalyzed IMCRs	146
5.6	Enantioselective P-3CR	149
5.7	Application in Medicinal Chemistry and in Natural Product Synthesis	151
5.8	Conclusion	152
	References	152
6	Flow Syntheses	159
	<i>Charlotte Wiles and Paul Watts</i>	
6.1	Introduction	159
6.1.1	Continuous Flow Reactors: What Are They and How are They Used?	159
6.2	Examples of Their Use as Tools for the Research Chemist	160
6.2.1	Liquid Phase	160
6.2.1.1	Solvent Free	160
6.2.1.2	Liquid–Liquid Phase	163
6.2.1.3	Elevated Reaction Temperatures	164
6.2.1.4	Reduced Reaction Temperatures	174
6.2.2	Solid–Liquid Phase	176
6.2.2.1	Solid-Supported Catalysts	177
6.2.2.2	Solid-Supported Reagents	185
6.2.2.3	Solid-Supported Scavengers	190
6.2.3	Gas–Liquid Phase	190
6.2.4	Gas–Liquid–Solid Phase	191
6.2.5	Biocatalysis	194
6.2.5.1	Liquid Phase	194
6.2.5.2	Immobilized Biocatalytic Flow Reactors	197
6.2.6	Photochemistry	199
6.2.6.1	Homogeneous Photochemical Reactions	199
6.2.6.2	Heterogeneous Photochemical Reactions	202
6.3	Process Intensification Achieved Through the Use of Flow Reactors	204
6.3.1	Synthesis of Azo Dyes	205
6.3.2	Synthesis of Ionic Liquids Under Continuous Flow	206
6.3.3	DSM Nitration	207
6.3.4	Synthesis of Rimonabant	208

6.3.5	Biocatalytic Synthesis of Vitamin A	209
6.4	Conclusions and Outlook	210
	References	210
7	Synthesis Without Protecting Groups	215
	<i>Reinhard W. Hoffmann</i>	
7.1	The Present Use of Protecting Groups	215
7.2	Protecting Group-Free Synthesis?	218
7.3	Use of <i>In Situ</i> Protections in Lieu of Short-Term Protecting Groups	220
7.4	Follow Nature's Biogenetic Routes to Avoid Protecting Groups	221
7.5	Apply Functional Group-Tolerant Construction Reactions to Avoid Protecting Groups	224
7.6	Aim for Higher Chemoselectivity to Avoid Protecting Groups	224
7.7	Change the Order of Synthesis Steps to Avoid Protecting Groups	227
7.8	Enlist Latent Functionality to Avoid Explicit Protecting Group Steps	229
7.9	Summary	231
	References	233
8	Biological Synthesis of Pharmaceuticals	237
	<i>Junhua Tao and Alex Chu</i>	
8.1	Introduction	237
8.2	New Enzymes for Chemical Synthesis	237
8.2.1	Enzymatic Halogenation	238
8.2.2	Macrocyclization	239
8.2.3	Glycosylation	241
8.2.4	Heterocyclization	242
8.2.5	Methylation	243
8.2.6	Oxygenation	244
8.3	Synthesis of Pharmaceuticals via Isolated Enzymes	244
8.3.1	Penicillins and Cephalosporins	244
8.3.2	Pregabalin	246
8.3.3	Atorvastatin	247
8.3.4	Levetiracetam	248
8.4	Synthesis of Pharmaceuticals via Whole Cells	249
8.4.1	Paclitaxel	249
8.4.2	Epothilones	251
8.4.3	Oseltamivir	251
8.4.4	Avermectins	252
8.5	Conclusion	254
	References	255

9	Syntheses via C–H Bond Functionalizations	259
	<i>Lutz Ackermann, Anant R. Kapdi, Harish K. Potukuchi, and Sergei I. Kozhushkov</i>	
9.1	Introduction	259
9.2	Direct Arylations of Arenes	261
9.2.1	“Green” Aspects of Direct Arylation of Aryl C–H Bonds	264
9.2.2	Chelation-Assisted Direct Arylations of Arenes	265
9.2.3	Non-Directed Direct Arylations of Arenes	275
9.2.4	Direct Arylations of Heteroarenes	279
9.2.4.1	Direct Arylations of Electron-Deficient Heteroarenes	279
9.2.4.2	Direct Arylations of Electron-Rich Heteroarenes	280
9.3	Catalytic Oxidative Arylations of (Hetero)arenes	293
9.3.1	Introduction	293
9.3.2	Oxidative Homocouplings	295
9.3.3	Cross-Dehydrogenative Arylations	296
9.4	Conclusion	298
	References	298
10	Synthesis Without Metals	307
	<i>Takahiko Akiyama</i>	
10.1	Introduction	307
10.2	Organic Reactions Promoted by Non-Metallic Catalysts	308
10.3	Asymmetric Organocatalysts	311
10.3.1	Introduction	311
10.3.2	Classification by Reaction Types	312
10.3.2.1	Covalent Organocatalysis	312
10.3.2.2	Non-Covalent Organocatalysis	312
10.3.3	Organocatalysts	312
10.3.3.1	Cinchona Alkaloids and Derivatives	312
10.3.3.2	Proline Derivatives and MacMillan’s Catalyst	313
10.3.3.3	Peptide Catalysts	318
10.3.3.4	Ketone Catalysts	319
10.3.3.5	Phase-Transfer Catalysts	319
10.3.3.6	Amine Catalysts	320
10.3.3.7	Guanidinium Salts	321
10.3.3.8	Hydrogen Bond Catalysts	321
10.3.3.9	Stronger Brønsted Acid Catalysts	324
10.3.3.10	Counteranion Catalysis	329
10.4	Conclusion	330
	References	331
11	Chemistry Beyond Functional Group Transformation	335
	<i>Zhiping Li and Rong Yu</i>	
11.1	Introduction	335
11.2	C–H Bond Activation	336

11.2.1	sp^3 C–H Bond Activation	337
11.2.1.1	C–C Bond Formation	337
11.2.1.2	C–N Bond formation	343
11.2.1.3	C–O Bond Formation	347
11.2.2	sp^2 C–H Bond Activation	349
11.2.2.1	C–C Bond Formation	349
11.2.2.2	C–N Bond Formation	353
11.2.2.3	C–O Bond Formation	353
11.3	C–C Bond Activation	353
11.3.1	Utilization of Strained Molecules	354
11.3.2	Utilization of Chelating Substrates	355
11.3.3	Utilization of Activating Functional Groups	357
11.4	C–O Bond Activation	357
11.5	C–F Bond Activation	357
11.6	C–N Bond Activation	359
11.7	Small Molecule Activation	359
11.7.1	H ₂	359
11.7.2	O ₂	360
11.7.3	CH ₄	361
11.8	Conclusions and Outlook	362
	References	363
12	Synthesis Assisted by Electricity	369
	<i>Jun-ichi Yoshida and Seiji Suga</i>	
12.1	Electroorganic Synthesis in Green Reaction Media (Homogeneous System)	369
12.1.1	Electroorganic Synthesis in Aqueous Solutions	369
12.1.2	Electroorganic Synthesis in Supercritical Carbon Dioxide	370
12.1.3	Electroorganic Synthesis in Ionic Liquids	370
12.2	Electroorganic Synthesis in Liquid–Liquid Biphasic Systems	372
12.3	Electroorganic Synthesis in Thermomorphic Liquid–Liquid Biphasic Systems	373
12.4	Electroorganic Synthesis in Solid–Liquid Biphasic Systems	374
12.4.1	Solid-Supported Mediators	375
12.4.2	Solid–Liquid Biphasic System for Electrolysis Without Intentionally Added Supporting Electrolyte	376
12.4.2.1	SPE Technology	376
12.4.2.2	Electrolysis Using Solid-Supported Bases	377
12.5	Electroorganic Synthesis in Microflow Systems	378
12.5.1	Electrochemical Microflow Cells	379
12.5.2	Paired Electrolysis in Microflow Systems	380

12.5.3	Electroorganic Synthesis in a Microflow System Without Using Intentionally Added Supporting Electrolyte	381
12.6	Future Outlook	383
	References	383
13	Parameterization and Tracking of Optimization of Synthesis Strategy Using Computer Spreadsheet Algorithms	387
	<i>John Andraos</i>	
13.1	Introduction	387
13.2	Synthesis Strategy Parameterization	390
13.3	Case Study: Lysergic Acid	393
	References	413
	Index	415

About the Editors

Series Editor



Paul T. Anastas joined Yale University as Professor and serves as the Director of the Center for Green Chemistry and Green Engineering there. From 2004–2006, Paul was the Director of the Green Chemistry Institute in Washington, D.C. Until June 2004 he served as Assistant Director for Environment at the White House Office of Science and Technology Policy where his responsibilities included a wide range of environmental science issues including furthering international public-private cooperation in areas of Science for Sustainability such as Green Chemistry. In 1991, he established the industry-government-university partnership Green Chemistry Program, which was expanded to include basic research, and the Presidential Green Chemistry Challenge Awards. He has published and edited several books in the field of Green Chemistry and developed the 12 Principles of Green Chemistry.

Volume Editor



Chao-Jun Li (FRSC, UK) received his PhD at McGill University (1992) and was an NSERC Postdoctoral Fellow at Stanford University (1992–1994). He was an Assistant Professor (1994), Associate Professor (1998) and Full Professor (2000–2003) at Tulane University, where he received a NSF CAREER Award (1998) in organic synthesis and the 2001 US Presidential Green Chemistry Challenge Award (Academic). In 2003, he became a Canada Research Chair (Tier I) in Organic/Green Chemistry and a Professor of Chemistry at McGill University in Canada. He serves as the Co-Chair of the Canadian Green Chemistry and Engineering Network, the Director of CFI Infrastructure for Green Chemistry and Green Chemicals, and Co-Director the FQRNT Center for Green Chemistry and

Catalysis (Quebec). He is the current Associate Editor for Americas for the journal of *Green Chemistry* (published by the Royal Society of Chemistry). He has been widely recognized as the leader in Green Chemistry for Organic Synthesis in developing innovative and fundamentally new organic reactions that defy conventional reactivities and have high synthetic efficiency.

List of Contributors

Lutz Ackermann

Georg-August-Universität Göttingen
Institut für Organische und
Biomolekulare Chemie
Tammannstrasse 2
37077 Göttingen
Germany

Geoffrey R. Akien

City University of Hong Kong
Department of Biology and Chemistry
Kowloon
Hong Kong

Takahiko Akiyama

Gakushuin University
Department of Chemistry
Mejiro 1-5-1, Toshima-ku
171-8588 Tokyo
Japan

John Andraos

CareerChem
504-1129 Don Mills Road
Toronto, ON M3B 2W4
Canada

Arno Behr

Technische Universität Dortmund
Fakultät Bio- und
Chemieingenieurwesen
Lehrstuhl Technische Chemie A
Emil-Figgestrasse 66
44227 Dortmund
Germany

Alex Chu

Metabornics, Inc.
SuZhou
Jiangsu 215600
China

David J.C. Constable

Lockheed Martin
Energy, Environment, Safety and Health
Gaithersburg, MD 20878
USA

Reinhard W. Hoffmann

Philipps-Universität
Fachbereich Chemie
Hans-Meerwein-Strasse
35032 Marburg
Germany

István T. Horváth

City University of Hong Kong
Department of Biology and Chemistry
Kowloon
Hong Kong

Concepción “Conchita”

Jiménez-González

GlaxoSmithKline
Operational Sustainability,
Sustainability and Environment
Research Triangle Park, NC 27709
USA

Leif Johnen

Technische Universität Dortmund
Fakultät Bio- und
Chemieingenieurwesen
Lehrstuhl Technische Chemie A
Emil-Figgestrasse 66
44227 Dortmund
Germany

Anant R. Kapdi

Georg-August-Universität Göttingen
Institut für Organische und
Biomolekulare Chemie
Tammannstrasse 2
37077 Göttingen
Germany

Sergei I. Kozhushkov

Georg-August-Universität Göttingen
Institut für Organische und
Biomolekulare Chemie
Tammannstrasse 2
37077 Göttingen
Germany

Zhiping Li

Renmin University of China
Department of Chemistry
No.59, Zhong Guan Cun Street
Beijing 100872
China

László Orha

Eötvös University
Institute of Chemistry
Pázmány Péter Sétány 1/A
1117 Budapest
Hungary

Harish K. Potukuchi

Georg-August-Universität Göttingen
Institut für Organische und
Biomolekulare Chemie
Tammannstrasse 2
37077 Göttingen
Germany

Seiji Suga

Okayama University
Graduate School of Natural Science and
Technology
Division of Chemistry and Biochemistry
3-1-1 Tsushima-naka, Kita-ku
700-8530 Okayama
Japan

Junhua Tao

Metabomics, Inc.
SuZhou
Jiangsu 215600
China

Barry M. Trost

Stanford University
Department of Chemistry
Stanford, CA 94305
USA

Mei-Xiang Wang

The Key Laboratory of Bioorganic
Phosphorus Chemistry & Chemical
Biology (Ministry of Education)
Department of Chemistry
Tsinghua University
100084 Beijing
China

Qian Wang

École Polytechnique Fédérale de
Lausanne
EPFL-SB-ISIC-LSPN
BCH 5304 (Bât BCH)
1015 Lausanne
Switzerland

Paul Watts

The University of Hull
Department of Chemistry
Cottingham Road
Hull HU6 7RX
UK

Charlotte Wiles

The University of Hull
Department of Chemistry
Cottingham Road
Hull HU6 7RX
UK

and

Chemtrix BV
Burgemeester Lemmensstraat 358
6163JT Geleen
The Netherlands

Jun-ichi Yoshida

Kyoto University
Graduate School of Engineering
Department of Synthetic Chemistry and
Biological Chemistry
Nishikyo-ku
615-8510 Kyoto
Japan

Rong Yu

Renmin University of China
Department of Chemistry
No.59, Zhong Guan Cun Street
Beijing 100872
China

Jieping Zhu

École Polytechnique Fédérale de
Lausanne
EPFL-SB-ISIC-LSPN
BCH 5304 (Bât BCH)
1015 Lausanne
Switzerland

Preface

Ever since the synthesis of urea by Friedrich Wöhler near two centuries ago, organic synthesis has become the foundation of modern medicines for human health, produced new agrochemicals to boost world food supply, created various synthetic fibers for daily usages, and bestowed a colorful enchantment through synthetic dyes. In spite of these great achievements, the general features of organic syntheses have been, by and large, unchanged over a century: e.g., non-renewable feedstock, batch reactor, and refluxing. In addition, classical organic syntheses often produce stoichiometric amount of waste, use organic solvents and sometimes dangerous reagents, require extensive protection-deprotection of functional groups, need pre-functionalized starting materials, and involve multi-step operations, which resulted in low efficiency in resource utilization and led to various concerns due to waste generations. While, in the past, the primary goal of organic syntheses is “to get the target product”, the sustainability of chemical synthesis becomes a more and more important issue. This volume of Green Syntheses illustrated some examples to address this issue ranging from starting materials, reaction design, choice of solvent, energy input, to reactor design. The chapter by Trost describes the general principle of greener synthesis; the chapter by Behr shows examples of using renewable feedstocks for making chemical products; the chapter by Horvath describes the use of alternative solvents for organic synthesis; the chapters by Zhu, Hoffman and Watts describe methods of reducing synthetic steps by running multi-component reactions, avoiding protecting groups, and in flow respectively; the chapters by Ackermann and Li show examples of direct conversion of C–H bonds; the chapters by Varma and Yoshida presents alternative energy input in chemical reactions through light and electricity; the chapters by Tao and Akiyama give examples of using enzymes and organo catalysts for synthetic purposes; and finally the chapter by Andraos uses computation methods to evaluate the relative efficiency of different synthetic routes. We hope that these examples will provide food-for-thought for further innovations in developing greener syntheses.

Montreal, April 2012

C-J Li

1

Atom Economy: a Challenge for Enhanced Synthetic Efficiency*Barry M. Trost*

The design of structure for function is the major task for helping to solve problems ranging from material science to human health. The demands and expectations for extremely high levels of performance frequently increases the molecular complexity needed. Thus, a major goal must be to allow the synthesis of such complex molecular arrays in a time-effective manner. The strategic design for the synthesis of complex molecules derives from the available basic tools – the reactions, reagents, and catalysts. Although some might think we have a pretty full toolbox, the reality is that, in most likelihood, only a very small fraction of the true total number of reactions possible is known today. Hence a great unknown awaits us, and chipping away at those unknown processes presents a great opportunity for discovery that will undoubtedly change the practice of the science.

In undertaking a program of discovery for new processes, the characteristics that defines the requirements for these new reactions/reagents/catalysts must be appreciated. In 1983, selectivity was noted as key to evolving reasonable efficiency in the synthesis of complex molecules [1]. The issue of chemoselectivity, defined as discriminating reactivity among various bond types in a molecule without employing activating or blocking groups, was placed at the top of the list! There is no question that problems of chemoselectivity are the single biggest factor in creating synthetic inefficiencies. More than 25 years later, the primacy of this selectivity issue was still noted [2]. It is so pervasive in the science that it undoubtedly will remain the greatest challenge for a long time to come. The second issue is regioselectivity, which is defined as orientational control in the joining of a reagent with an unsymmetrical functional group. Controlling stereochemistry constitutes the third major challenge. There are two fundamentally different issues embodied within this topic – controlling relative stereochemistry or diastereoselectivity and absolute stereochemistry or enantioselectivity.

By and large, selectivity was equated with efficiency. However, just a little further thought makes us realize that we are missing one key aspect of efficiency by focusing only on selectivity, that is, by ignoring an obvious but neglected aspect, which is, simply put, how much of what you put into your pot ends up in your product? In 1991, this fundamental and critical issue was explicitly recognized and referred to as “atom economy” [3]. In 1992, the *E* factor was introduced, which also

provided a quantitative metric to evaluate the degree of atom economy [4]. Atom economy, which basically emphasizes maximal use of raw materials and minimization of waste, has become one of the 12 principles of Green Chemistry [5]. Making synthetic chemistry more “environmentally benign by design” has become a mantra.

The ideal reaction is one in which 100% of what is introduced into a reaction ends up as product and, if anything else is required, it is needed only catalytically. Thus, a bimolecular reaction should be a single addition and a unimolecular reaction should be an isomerization. Such processes are clearly known and heavily used, such as a Diels–Alder reaction and Claisen and Cope rearrangements. However, such processes constitute only a very small fraction of our toolbox. In the late 1980s, we began a deliberate program to invent processes that theoretically are 100% atom economic. I note this goal as theoretical since to achieve it fully requires the yield to be quantitative, which rarely occurs, but at least the possibility does exist. A reaction of the type $A + B \rightarrow C + D$ suffers from the fact that it theoretically cannot be 100% atom economic and also suffers from the issue of yields typically being less than quantitative. This overview reports the evolution of one of our programs for semi-rationally inventing atom economic processes based upon catalysis with ruthenium complexes.


1.1

Vinylidenes


1.1.1

Cycloisomerization of Hydroxyalkynes


The formation of reactive intermediates provides possible opportunities for new reaction design. An attractive highly reactive intermediate, carbenes, which demonstrate numerous useful synthetic pathways, most notably by addition to alkenes and alkynes and also insertion into X–H bonds, where X is both carbon and heteroatoms, suffers from problems associated with their accessibility. Undoubtedly, the most useful class of precursor is the diazo compounds, whose safety problems restrict their use. For the specific case of vinylidenes, an attractive possibility is a terminal alkyne which is isomeric with a vinylidene. Although the thermolysis appears to effect this transformation (Equation 1.1, path a), the extraordinarily high temperatures required make the prospect of a transition metal-catalyzed version (Equation 1.1, path b) attractive. The early studies of Werner [6] using Rh and Bruce and co-workers [7] using Ru proved the facility with which such species would form; however, the studies focused on the formation and isolation of the vinylidene–metal complexes and their stoichiometric reactions.



Based on these studies and choosing to focus on Ru for practical considerations such as cost, we envisioned a possible catalytic cycle shown in Scheme 1.1, wherein an ω -alkynyl alcohol **1** would cycloisomerize to the dihydropyran. Although McDonald's group has pioneered the use of molybdenum- and tungsten-mediated processes, several issues related to chemoselectivity and the common need for stoichiometric amounts make the development of other catalysts for such processes desirable [8]. Using $\text{CpRu}(\text{PAr}_3)_2\text{Cl}$ -based complexes, less electron-rich arylphosphine ligands such as *m*- or *p*-fluorophenylphosphines promote such processes (Equation 1.2) [9].






Scheme 1.1 Cycloisomerization of hydroxyalkenes.







While the presence of propargylic hydroxy groups has proven problematic (see below), the presence of a propargylic *N*-tosylamido group is tolerated (Equation 1.3).

Interestingly, the presence of a bis-homopropargylic secondary amide does not lead to insertion into the N–H (as happens with a tungsten catalyst [8c]) competing with formation of the dihydropyran as shown in Equation 1.4 [10]. This reaction extends to the insertion into a phenolic OH to form benzofurans (Equation 1.5). In this case, the presence of an amine such as *n*-butylamine or pyridine appears to be required [11]. It should be noted that insertion into the benzylic OH to form the pyran system does not compete.




Interestingly, making the reaction less basic, increasing the amount of *N*-hydroxysuccinimide to 3 equiv., and changing the phosphine reorients the course of the reaction to an oxidative cyclization to form lactones. For example, γ -butyrolactones form fairly readily from homopropargylic alcohols (Equation 1.6) and even preferentially when the possibility for forming the thermodynamically more stable six-membered ring lactones could occur (Equation 1.7) [12]. The synthesis of the butyrolactones is optimized using trifurylphosphine as the ligand. Nevertheless, bis-homopropargyl alcohols also undergo oxidative cyclization. Thus, the homopropargyl alcohol **3** in Equation 1.2 under the conditions of Equation 1.6 except that the phosphine ligand is changed to tris(*p*-methoxyphenyl) phosphine gives the δ -lactone **4** in 65% yield (Equation 1.8) [9]. As expected, competing a six- versus a seven-membered ring leads exclusively to cyclization to form the six-membered ring (Equation 1.9).





After esterification of alcohol **4** with inversion of configuration, the oviposition attractant pheromone of the mosquito *Culex pipens fatigans* **5** is formed. The stereochemistry of this sequence originated from the asymmetric dihydroxylation of the *trans*-alkene. Although the oxidative cyclization is significantly less atom economic than the cycloisomerization, since it requires stoichiometric amounts of *N*-hydroxysuccinimide, the recyclability of the resultant succinimide mitigates this defect somewhat. This switch in pathway as a function of pH can be understood by the sequence depicted in Scheme 1.2. In particular, the initially formed vinyl Ru complex **6**, whose coordination with the anion derived from hydroxysuccinimide, thus facilitating protonation to the enol ether, preferentially protonates to form the carbene complex **7**. Now the electrophilic carbene carbon undergoes nucleophilic addition of this same anion, leading ultimately to fragmentation with generation of the lactone.






Scheme 1.2 Mechanistic rationale for oxidative cyclization.

1.1.2



Reconstitutive Condensation

The intermolecular variant of the O–H insertion reaction gets stuck at the stage of the initial adduct **8**. We envisioned that if $R^1 = \text{allyl}$, coordination of the double bond to the metal would initiate a Claisen-type process to form the π -allylruthenium complex **9**, whose reductive elimination would form the allyl ketone starting from terminal alkynes and allyl alcohols (Equation 1.10). Gratifyingly, this prediction was fully realized as shown in Equation 1.11 [13]. A tertiary ester does not undergo elimination under these reaction conditions (Equation 1.12). Dihydroxylation of the double bond and subsequent acidification effect cyclodehydration to form furans in two overall “steps.” Subsequent elimination of the elements of acetic acid completes a synthesis of rosefuran **10**, one of the most prized fragrances [14].



With terminal alkynes bearing propargylic alcohols, formation of the vinylidene is complicated by the competitive formation of the allenylidene as shown in Equation 1.13.

In the case of a substrate bearing a second hydroxyl group, the allenylidene can undergo nucleophilic attack at the γ -carbon to re-form a vinylidene [15]. In such an event, the newly formed vinylidene can then undergo the reconstitutive condensation shown in Equation 1.14 [16]. This reaction provides access to the spiroketal of calyculin A, a nanomolar inhibitor of serine/threonine phosphatases.





1.2

Redox Isomerization




1.2.1

Allyl Alcohols

In the reconstitutive addition in which we used the allyl alcohol as both reactant and solvent, we noted that propanal was detected as a by-product. This observation led us to consider the use of CpRuL_2Cl as a catalyst for redox isomerization of allyl alcohols in general. Indeed, subjecting allyl alcohol **12** to the standard Ru catalyst effects chemoselective isomerization to the saturated ketone **13** (Equation 1.15) [17]. Normally starting from aldehyde **11**, the typical route to saturated ketone **13**, path b, involves a stoichiometric oxidant for one step. Furthermore, there is also a chemoselectivity issue in this example, thereby further enhancing the inefficiency by requiring a protecting group operation. By simply switching to an unsaturated organometallic for the first step, a stoichiometric oxidant is avoided, as is any need for protecting groups, thereby improving the overall atom economy. For more substituted double bonds, the (indenyl) $\text{Ru}(\text{PPh}_3)_2\text{Cl}$ catalyst, which can slip from an η^5 to an η^3 complex, allows a decrease in steric hindrance in addition to being accompanied by the formation of a coordinatively unsaturated ruthenium which improves the reactivity. Thus, the 1,2-disubstituted alkene substrate **14** (Equation 1.16) undergoes redox isomerization within 3 h, whereas such substrates barely react at all with the $\text{CpRu}(\text{PPh}_3)_2\text{Cl}$ complex.



Application of this catalyst system to propargyl alcohols provides α,β -unsaturated aldehydes (Equation 1.17) and ketones (Equation 1.18) [18]. The ease of accessibility of the substrates by simple addition of terminal alkynes to aldehydes followed by this redox isomerization constitutes a highly chemoselective and atom economic strategy to these unsaturated carbonyl compounds. The chemoselectivity problems of the direct aldol condensation and the poor atom economy of olefination methods make this new strategy the most efficient and reliable approach to these units.



A significant improvement in the catalyst adds further utility to this strategy. Using 1–5 mol% of indenyl complex 15 along with indium triflate and camphorsulfonic acid (CSA) as cocatalysts, the redox isomerization in Equation 1.19 was completed in 20 min compared with 1.5 h under the indium chloride cocatalyst conditions.