



The Schema Therapy Clinician's Guide

A Complete Resource for Building and Delivering Individual,
Group and Integrated Schema Mode Treatment Programs

Joan M. Farrell • Neele Reiss • Ida A. Shaw



WILEY Blackwell

The Schema Therapy Clinician's Guide

The Schema Therapy Clinician's Guide

*A Complete Resource for Building
and Delivering Individual, Group
and Integrated Schema Mode
Treatment Programs*

Joan M. Farrell, Neele Reiss,
and Ida A. Shaw

Illustrations by Britta Finkelmeier

WILEY Blackwell

This edition first published 2014
© 2014 John Wiley & Sons, Ltd
Except for illustrations © 2014 Britta Finkelmeier.

Registered Office

John Wiley & Sons, Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Offices

350 Main Street, Malden, MA 02148-5020, USA

9600 Garsington Road, Oxford, OX4 2DQ, UK

The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

For details of our global editorial offices, for customer services, and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell.

The right of Joan M. Farrell, Neele Reiss, and Ida A. Shaw to be identified as the authors of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book.

Limit of Liability/Disclaimer of Warranty: While the publisher and author(s) have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. It is sold on the understanding that the publisher is not engaged in rendering professional services and neither the publisher nor the author shall be liable for damages arising herefrom. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

Library of Congress Cataloging-in-Publication Data

Farrell, Joan M., author.

The schema therapy clinician's guide : a complete resource for building and delivering individual, group and integrated schema mode treatment programs / Joan M. Farrell, Neele Reiss, and Ida A. Shaw.

p. ; cm.

Includes bibliographical references and index.

ISBN 978-1-118-50917-3 (pbk.) – ISBN 978-1-118-50918-0 (cloth)

I. Reiss, Neele, author. II. Shaw, Ida A., author. III. Title.

[DNLM: 1. Psychotherapy–methods–Handbooks. 2. Personality Disorders–therapy–Handbooks. 3. Psychotherapy, Group–methods–Handbooks. WM 34]

RC480.5

616.89'14–dc23

2014002676

A catalogue record for this book is available from the British Library.

Cover image: Alamy, ID C84W3F

Cover design by Design Deluxe

Set in 10.5/13pt MinionPro by SPi Publisher Services, Pondicherry, India

Contents

List of Figures and Tables	ix
About the Authors	xi
Foreword	xiii
Acknowledgments	xvii
About the Companion Web Site	xix
1 Introduction	1
1.1 Background	1
1.2 The Chapters	3
2 The Basics of Schema Therapy	5
2.1 The Theoretical Model	5
2.2 Goals and Stages of Schema Therapy	9
2.3 Limited Reparenting	10
2.4 The Components of Schema Therapy	20
3 The Integrated Schema Therapy Treatment Program	21
3.1 The Patients: Who Are Good Candidates?	26
3.2 The Group Schema Therapy Sessions	27
3.2.1 The “Welcome” session	28
3.2.2 Schema Therapy Education sessions	28
3.2.3 Mode Awareness sessions	28
3.2.4 Mode Management sessions	29
3.2.5 Experiential Mode Work sessions	29
3.3 The Individual Schema Therapy (IST) Sessions	30
3.4 Options for the Delivery of the Treatment Program	31
3.5 Structural Aspects of the Group Sessions	35
3.6 Assessment and Orientation	36
3.7 The Milieu of the Treatment Program	43
3.7.1 Multidisciplinary teams in intensive settings	43
3.7.2 The physical environment	45

4	The Group Schema Therapy Sessions	47
4.1	General Therapist Tips for Group Schema Therapy	52
4.2	The Welcome Group Session	59
4.3	Five Schema Therapy Education (STE) Sessions	70
4.4	The 12 Mode Awareness Sessions	95
4.5	The 12 Mode Management Sessions	114
4.6	The 12 Experiential Mode Work Sessions	195
4.6.1	Maladaptive Coping Mode sessions (EMW 1&7)	195
4.6.2	Dysfunctional Parent Mode sessions (EMW 2&8)	213
4.6.3	Vulnerable Child Mode sessions (EMW 3&9)	227
4.6.4	Angry/ Impulsive Child Mode sessions (EMW-ACM 4&10)	240
4.6.5	Happy Child Mode sessions (EMW 5&11)	247
4.6.6	Healthy Adult Mode sessions (EMW 6&12)	255
5	The Individual Schema Therapy Sessions	263
5.1	The Schema Therapy Conceptualization and Goals	265
5.2	The Maladaptive Coping Modes (MCM)	267
5.2.1	Schema Therapy behavioral pattern-breaking interventions	267
5.2.2	Schema Therapy cognitive interventions	275
5.2.3	Schema Therapy experiential interventions	278
5.3	Dysfunctional Parent Modes (DyPMs)	280
5.3.1	Schema Therapy behavioral pattern-breaking interventions	280
5.3.2	Schema Therapy cognitive interventions (Handout: IST-DyPM3)	282
5.3.3	Schema Therapy experiential interventions	288
5.4	The Vulnerable Child Mode	292
5.4.1	Schema Therapy behavioral pattern-breaking interventions	292
5.4.2	Schema Therapy cognitive interventions	295
5.4.3	Schema Therapy experiential interventions	299
5.5	The Angry or Impulsive Child Mode (ACM, ICM)	302
5.5.1	Schema Therapy behavioral pattern-breaking interventions	302
5.5.2	Schema Therapy cognitive interventions	306
5.5.3	Schema Therapy experiential interventions	309
5.6	The Happy Child Mode	311
5.6.1	Schema Therapy behavioral pattern-breaking interventions	311
5.6.2	Schema Therapy cognitive interventions	313
5.6.3	Schema Therapy experiential interventions	315

5.7	The Healthy Adult Mode (HAM)	316
5.7.1	Schema Therapy behavioral pattern-breaking interventions	316
5.7.2	Schema Therapy cognitive interventions	317
5.7.3	Schema Therapy experiential interventions	319
6	Training, Supervision, Research, and Concluding Remarks	321
6.1	Training	321
6.2	Supervision	321
6.3	Research on Schema Therapy	323
6.4	Concluding Remarks	324
	References	326
	Index	328

List of Figures and Tables

Figures

2.1	Schema Therapy: Etiology of psychological disorders	6
4.1	Mode role-play of schema modes in action. Scene 1	202
4.2	Mode role-play of schema modes in action. Scene 2	204
4.3	Mode role-play of schema modes in action. Scene 3	206

Tables

2.1	Schemas organized by content area	6
2.2	Basic schema modes	7
2.3	Suggested therapist interventions in relationship to patient modes and needs	12
2.4	Models of group therapy	17
3.1	Integrated Schema Therapy Program by session: group and individual	22
3.2	Treatment components, goals and focus, patient materials list	32
3.3	Formats and lengths for the Integrated Schema Therapy treatment program	34
3.4	Schema therapist tool kit	46
4.1	List of group handouts, exercises, and assignments by session, mode, and location	48
5.1	Patient materials for Individual Schema Therapy sessions	264

About the Authors

Joan M. Farrell, Ph.D. is a Clinical Psychologist, Research and Training Director of the Center for Borderline Personality Disorder Treatment and Research, Indiana University School of Medicine, USA, and Adjunct Professor in the Department of Psychology, Indiana University-Purdue University Indianapolis (IUPUI). She is an International Society for Schema Therapy (ISST) Certified Schema Therapy Trainer and Supervisor, the Coordinator for Training & Certification of the ISST, and an Executive Board member.

Neele Reiss is a Researcher, Clinical Psychologist, and Schema Therapist in the Department of Differential Psychology and Psychological Diagnostics, Goethe-University Frankfurt, Germany, and a Director of the Institute for Psychotherapy in Mainz, Germany. She is an International Society for Schema Therapy (ISST) Certified Schema Therapy Trainer and Supervisor in Adult and Group.

Ida A. Shaw, M.A. is the Director of the Schema Therapy Institute Midwest, Indianapolis, USA, and Senior Clinical Supervisor at the Center for BPD Treatment and Research, Indiana University School of Medicine. She is an ISST Certified Schema Therapy Trainer and Supervisor in Adult, Child-Adolescent and Group.

Foreword

I am very pleased to have been invited to write the foreword for this groundbreaking resource, which will allow clinicians to integrate Individual and Group Schema Therapy (GST) sessions into complete treatment programs that can be delivered over a range of patient populations, treatment lengths, and levels of care.

Since I first heard about the extremely positive results of the authors' randomized controlled trial of GST for patients with borderline personality disorder (BPD) in 2008, I have been very excited about the potential of the group model to make Schema Therapy more available and affordable for patients. Given the worsening climate for mental health reimbursement in this era of managed care in the United States and elsewhere in the world, Group Schema Therapy has the potential to deliver the powerful treatment strategies of the schema approach in a more cost-effective manner than has been possible with individual schema therapy – with equivalent or perhaps superior results. I am especially excited about the large-scale clinical trial that is under way at 14 sites in six different countries. Arnoud Arntz and Joan Farrell serve as the co-principal investigators of the study, testing the efficacy and cost-effectiveness of the Group Schema Therapy model for BPD patients combined with two different numbers of individual sessions.

This book, with the collaboration of Neele Reiss, the psychologist and schema therapist who pioneered GST in Germany, extends the integrated Individual and Group Schema Therapy program for use with a wide range of patient groups – personality disorder, complex trauma, chronic difficulties, those who have failed in other treatments, and those who need a higher level of care. It includes the novel concept of giving patients a “bank account” of individual ST sessions to draw on, as they need to; this concept was developed for the BPD multisite trial. Since the publication of the first book on GST (Farrell & Shaw, 2012), there have been many initiatives internationally to apply the GST model to other diagnostic groups of patients. Like individual ST, GST is trans-diagnostic – in the sense that the focus of interventions is the mode profile of the patient rather than specific symptoms. Thus, GST,

like ST, should be effective for disorders other than BPD. Of course, any application needs empirical validation and I am happy to say that preliminary findings are promising. This integrated program has been tested in inpatient and day hospital settings for BPD (Reiss *et al.*, 2013a) and mixed Cluster B and C disorders (Muste, 2012; Fuhrhans, 2012). Its use in forensic settings is being evaluated in the UK. A randomized controlled trial comparing ST to cognitive-behavioral therapy (CBT) for avoidant PD and social phobia and an experimental case series for dissociative disorders are under way in the Netherlands.

The integration of Group and Individual Schema Therapy presented here feels entirely consistent with my own individual model, in terms of the conceptual model, therapeutic alliance, and treatment interventions. The GST model encourages group members to become like a healthy family in which they can “re-family” each other, under the watchful guidance of two highly skilled therapist-parents. The sense of belonging and acceptance provided by this group analogue of a loving family seems to catalyze both the limited re-parenting and emotion-focused components of ST. Furthermore, by utilizing two co-therapists for each group, GST has found a way to free up one therapist to move fluidly around the group, often working with one or two members at a time, creating novel experiential exercises to bring about change. At the same time, the second therapist serves as the “stable base” for the rest of the group, maintains an ongoing emotional connection with each member, monitors the reactions of all members, explains what is happening to educate them about what is taking place, and intervenes to shift the direction of the group to focus on the needs of other group members. I am also impressed that GST goes well beyond the traditional CBT/DBT (dialectical behavioral therapy) group format, in which members are taught skills in a seminar-like setting; and beyond non-CBT groups, in which the therapist does individual work with one member while the rest of the group primarily watches. In GST, the techniques used in Individual Schema Therapy, like imagery change work and mode role-plays, have been adapted to engage all of the members in unique exercises that make use of the power of group interaction and support. These group therapeutic factors, combined with the broad range of integrative techniques that are already part of ST, may account for the large treatment effects in the controlled outcome study I mentioned earlier, as well as in preliminary data from other ongoing studies of GST.

The authors describe a systematic approach to ST treatment, while retaining the flexibility that I have always valued so highly in developing Individual Schema Therapy. The treatment suggestions are specific and well organized, while avoiding the temptation to write a therapeutic

“cookbook” for therapists to follow in a rote manner. The authors have preserved the core elements of ST by developing “limited reparenting” intervention strategies for each mode that arises, seizing “experiential moments” to do emotion-focused work that brings about change at a deep level. Like individual ST, their group model blends experiential, cognitive, interpersonal, and behavioral work. The program presented here divides ST interventions into four main components: Schema Therapy Education, Mode Awareness, Mode Management, and Experiential Mode Work. There are sessions of each component that target the main schema modes one at a time. The individual and group sessions are coordinated by mode. The individual ST sessions give therapists the option of a cognitive, experiential, or behavioral pattern-breaking intervention. The authors provide specific therapist script examples that allow access for the beginner using ST, yet have the essential flexibility of ST so that they can be adapted to meet an individual patient’s mode and need. The combination of structure and flexibility in this manual make it accessible to practitioners at all levels of experience with ST. The book is written at a level that should appeal to a very broad range of mental health professionals, including psychologists, social workers, psychiatrists, counselors, and psychiatric nurses, as well as interns and residents.

The experience that the authors have gained over 30 years of training therapists throughout the world, and of leading GST groups with a broad range of clinical populations, is evident throughout the volume. This book is the first published treatment manual for integrating Individual and Group Schema Therapy, and succeeds in providing the most essential information that clinicians will need in order to develop and implement such programs. The program sessions can be delivered in intensive form for higher levels of care, such as inpatient and day therapy, or spread over a year of outpatient treatment. The program could begin with more sessions per week, then taper in intensity and be moved to outpatient care. The user-friendly format of the book includes patient examples, descriptions of group and individual sessions, and examples of therapist scripts to explain core ST concepts in language easily understood by patients, along with patient handouts, exercises, and assignments. This material is presented in the manual itself as well as being available in downloadable form on the Wiley website for use with patients.

On a more personal level, I had the opportunity to experience GST first hand as a participant at an advanced training workshop that I invited Joan and Ida with Neele to teach for the senior schema therapists at our New York institute. I am even more excited about the potential of ST in a group after this experience, and would love to conduct an ST group like this

myself once I have learned the necessary skills. Joan Farrell is an outstanding schema therapist who serves as the “stable base,” emotional center, and “educator” for the group as a whole – a role I can imagine myself learning to fill, given enough time and experience. What truly amazed me – perhaps because her style is so different from mine and Joan’s – was the remarkable group work of Ida Shaw. It is hard to convey the level of originality, creativity, and spontaneity she brings to the group experience. She is able to blend elements of gestalt, psychodrama, role-playing, and her own infectious style of play into an approach that perfectly fits the intensive demands of schema mode work, cajoling patients to change in profound ways. The group exercises in this manual will allow schema therapists to try out some of her unique work. Neele Reiss adds the perspective of the “next generation” of schema therapists who are intent upon practicing and empirically validating ST. She collaborated on inpatient studies of the integrated model for BPD patients (Reiss *et al.*, 2013a) and her current work applies ST interventions to problems like test phobia and eating disorders.

I highly recommend this outstanding manual to all mental health professionals working with more complex, chronic, and hard-to-treat patient populations – especially those who are looking for an evidence-based, cost-effective alternative to existing therapies. This book is essential reading for professionals interested in schema therapy, BPD and other personality disorders, group therapy, and new approaches to expanding CBT.

Jeffrey Young, PhD
Schema Therapy Institute of New York
Columbia University, Dept. of Psychiatry
March 2014

Acknowledgments

This book is the culmination of our collaboration with Neele Reiss that began in 2008 when we discovered at the International Society of Schema Therapy Conference in Portugal that someone else in the world was doing Schema Therapy in groups. We have had hours of discussion, fun, and ice cream since then and the development of an important friendship which we treasure. We thank Arnoud Arntz for his mentorship, personal support, and friendship; Jeff Young for his thought-provoking discussions, support, and encouragement and Wendy Behary for commiserating about the arduous task of writing and for humor at needed moments.

The therapists who trained with us made important contributions to this treatment manual as teaching them forced us to make explicit and clear the way we practice Group Schema Therapy. Most of all we thank our patients, who taught us what we needed to understand about their needs and struggles and what worked to help them.

Special thanks from Joan to Elke and Siegbert Reiss for their warm hospitality, lovely meals, and fine wine during a heavy writing period as well as German history lessons and sightseeing during much needed breaks.

Joan Farrell and Ida Shaw

Writing this book with my close friends Joan and Ida has been a wonderful journey with many new discoveries. I want to thank you for the collaboration and the warm friendship over the past years.

I would also like to thank Friederike Vogel, who has been a very supportive colleague ever since we started working together and has become a dear friend to me. Furthermore I would like to thank all my patients – without them I would never have learned to practice Schema Therapy and Group Schema Therapy.

Last but not least, I would like to thank my family, especially my father Siegbert and my mother Elke, and my best friend Steffi – for your patience, your advice, for grounding me whenever needed, and for your unconditional support at all times. I know I can count on you.

Neele Reiss

About the Companion Web Site

A companion web site is available for this book at:



www.wiley.com/go/farrell

Visit the site to download the patient materials for your personal use.

Introduction

This manual presents an integrated Individual and Group Schema Therapy Program that is transdiagnostic, designed to be implemented in a variety of treatment settings and in programs of varied length. It is written for psychotherapists in a practical manner with a clinical focus. Sample therapist scripts, detailed session instructions, and handouts for each individual and group session are provided.

1.1 Background

Schema Therapy (ST), developed originally for individual psychotherapy by Young (1990; Young, Klosko, & Weishaar, 2003), is an approach to the treatment of a wide variety of psychological disorders that integrates cognitive, experiential, or emotion-focused and behavioral pattern-breaking interventions. It is a comprehensive model that strategically integrates aspects of other approaches but remains unique. A group version of ST was developed (Farrell & Shaw, 2012; Reiss & Vogel, 2010; Muste, Weertman, & Claassen, 2009). The effectiveness of individual (IST) and group (GST) therapy for borderline personality disorder (BPD) has been validated empirically (Giesen-Bloo *et al.*, 2006; Farrell, Shaw, & Webber, 2009; Reiss, Lieb, Arntz, Shaw, & Farrell, 2013a; Nadort *et al.*, 2009). The model of ST is transdiagnostic. The use of ST for avoidant personality disorder (PD), social anxiety, eating disorders, post-traumatic stress disorder, narcissism, anti-social PD, substance abuse, and psychopathy is being explored and evaluated internationally. It is an approach that is rated positively by both patients and therapists (Spinhoven, Giesen-Bloo, van Dyck, Kooiman, & Arntz, 2007). In addition, ST has growing evidence of its cost-effectiveness for the individual modality (IST) (Giesen-Bloo *et al.*, 2006; Nadort *et al.*, 2009). However, the limited availability of specialized individual psychotherapies,

including ST, in the current healthcare economy has prevented more wide-scale use of IST in clinical settings. An integrated program combining multiple GST sessions with a limited number of IST sessions in a structured program has been evaluated, with promising results (Muste, Weertman, & Claassen, 2009; Reiss *et al.*, 2013a). Combining more frequent group ST sessions with strategically planned individual sessions has the potential for making this promising treatment more widely available in a larger number of settings.

The concept of an integrated and structured program was originally developed for use with severe patients in hospital settings (Reiss *et al.*, 2013a), but it can be implemented across a wide range of patient populations, treatment settings, and treatment lengths. One patient population for whom it is ideal are those with personality disorders or PD features, those with chronic or complex psychological problems, those with multiple trauma histories, and those for whom other treatment approaches have failed. In general, these are patients who are treated at the higher levels of care and whose life potential is tragically not realized in the quality of their lives. An advantage of Schema Therapy is that it approaches treatment by targeting maladaptive schema modes rather than specific symptoms or disorders, thus transcending psychiatric diagnoses and impending changes in diagnostic classification. This program can be implemented in inpatient, day therapy, and intensive outpatient settings, as well as in general outpatient treatment with varying session frequency. A program of intensive sessions as the beginning of outpatient treatment may act to “jump-start” the treatment of patients with entrenched maladaptive coping strategies that produce the severe and sometimes life-threatening symptoms of disorders like BPD. It is possible to use all the handouts provided throughout this book within the context of individual ST treatment or selectively within other psychotherapy models. For example, the experiential interventions can augment cognitive therapy, filling a gap in that approach to treating patients with personality disorders and complex trauma.

The program of the manual is theoretically consistent with individual ST (Young *et al.*, 2003) and group ST (Farrell & Shaw, 2012). Like most approaches to psychotherapy that go beyond skills training, ST requires specialized training to meet adherence and competence standards. This manual was designed to make IST, GST, and their integrated combination accessible for a wide range of psychotherapists of multiple theoretical orientations, including those working in intensive settings like inpatient and day therapy programs. The manual provides a detailed step-by-step guide for an ST fundamentals program that integrates IST and GST sessions. The program includes the core components of ST: limited reparenting,

education about the ST model, mode awareness, mode management and cognitive, experiential, and behavioral pattern-breaking change work. It can be used across diagnostic groups. There are 12 individual sessions and 42 group sessions. Each session is described in terms of goals, therapist interventions, tips on management, sample session scripts for therapists to adapt, and corresponding information handouts, ST exercises, and therapy assignments. The individual sessions are designed to be complementary to the group work and the schema mode being focused on, while still allowing individual needs to be met. How to balance individual and group focus is discussed throughout. This level of detail will allow therapists new to Schema Therapy to conduct sessions with confidence and to coordinate group work effectively with individual work. It will also allow more experienced individual schema therapists to begin leading ST groups effectively. The program is not a rigid protocol, but rather a framework combining flexibility with standardization. The structured yet flexible format serves a number of purposes: the manual can be used as a detailed plan for implementing a structured, integrated program of individual and group ST in higher levels of care settings with multiple sessions per week; as a research protocol for psychotherapy outcome studies; or clinicians can select individual or group sessions, or the combination, to work on specific schema modes. Therapists can choose whether to implement the program as a whole, or to select individual sessions, group sessions, or a combination of both to suit their group and its needs.

1.2 The Chapters

Chapter 2 presents the basic concepts of the ST model originally developed for individuals by Young (Young *et al.*, 2003) and the adaptation for groups developed by Farrell and Shaw (2012). The goals of ST, the therapist approach of limited reparenting, and the core components of the model are described in this chapter. The approach of the integrated ST program to combining individual and group sessions is discussed. The general course and stages of the treatment program are outlined here. Chapter 3 describes some of the essentials of the treatment program: the patients, with inclusion and exclusion factors; the therapists; the milieu – both the physical setting and the multidisciplinary treatment team; the length of sessions and treatment; and the possible schedules and format for delivering the program in different settings. Chapter 4 presents the group sessions by component, with sample therapist scripts and the patient materials – handouts, group exercises, and session assignments. Chapter 5 presents the 12 individual sessions – with specific content, sample scripts, and handouts. Chapter 6

describes the ST training and supervision recommended for therapists. The research that has been conducted is summarized, with a brief description of the investigations that are in progress at the time of writing.

This book is unique in the ST literature as its goal is to provide an implementable program in its entirety. It is not focused on a particular disorder. It is not limited to one of the modalities of treatment delivery; it covers both individual and group ST. The reader is referred to the ST volumes in the References for theory and focus on specific disorders.

The Basics of Schema Therapy

2.1 The Theoretical Model

The model presented in this manual is consistent with the theory, components of treatment, and goals presented by Young, Klosko, and Weishaar (2003). The Schema Therapy (ST) model is summarized here and the reader is referred to that volume for additional elaboration of the individual ST model and its application. ST grew out of efforts by Young *et al.* (2003) to treat patients with personality disorders more effectively, and also those who either did not respond to traditional cognitive therapy or relapsed. ST is based upon a unifying theory and a structured and systematic approach. Because ST is an integrative treatment, there is overlap with other psychotherapy models such as cognitive and psychodynamic psychotherapy, object relations theory, and Gestalt psychotherapy, but total overlap with no other model.

Figure 2.1 summarizes the model for the etiology of psychopathology posited by ST. When the normal, healthy developmental needs of childhood are not met, maladaptive schemas develop. Maladaptive schemas are psychological constructs that include beliefs we have about ourselves, the world, and other people, which result from interactions of unmet core childhood needs, innate temperament, and early environment. ST views this interaction in terms of a plasticity or differential susceptibility model. Schemas are comprised of memories, bodily sensations, emotions, and cognitions that originate in childhood and are elaborated through a person's lifetime. These schemas often have an adaptive role in childhood (e.g., in terms of survival in an abusive situation – it engenders more hope for children if they believe they are defective as opposed to the adult being defective). By adulthood, maladaptive schemas are inaccurate, dysfunctional, and limiting, although strongly held and frequently not in the person's conscious awareness. Young (1990; Young *et al.*, 2003) identified 18 early maladaptive schemas




Figure 2.1 Schema Therapy: Etiology of psychological disorders.

Table 2.1 Schemas organized by content area

<i>Disconnection and rejection</i>	<i>Impaired autonomy and performance</i>
Mistrust/Abuse	Dependence/Incompetence
Emotional Deprivation	Vulnerability to Harm/Illness
Defectiveness/Shame	Enmeshment/Undeveloped Self
Social Isolation/Alienation	Abandonment/Instability
Emotional Inhibition	Subjugation
	Failure
<i>Impaired limits</i>	<i>Exaggerated expectations</i>
Entitlement	Self-Sacrifice
Insufficient Self-Control	Unrelenting Standards
<i>Other</i>	
Negativity/Pessimism	Approval-Seeking/Recognition-Seeking
Punitiveness	

Table 2.2 Basic schema modes

Innate Child Modes	Vulnerable Child Angry/Impulsive Child	Innate responses to unmet needs
Maladaptive Coping Modes	Avoidant Protector Overcompensator Compliant Surrenderer	Overused survival responses to trauma or unmet needs – flight, fight, and freeze
Dysfunctional Parent Modes	Punitive Parent Demanding Parent	Selective internalization of negative aspects of early others
Healthy Modes	Happy Child Healthy Adult	Underdeveloped

(EMS) in patients with personality disorders (Table 2.1). The schemas are defined individually in the patient materials for the ST Education group (ST-Education Sessions 1–5).

When maladaptive schemas are activated, intense states occur, described in ST as “schema modes.” A schema mode is defined as the current emotional, cognitive, and behavioral state that a person is in. Dysfunctional modes occur most frequently when multiple maladaptive schemas are activated. Four basic categories of modes are defined (Table 2.2).

Primary or innate child modes (Vulnerable Child, Angry Child, Impulsive Child) are said to develop when basic emotional needs in childhood (such as safety, nurturance, or autonomy) are not adequately met. These “child modes” are defined by intense feelings such as fear, helplessness, or rage, and involve the innate reactions a child has. Dysfunctional Parent Modes (Punitive Parent or Demanding Parent) comprise the second category of modes. Dysfunctional Parent Modes reflect the selective internalization of negative aspects of attachment figures (e.g., parents, teachers, peers, etc.) during childhood and adolescence. Maladaptive Coping Modes, a third category of mode, are defined by an overuse of unhealthy coping styles (fight – overcompensation; flight – avoidance; or freeze – surrender). All have the goal of protecting the person from experiencing pain, anxiety, or fear. These Maladaptive Coping Modes operate outside of conscious awareness and a goal of ST is that patients become aware of their coping styles and learn healthier, more adaptive coping responses. Maladaptive Coping Modes incorporate the concept of defense mechanisms, a concept previously missing in cognitive therapy, and allow for a better understanding of personality disorders by clinicians and patients. The overcompensating coping style contains modes in which a person acts in opposition to the

schema or schemas that are triggered. One example is the Bully-Attack Mode in which the person experiencing the pain resulting from a schema retaliates by causing pain. The avoidant coping style involves physical, psychological, and social withdrawal and avoidance. Avoidant Modes include the Detached Protector, a mode that ranges from a person being “spacey,” or briefly losing focus in an interaction, to severe dissociation. This mode is very commonly present when patients enter therapy as it operates to protect the Vulnerable Child Mode from overwhelming fear or painful feelings. Surrender is the third coping style and it represents giving in or giving up to the schema present. For example, if the triggering schema is defectiveness, a surrender response would be to accept that you are defective and behave accordingly: never taking on challenges, working not to be exposed as incompetent.

In a fourth category, Healthy and Functional Modes, the Healthy Adult Mode and Happy Child Mode are found. The Healthy Adult Mode includes functional thoughts and behaviors, and the skills needed to function in adult life. The Happy Child Mode is a resource for playful and enjoyable activities, especially in social networks. Many patients were neither allowed nor encouraged to play, thus missing opportunities to explore their likes and dislikes and take part in our earliest social interactions with peers. The healthy modes tend to be severely underdeveloped in patients with personality disorders or features.

Modes are often triggered by events that patients experience as highly emotional. Modes can switch rapidly in patients suffering from severe personality disorders, resulting in the sudden changes in behavior or seemingly disproportionate reactions that are one source of patients’ interpersonal difficulties and emotional and behavioral instability. Modes can also stay rigidly entrenched, as is the case with many avoidant patients. Common negative coping responses – aggression, hostility, manipulation, exploitation, dominance, recognition-seeking, stimulation-seeking, impulsivity, substance abuse, compliance, dependence, excessive self-reliance, compulsivity, inhibition, psychological withdrawal, social isolation, and situational and emotional avoidance – can be understood in mode terms.

Symptoms of personality disorder can be described and understood in terms of the operation of modes. One example is the ST conceptualization of borderline personality disorder (BPD). Abandonment fears describe the emotional state of the Vulnerable Child Mode. Intense anger, at times accompanied by uncontrolled expressions of anger, occurs in the Angry Child and Impulsive Child Modes. The Impulsive Child Mode fuels action that is potentially damaging as well as being one source of self-injurious behavior. The Dysfunctional Parent Modes are another source of self-injurious