

Lori A. Erickson

# Atlas of Endocrine Pathology



 Springer

---

# Atlas of Anatomic Pathology

*Series Editor*  
Liang Cheng

---

For further volumes:  
<http://www.springer.com/series/10144>



---

Lori A. Erickson

# Atlas of Endocrine Pathology

 Springer

Lori A. Erickson, MD  
Mayo Clinic College of Medicine  
Rochester, MN  
USA

ISBN 978-1-4939-0442-6      ISBN 978-1-4939-0443-3 (eBook)  
DOI 10.1007/978-1-4939-0443-3  
Springer New York Heidelberg Dordrecht London

Library of Congress Control Number: 2014935862

© Springer Science+Business Media New York 2014

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media ([www.springer.com](http://www.springer.com))

---

## Series Preface

*One Picture is Worth Ten Thousand Words*

– Frederick Barnard, 1927

Remarkable progress has been made in anatomic and surgical pathology during the last 10 years. The ability of surgical pathologists to reach a definite diagnosis is now enhanced by immunohistochemical and molecular techniques. Many new clinically important histopathologic entities and variants have been described using these techniques. Established diagnostic entities are more fully defined for virtually every organ system. The emergence of personalized medicine has also created a paradigm shift in surgical pathology. Both promptness and precision are required of modern pathologists. Newer diagnostic tests in anatomic pathology, however, cannot benefit the patient unless the pathologist recognizes the lesion and requests the necessary special studies. An up-to-date Atlas encompassing the full spectrum of benign and malignant lesions, their variants, and evidence-based diagnostic criteria for each organ system is needed. This Atlas is not intended as a comprehensive source of detailed clinical information concerning the entities shown. Clinical and therapeutic guidelines are served admirably by a large number of excellent textbooks. This Atlas, however, is intended as a “first knowledge base” in the quest for definitive and efficient diagnosis of both usual and unusual diseases.

The *Atlas of Anatomic Pathology* is presented to the reader as a quick reference guide for diagnosis and classification of benign, congenital, inflammatory, nonneoplastic, and neoplastic lesions organized by organ systems. Normal and variations of “normal” histology are illustrated for each organ. The Atlas focuses on visual diagnostic criteria and differential diagnosis. The organization is intended to provide quick access to images and confirmatory tests for each specific organ or site. The Atlas adopts the well-known and widely accepted terminology, nomenclature, classification schemes, and staging algorithms.

This book Series is intended chiefly for use by pathologists in training and practicing surgical pathologists in their daily practice. It is also a useful resource for medical students, cyto-technologists, pathologist assistants, and other medical professionals with special interest in anatomic pathology. We hope that our trainees, students, and readers at all levels of expertise will learn, understand, and gain insight into the pathophysiology of disease processes through this comprehensive resource. Macroscopic and histological images are aesthetically pleasing in many ways. We hope that the new Series will serve as a virtual pathology museum for the edification of our readers.

Indianapolis, IN, USA

Liang Cheng, MD



---

## Preface

The *Atlas of Endocrine Pathology* provides a comprehensive compendium of photomicrographs of common and uncommon entities in endocrine pathology. Histologic features of normal features, reactive conditions, hyperplasia, and tumors are included. The most helpful diagnostic features are illustrated to provide direction and clues to the diagnosis of endocrine tumors. The photomicrographs highlight the most pertinent diagnostic features in problematic diagnoses in endocrine pathology. This Atlas serves as a learning tool for those becoming familiar with the diverse entities encountered in endocrine pathology and a reference for practicing pathologists faced with challenging diagnoses in endocrine pathology.

Rochester, MN, USA

Lori A. Erickson, MD





---

# Contents

<b>1 Thyroid Histology</b> .....	1
Normal Thyroid .....	2
Solid Cell Nests .....	4
Radiation Changes .....	5
Minocycline and Amiodarone Thyroid .....	5
Palpation Thyroiditis .....	7
Thyroid Needle-Biopsy Site .....	7
Ectopic Thyroid Tissue .....	8
Thyroglossal Duct Cyst .....	9
Struma Ovarii .....	9
References .....	10
<b>2 Thyroiditis</b> .....	13
Acute Thyroiditis .....	14
Granulomatous (de Quervain) Thyroiditis .....	14
Fibrous (Riedel) Thyroiditis .....	16
Hashimoto Thyroiditis .....	17
Focal Lymphocytic Thyroiditis .....	18
References .....	19
<b>3 Graves Disease (Diffuse Hyperplasia)</b> .....	21
References .....	24
<b>4 Thyroid Goiter</b> .....	25
Multinodular Goiter .....	26
Dyshormonogenetic Goiter .....	27
Amyloid Goiter .....	28
References .....	29
<b>5 Papillary Thyroid Carcinoma</b> .....	31
Classic PTC .....	32
Papillary Thyroid Microcarcinoma .....	36
FVPTC .....	37
Cribriform-Morular Variant of PTC .....	40
Warthin-Like Variant of PTC .....	42
Diffuse Sclerosing Variant of PTC .....	43
Oxyphilic (Oncocytic/Hurthle Cell) Variant of PTC .....	44
Solid Variant of PTC .....	45
Tall Cell Variant of PTC .....	46
Columnar Variant of PTC .....	47
Hobnail Variant of PTC .....	48
References .....	49
<b>6 Hyalinizing Trabecular Tumor</b> .....	51
References .....	53

<b>7 Follicular Thyroid Neoplasms</b> .....	55
Follicular Adenoma .....	56
Follicular Carcinoma .....	60
References .....	62
<b>8 Hurthle Cell Thyroid Neoplasms</b> .....	63
Hurthle Cell Adenoma .....	64
Hurthle Cell Carcinoma .....	65
References .....	66
<b>9 Poorly Differentiated Thyroid Carcinoma</b> .....	67
References .....	69
<b>10 Anaplastic Thyroid Carcinoma</b> .....	71
References .....	75
<b>11 Tumors of C Cells</b> .....	77
C-Cell Hyperplasia .....	78
Medullary Thyroid Microcarcinoma .....	78
Medullary Thyroid Carcinoma .....	79
References .....	84
<b>12 Unusual Thyroid Tumors</b> .....	87
Sclerosing Mucoepidermoid Carcinoma with Eosinophils .....	87
Squamous Cell Carcinoma .....	89
Lymphoma .....	90
Langerhans Cell Histiocytosis .....	91
Solitary Fibrous Tumor .....	91
Angiosarcoma .....	92
Synovial Sarcoma .....	93
Low-Grade Fibromyxoid Sarcoma .....	93
Rhabdomyosarcoma .....	94
Carcinoma Showing Thymus-Like Differentiation .....	94
Teratoma .....	95
Paraganglioma .....	96
References .....	96
<b>13 Metastases to Thyroid</b> .....	99
References .....	101
<b>14 Parathyroid Histology</b> .....	103
Normal Parathyroid Histology .....	104
Parathyroid Cysts .....	105
Parathyroiditis .....	106
References .....	107
<b>15 Parathyroid Hyperplasia</b> .....	109
Primary Parathyroid Hyperplasia .....	110
Primary Clear Cell Hyperplasia .....	113
Lipoadenomatous Hyperplasia .....	114
Secondary Parathyroid Hyperplasia .....	115
Tertiary Parathyroid Hyperplasia .....	118
References .....	119
<b>16 Parathyroid Adenoma</b> .....	121
Chief Cell Parathyroid Adenoma .....	122
Oxyphil Cell Parathyroid Adenoma .....	125

Clear Cell Parathyroid Adenoma . . . . .	126
Parathyroid Lipoadenoma . . . . .	127
Atypical Parathyroid Adenoma . . . . .	128
References . . . . .	130
<b>17 Parathyroid Carcinoma . . . . .</b>	<b>131</b>
References . . . . .	138
<b>18 Parathyromatosis . . . . .</b>	<b>139</b>
References . . . . .	141
<b>19 Metastases to Parathyroid . . . . .</b>	<b>143</b>
References . . . . .	145
<b>20 Adrenal Gland Histology . . . . .</b>	<b>147</b>
Normal Adrenal Histology . . . . .	148
Heterotopia of Adrenal Tissue. . . . .	152
Adrenal Infection. . . . .	153
References . . . . .	153
<b>21 Adrenal Cysts . . . . .</b>	<b>155</b>
References . . . . .	157
<b>22 Adrenal Cortical Hyperplasia . . . . .</b>	<b>159</b>
Adrenal Cortical Hyperplasia, Diffuse . . . . .	160
Adrenal Cortical Hyperplasia, Nodular. . . . .	161
Primary Pigmented Nodular Adrenocortical Disease . . . . .	162
References . . . . .	164
<b>23 Adrenal Cortical Adenoma . . . . .</b>	<b>165</b>
References . . . . .	172
<b>24 Adrenal Cortical Carcinoma . . . . .</b>	<b>173</b>
References . . . . .	182
<b>25 Adrenal Myelolipoma . . . . .</b>	<b>183</b>
References . . . . .	185
<b>26 Pheochromocytoma and Adrenal Medullary Hyperplasia . . . . .</b>	<b>187</b>
Adrenal Medullary Hyperplasia . . . . .	188
Pheochromocytoma. . . . .	189
Composite Pheochromocytoma. . . . .	194
References . . . . .	195
<b>27 Ganglioneuroma and Neuroblastoma . . . . .</b>	<b>197</b>
References . . . . .	200
<b>28 Unusual Adrenal Tumors . . . . .</b>	<b>201</b>
Angiosarcoma of the Adrenal Gland. . . . .	202
Leiomyosarcoma of the Adrenal Gland . . . . .	203
Schwannoma of the Adrenal Gland. . . . .	203
Teratoma of the Adrenal Gland . . . . .	204
Adenomatoid Tumor of the Adrenal Gland. . . . .	204
References . . . . .	205
<b>29 Metastases to the Adrenal Gland . . . . .</b>	<b>207</b>
References . . . . .	209
<b>Index . . . . .</b>	<b>211</b>

The thyroid gland weighs 15–25 g and is composed of two lobes joined by the isthmus, with approximately 40 % of people having a pyramidal lobe [1]. The thyroid is composed of lobules, each of which is composed of 20–40 follicles. Each follicle is surrounded by a basement membrane and lined by follicular cells. Colloid is present in the lumen of the follicles. The presence of calcium oxalate crystals in the colloid is a helpful feature differentiating it from parathyroid. Thyrocytes also may be distinguished from parathyroid cells by reactivity to thyroid transcription factor 1 (TTF1) and thyroglobulin and the absence of parathyroid hormone and the neuroendocrine markers chromogranin and synaptophysin in thyrocytes. C cells are calcitonin-producing cells in the thyroid. They are present in interfollicular areas but are difficult to identify in normal thyroid tissue with hematoxylin and eosin (H&E) alone. The immunohistochemical stains calcitonin, chromogranin, synaptophysin, somatostatin, calcitonin gene-related peptide, and bombesin will highlight these neuroendocrine cells of the thyroid. Thyroid follicular cells produce thyroxine (T4) and triiodothyronine (T3), and C cells produce calcitonin. C cells migrate from the neural crest to the

ultimobranchial bodies, which are derived from branchial pouch complexes IV and V and develop in the first five to seven fetal weeks [2, 3]. C cells have clear cytoplasm and oval to round nuclei; they are positive for calcitonin, chromogranin, synaptophysin, calcitonin, calcitonin gene-related peptide, somatostatin, and bombesin [4]. Calcitonin mRNA has been localized in C cells by in situ hybridization studies [5]. Calcitonin is present in both 280-nm type I and 130-nm type II secretory granules; it is produced by C cells. Calcitonin interacts with bone, kidney, and the gastrointestinal tract to lower serum calcium [6]. Thyroid follicular cells and tumors are immunopositive for thyroglobulin, TTF1, and keratin. Thyroid C cells and tumors are immunopositive for calcitonin, chromogranin, synaptophysin, keratin (Cam5.2), and TTF1, although the intensity of TTF1 immunostaining often is less than in follicular thyroid cells and tumors. Interesting histologic features occasionally may be recognized in thyroid tissue, including solid cell nests, fatty metaplasia, radiation changes, drugs such as minocycline and amiodarone, and palpation thyroiditis. Thyroid tissue also may occur ectopically, and tumors may develop in this tissue.

## Normal Thyroid




**Fig. 1.1** Normal thyroid. Gross image of a normal thyroid gland showing the bilobed gland joined by the isthmus in the center. The isthmus lies anterior to the trachea and inferior to the cricoid cartilage. The thyroid is in front of the larynx and trachea. The thyroid descends with the thyroglossal duct into the neck and expands after the thyroglossal duct atrophies. The pyramidal lobe, present in approximately 40 % of people, is a vestige of the thyroglossal duct. The anlage is visible in the pharynx in association with the heart by day 17 of fetal life [7, 8]. Follicles develop from 9 to 12 fetal weeks, and well-developed colloid follicles are seen at 14 weeks' gestation [7, 8]




**Fig. 1.2** Normal thyroid. Gross image of a cut section of a normal thyroid gland, showing the bilobed gland joined by the isthmus in the center. Normal thyroid glands in adults weigh 15–25 g and may vary with constitutional factors, age, sex, size of the individual, and functional status. The thyroid may be larger in women and may change with hormonal effects such as pregnancy and the menstrual cycle [9, 10]. An enlarged thyroid gland is referred to as a goiter







**Fig. 1.3** Normal thyroid. The thyroid gland is made up of lobules of 20–40 follicles. The follicles have an average size of 200  $\mu\text{m}$  (range of 50–500  $\mu\text{m}$ ) and have colloid in their lumens, which is made up mostly of thyroglobulin. Calcium oxalate crystals are present in normal colloid and allow a distinction from colloid-like material in parathyroid tissue that lacks calcium oxalate [11]. Normal thyroid tissue may be present in soft tissues next to the thyroid as ectopic thyroid tissue [12]. The follicular cells have round nuclei with a diffuse chromatin pattern and amphophilic to eosinophilic cytoplasm. Thyroglobulin mRNA has been identified in follicular cells by in situ hybridization studies [13]. Follicular cells are positive for the immunohistochemical markers thyroglobulin, keratin, and TTF1



**Fig. 1.5** Normal thyroid. Immunopositivity for thyroglobulin is present in thyroid tissues and folliculogenic thyroid tumors. The thyrocytes stain with thyroglobulin, as does the colloid. Medullary thyroid tumors are negative for thyroglobulin. Thyroglobulin is helpful in identifying tissues and tumors as thyroid and differentiating them from tumors from other sites




**Fig. 1.4** Normal thyroid. Thyroid follicles are present with a central lumen containing colloid. Thyroid follicular cells are polygonal, with central nuclei and round nuclei with diffuse chromatin and moderate amounts of amphophilic to eosinophilic cytoplasm. The follicular cells produce thyroxine (T4) and triiodothyronine (T3) from exogenous iodine, which is oxidized and catalyzed by thyroid peroxidase. Thyroid-stimulating hormone (TSH) regulates the release of thyroglobulin, stored as colloid, endocytosed, and hydrolyzed with release of T3 and T4. Thyroglobulin breakdown and release of T3 and T4 are inhibited by iodine by inhibiting TSH stimulation of thyroid adenylate cyclase




**Fig. 1.6** Normal thyroid. Thyroid follicular cells are immunopositive for TTF1, which is helpful in differentiating thyroid from parathyroid and other tissues. TTF1 is not specific for thyroid follicular cells or folliculogenic tumors as it also stains, although with less intensity, medullary thyroid carcinoma. TTF1 also stains most adenocarcinomas of the lung and papillary adenocarcinoma of the sinuses. High-grade neuroendocrine tumors from the lung and some other sites may stain from TTF1. Thus, this marker is not specific for thyroid, although it is very helpful in determining the differentiation or primary site of difficult tumors, particularly when used as part of a panel of immunostains






**Fig. 1.7** Normal thyroid. This chromogranin immunostain highlights scattered C cells in interfollicular areas in normal thyroid. C cells of the thyroid are difficult to identify in normal thyroid tissue with H&E staining. C cells have clear cytoplasm and oval to round nuclei. They are positive for calcitonin, chromogranin, synaptophysin, calcitonin, calcitonin gene-related peptide, somatostatin, and bombesin [4]




**Fig. 1.8** Normal thyroid. Foci of fat cells can be seen in the thyroid parenchyma. Fat also may be seen in thyroid neoplasms. This finding of adipocytes in the thyroid is not known to be of pathologic significance

## Solid Cell Nests




**Fig. 1.9** Solid cell nests. Solid cell nests are small nests of oval cells with fine chromatin identified in 3 % of thyroid glands. They are thought to be ultimobranchial body rests [14]




**Fig. 1.10** Solid cell nests. Solid cell nests may be mistaken for papillary thyroid carcinomas (PTCs) and for squamous metaplasia. They stain for low molecular weight keratin and polyclonal carcinoembryonic antigen, and may show variable staining for calcitonin [15]



## Radiation Changes




**Fig. 1.11** Radiation changes. Radiation to the thyroid, neck, or tonsils may affect the thyroid. The effects depend on the dose and isotope [16–18]. Historical irradiation of the tonsils was associated with nodular hyperplasia of the thyroid. Radiation for acne also was associated with thyroid carcinoma [19]. More extensive irradiation, as with lymphoma treatment, is associated with increased cellularity, increased nodular size, fibrosis, chronic thyroiditis, and cytologic atypia [16]




**Fig. 1.12** Radiation changes. Acute radiation changes include hemorrhage, neutrophilic infiltration, necrosis, and follicular disruption [18]. Radiation may be associated with cytologic atypia, pleomorphism, and an increased incidence of thyroid carcinoma, predominantly PTCs [20, 21]. The latency may vary from decades to a short latency, as in childhood cancers after the Chernobyl accident [17, 22]. It is speculated that dietary iodine levels may have implications in radiation-associated thyroid carcinogenesis, and iodine deficiency might increase incidence, reduce latency, and influence tumor morphology and aggressiveness [23]


## Minocycline and Amiodarone Thyroid




**Fig. 1.13** Minocycline thyroid. Drugs may have various effects on the thyroid. Minocycline is a tetracycline antibiotic that produces a black pigment in the thyroid, which may be related to its interaction with thyroid peroxidase [24]. Minocycline pigmentation also may affect the skin, sclera, teeth, bone, heart valves, oral mucosa, and atherosclerotic plaques [25–30]. The antithyroid compounds thiocyanate (both in its chemical form and in cabbage and turnips) and perchlorate can inhibit thyroid iodine transport, whereas propothiouracil, methimazole, phenylbutazone, and lithium can inhibit binding reactions and thyroid hormone release [4, 31]




**Fig. 1.14** Minocycline thyroid. Minocycline pigment in the colloid and localized within the cytoplasm of thyroid follicular epithelial cells. Lipofuscin and hemosiderin also may localize in the cytoplasm but are light brown and brown yellow, respectively. Minocycline is dark brown/black



**Fig. 1.15** Minocycline thyroid. Minocycline is present in the follicular lumen and as pigmented granules in the follicular cells. In addition to pigments, other substances, such as calcium oxalate crystals, may be seen in colloid and are highlighted with polarization. Crystals are not specific for a particular disorder but are helpful in separating thyroid from parathyroid, which lacks calcium oxalate crystals




**Fig. 1.17** Amiodarone thyroid. Amiodarone is a class III antiarrhythmic agent associated with changes in thyroid function tests due to inhibition of 5'-deiodinase, resulting in decreased T3 generation from T4 with an associated increase in reverse T3 [33]. Amiodarone is associated with amiodarone-induced thyrotoxicosis related to excess iodine-induced thyroid hormone synthesis in an abnormal thyroid or is the result of destructive thyroiditis [33]. Amiodarone-induced hypothyroidism is thought to result from escape from the acute Wolff-Chaikoff effect, from defects in thyroid hormonogenesis, or from concomitant Hashimoto thyroiditis [33]




**Fig. 1.16** Amiodarone thyroid. Histologic features of amiodarone thyroid disease include involutional changes, degenerative and destructive follicular lesions, and zones of fibrosis [32]. Small groups of involuted follicles may have damage varying from changes in a few lining cells to total follicular destruction and follicular cell injury with swelling and granular or vacuolated cytoplasm [32]. The follicular damage and release of hormones may contribute to the associated thyrotoxicosis [32]



## Palpation Thyroiditis




**Fig. 1.18** Palpation thyroiditis. Multiple foci of small groups or single follicles are disrupted and associated with a mixed inflammatory infiltrate with prominent histiocytes, appearing as multiple small granulomatous foci in this case of palpation thyroiditis. Palpation thyroiditis is a common finding in surgically resected thyroids from palpation of the thyroid gland




**Fig. 1.19** Palpation thyroiditis. Palpation thyroiditis with disrupted follicles with mixed inflammatory infiltrate forming a small granulomatous focus. Palpation folliculitis is believed to be caused by traumatic rupture of isolated thyroid follicles as a result of palpation of the gland [34]

## Thyroid Needle-Biopsy Site




**Fig. 1.20** Thyroid needle-biopsy site. Shown is an area of fibrosis where prior fine-needle aspiration was performed. Needle-biopsy sites are not an uncommon finding in the thyroid. The fibrosis in these cases may entrap follicular cells and be mistaken for infiltrative growth. Inflammation and hemosiderin in these fibrous foci are helpful in identification




**Fig. 1.21** Thyroid needle biopsy site. Inflamed focus with associated fibrosis and histiocytic reaction in a fine-needle aspiration site. Cells associated with inflammation may have cytologic clearing and irregularity and should not be mistaken for PTC. Also, in follicular neoplasms, identifying the perpendicular and linear appearance of the needle-biopsy site with associated inflammation and hemosiderin prevents misinterpretation of capsular invasion




## Ectopic Thyroid Tissue



**Fig. 1.22** Ectopic thyroid tissue. This is an unusual example of ectopic thyroid tissue in the adrenal gland. Ectopic thyroid tissue has been reported from a variety of sites; however, historic reference to ectopic thyroid tissue involving lymph nodes now is thought most likely to represent metastatic PTC, particularly if lateral to the jugular vein [35–37]




**Fig. 1.23** Ectopic thyroid tissue. This ectopic thyroid tissue in the heart was identified along the left ventricular outflow tract. Ectopic thyroid tissue has been reported along the thyroglossal tract and in unusual locations such as the heart and pericardium, chest wall, porta hepatis, and vagina [38–40]




**Fig. 1.24** Ectopic thyroid tissue. This ectopic thyroid tissue is in the wall of the gallbladder. Although this is a rare location for ectopic thyroid tissue, it is important to recognize that ectopic thyroid tissue may occur in unusual locations and not to mistake it for another tissue type of a metastasis

## Thyroglossal Duct Cyst




**Fig. 1.25** Thyroglossal duct cyst. Thyroglossal duct cyst developmental anomalies due to a persistent connection between the foramen cecum and the thyroid gland. The cystic structures, lined by ciliated respiratory epithelium, may show squamous metaplasia, particularly if inflamed, and may have thyroid follicles in the wall. Complications include infections with fistula formation; rarely, carcinoma, usually papillary carcinoma, may develop in these lesions

## Struma Ovarii



**Fig. 1.26** Struma ovarii. Struma ovarii are uncommon mature teratomas in which thyroid tissue is present exclusively or predominantly. In this photomicrograph, the thyroid tissue is identified adjacent to ovarian tissue. No other components are present. Struma ovarii usually present in adults in the fifth or sixth decade as an ovarian mass and may have unusual clinical manifestations, such as hyperthyroidism, ascites, and Meigs syndrome [41]. Strumal carcinoids are uncommon, comprising approximately 1 % of ovarian tumors and 2.7 % of teratomas [41]



**Fig. 1.27** Struma ovarii. Struma ovarii with thyroid tissue adjacent to ovarian tissue. The thyroid tissue is present in follicles with eosinophilic colloid. The follicular cells are cuboid to slightly flattened cells with round nuclei and lack nuclear atypia. In this case, other tissue components are not present. When struma ovarii have a component of carcinoid tumor, they are referred to as strumal carcinoid. Carcinoid elements are not identified in this patient's tumor