

International Practices in Pancreatic Surgery

René Mantke · Hans Lippert
Markus W. Büchler · Michael G. Sarr
Editors

With contributions by
Adams, Andrén-Sandberg, Bassi, Beger, Bornman,
Farnell, Fernández-del Castillo, Fingerhut,
Gooszen, Imrie, Jaeck, Neoptolemos, Schmidt,
Takada, Tanaka, Warshaw

 Springer

International Practices in Pancreatic Surgery

René Mantke • Hans Lippert
Markus W. Büchler • Michael G. Sarr
Editors

International Practices in Pancreatic Surgery

Editors

René Mantke
Department of Surgery
City Hospital Brandenburg
Brandenburg
Germany

Markus W. Büchler
Department of Surgery
University of Heidelberg
Heidelberg
Germany

Hans Lippert
Department of Surgery
Otto von Guericke University
of Magdeburg
Magdeburg
Germany

Michael G. Sarr
College of Medicine
Department of Surgery
Mayo Clinic
Rochester
Minnesota
USA

ISBN 978-3-540-74505-1 ISBN 978-3-540-74506-8 (eBook)
DOI 10.1007/978-3-540-74506-8
Springer Heidelberg New York Dordrecht London

Library of Congress Control Number: 2012953477

© Springer-Verlag Berlin Heidelberg 2013

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media (www.springer.com)

Contents

Part I Surgery of the Pancreatic Carcinoma

1 Basic Chapter	3
René Mantke and Hans Lippert	
2 Commentary	23
Claudio Bassi, Giovanni Butturini, and Roberto Salvia	
3 Commentary	27
Thilo Hackert, Jens Werner, and Markus W. Büchler	
4 Commentary	35
Philippe Bachellier, E. Rosso, and D. Jaeck	
5 Commentary	43
Michael G.T. Raraty and John P. Neoptolemos	
6 Commentary	47
Tadahiro Takada, Keita Wada, and Keiji Sano	
7 Commentary	53
Andrew L. Warshaw	

Part II Chronic Pancreatitis Surgery

8 Basic Chapter	65
Hans-Ulrich Schulz, René Mantke, and Hans Lippert	
9 Commentary	79
David B. Adams and Katherine A. Morgan	
10 Commentary	89
P.C. Bornman and J.E.J. Krige	
11 Commentary	97
Thilo Hackert and Markus W. Büchler	
12 Commentary	105
Michael B. Farnell and Ashleigh L. Levendale	
13 Commentary	117
A. Fingerhut, S.V. Shrikhande, and P.J. Shukla	

14	Commentary	121
	Michael G.T. Raraty and John P. Neoptolemos	

Part III Cystic Neoplasms of the Pancreas: SCN, MCN, IPMN

15	Basic Chapter	127
	Thilo Hackert, Stefan Fritz, and Markus W. Büchler	
16	Commentary	135
	Katherine A. Morgan and David B. Adams	
17	Commentary	141
	Roberto Salvia, Marco Dal Molin, and Claudio Bassi	
18	Commentary	147
	C. Max Schmidt	
19	Commentary	151
	Masao Tanaka	

Part IV Surgery of Acute Pancreatitis

20	Basic Chapter	157
	René Mantke, Hans-Ulrich Schulz, and Hans Lippert	
21	Commentary	165
	Hans G. Beger, B.M. Rau, and B. Poch	
22	Commentary	173
	Peter Fagenholz and Carlos Fernández-del Castillo	
23	Commentary	179
	H.G. Gooszen	
24	Commentary	187
	C.W. Imrie and C.R. Carter	
25	Commentary	191
	Åke Andrén-Sandberg	
26	Commentary	197
	Michael G. Sarr	

Index	201
------------------------	-----

Introduction

This book, entitled *International Practices in Pancreatic Surgery*, represents the somewhat unique method of allowing high-visibility pancreatic surgeons from around the globe to discuss and compare their approaches to pancreatic cancer, cystic neoplasms of the pancreas, chronic pancreatitis, and acute pancreatitis. This book may serve as a code of practice for residents as well as specialized pancreatic surgeons in the clinical practice of pancreatic surgery. Diagnosis and subsequent management, from evaluation/staging to operative treatment, are described clearly, and the many forms of operative techniques are described, concentrating on the important details. The prevention and the treatment of common complications are included. Superb photos and drawings complement the comprehensive descriptions of the various surgical treatments for pancreatic cancer, cystic neoplasms of the pancreas, and chronic and acute pancreatitis. We specifically avoided needless theoretical information and discussions of the literature. This book was designed to be a practical guide, including information used on a day-to-day basis, for the surgeon, i.e., planning for resection, the instruments or sutures to be used, details of the possible reconstruction, the drugs required or that should be available, and perioperative management. Many well-known experts from all over the world describe their individual techniques and thoughts about pancreatic surgery. This discussion is combined with the latest results in their departments and contrasts with the four basic chapters from the Otto-von-Guericke-University in Magdeburg and the Ruprecht-Karls-University in Heidelberg.

The concept of this book is unique and immediately relevant to the daily routine of a surgeon. We hope you find it interesting, and we believe that it will provide new insights into the diagnosis and management of the more common disorders that require pancreatic surgery.

Brandenburg, Germany
Magdeburg, Germany
Heidelberg, Germany
Rochester, MN, USA

R. Mantke
H. Lippert
M.W. Büchler
M.G. Sarr

Part I

Surgery of the Pancreatic Carcinoma

1.1 Carcinoma of the Pancreatic Head/Periampullary Adenocarcinoma

1.1.1 Relevant Basic Information, Indication and Contraindication

Periampullary adenocarcinoma includes cancer of the pancreatic head, ampullary cancer, distal bile duct cancer, and duodenal cancer. About 70 % are carcinomas of the pancreatic head. The operation procedure is essentially the same for all of these types of neoplasms. The frequency of lymph node metastases in patients with pancreatic cancer is associated with a 5-year survival rate of only 5 % or less. This very aggressive tumor biology is the rationale for an extended lymph node resection in pancreatic cancer. Another major problem with performing operations for pancreatic head carcinoma is also the frequent presence of perineural invasion; perineural involvement is also associated with a very poor prognosis. Because involvement of the mesenteric neural plexus is extensive,

it is difficult to achieve a negative, retroperitoneal margin even with a radical resection.

Clinical symptoms (usually jaundice related to obstruction of the distal bile duct), computed tomography (CT), and endoscopic retrograde cholangiopancreatography (ERCP) or magnetic resonance imaging (MRI)/magnetic resonance cholangiopancreatography (MRCP) lead to the diagnosis of a periampullary carcinoma. We prefer a high-quality, multiphase, contrast-enhanced, thin-section, helical CT including angio-CT (Table 1.1). MRI requires more time to perform, is more expensive, less available, and more difficult to read, yet MRI is less “invasive” compared to CT and ERCP; moreover, CT is usually easier for a surgeon to interpret than MR images. Preservation of fat around the major peripancreatic vascular structures suggests a lack of direct tumor invasion and is consistent with the clinical prediction of “resectability”. Isolated involvement of the superior mesenteric vein or the portal vein is not necessarily a contraindication for resection. We believe that circumferential vessel involvement by tumor, infiltration of the hepatic or mesenteric artery, or occlusion by the tumor of these vessels should be absolute contraindications for resection. We use ERCP and biliary stents only selectively. If the diagnosis is clear and the operation can be done in a short time window, we see no need for insertion of an endoscopically placed endo-biliary stent when the findings on CT or MRI/MRCP are unclear, we will often proceed to ERCP including cytologic investigation. In the case of cholangitis or bilirubin levels $>300 \mu\text{mol/l}$ ($>18 \text{ mg/dl}$),

R. Mantke (✉)
Professor of Surgery, Chair, Department of Surgery,
City Hospital Brandenburg, 14770 Brandenburg,
Germany
e-mail: mantke@klinikum-brandenburg.de

H. Lippert, M.D.
Professor of Surgery, Chair, Department of Surgery,
Otto von Guericke University of Magdeburg,
39129 Magdeburg, Germany
e-mail: hans.lippert@med.ovgu.de

Table 1.1 Diagnostic in patients with suspicion of a periampullary carcinoma

Diagnostic method	Questions
Clinic	Jaundice, pain, vomiting, weight loss, glucose intolerance, new diabetes, gastrointestinal bleeding, steatorrhea, palpable abdominal mass
Laboratory evaluation	Standard parameters
Standard chest X-ray	Pulmonary lesions, metastases
Ultrasound	Pancreatic tumor, ascites, liver metastases
CT (high quality multiphase contrast enhanced, thin-section helical CT), Angio-CT	Size and location of the tumor, relationship to the mesenteric and portal vein and hepatic and superior mesenteric artery, liver or lymph node metastases, peritoneal metastases
ERCP	Strictures and obstruction of pancreatic or bile duct, endobiliary stenting, biopsy – cytologic investigations
Endoscopic ultrasonography	Size and location of the tumor, relationship to the mesenteric and portal vein and hepatic and superior mesenteric artery, local lymph node metastases, strictures and obstruction of pancreatic or bile duct
<i>MRT + MRCP + Angio-MRI</i>	<i>All-in-one procedure, size and location of the tumor, relationship to the mesenteric and portal vein and hepatic and superior mesenteric artery, liver or lymph node metastases, peritoneal metastases, strictures and obstruction of pancreatic or bile duct</i>
<i>Diagnostic laparoscopy (in combination with the resection procedure – in one step)</i>	<i>Small liver metastases (which are not seen with CT or MRI), peritoneal metastases</i>
<i>Preoperative biopsy</i>	<i>Histological confirmation of the tumor, only important for nonresectable patients</i>

Italic = optional tests for specification of the diagnosis

endoscopic papillotomy and stenting is indicated. In practice, we often get patients with suspected pancreatic cancer referred from other hospitals or gastroenterology departments with biliary stenting already performed for the treatment of jaundice. In contrast to some authors, we have not seen a greater rate of postoperative complications (fistulas, infections, mortality) in patients with endobiliary stents placed preoperatively.

Endoscopic ultrasonography is a relatively non-invasive diagnostic tool to investigate the primary neoplasm. If the CT or MRI is unclear, the patient will get an endoscopic ultrasonography to confirm or supplement the other diagnostic results. We see no need for a preoperative biopsy in potentially resectable patients because of high rates of false-negative biopsies and the risk of tumor seeding. Pancreas biopsy is usually only necessary in patients who will not be resected and palliative therapy will be done.

Staging: Operative resection only benefits patients with loco-regional disease. Preoperative imaging/staging should be undertaken to exclude distant metastases. There should be no evidence of involvement of the hepatic, celiac, or superior mesenteric arteries and no diagnostic evidence of

occlusion of the superior mesenteric or portal vein. Extensive resection should be avoided in patients with occlusion of the superior mesenteric vein (SMV) or portal vein (PV) and collateralization because of a strong risk of bleeding and a high mortality. Segmental vein resection and reconstruction of the portal and/or the superior mesenteric veins in selected patients provided a complete resection can be achieved with this procedure. Candidates for resection of the pancreatic head should have a good functional status and physiologic reserve to withstand the resection procedure. Chronologic age alone should not be a contraindication for pancreatic resection.

General contraindications for pancreas head resection are (exceptions are possible):

- Liver metastases
- Peritoneal metastases (malignant ascites)
- Other distant metastases
- Tumor involvement of the superior mesenteric artery (SMA) or hepatic artery
- Circumferential tumor involvement or occlusion of the SMV or PV
- Patient in an unsatisfactory medical condition, other relevant diseases limiting expected survival

Table 1.2 Steps of a standard procedure (Traverso–Longmire)

<i>Resection</i>	
1	Exploration
2	Biopsy of liver or peritoneal metastases if necessary
3	Elevation of the duodenum and pancreatic head (Kocher maneuver)
4	Division of the right half of the gastrocolic ligament
5	Mobilization of the right colon flexure
6	Division of the gastrohepatic ligament
7	Division of the gastroduodenal artery and identification of the portal vein on the superior border of the pancreas (attention of a relevant stenosis of the common hepatic artery or atypical arterial perfusion of the liver)
8	Exposure of the SMV at the inferior border of the pancreas
9	Division of the postpyloric duodenum
10	Freeing of the gallbladder and transection of the common hepatic bile duct
11	Division of the jejunum distal the ligament of Treitz and delivery the jejunum and the distal duodenum to the right of the superior mesenteric vessels
12	Division of the pancreas
13	Freeing the uncinate process and division of the lateral branches of the SMV and SMA
14	Complete lymphadenectomy
<i>Reconstruction</i>	
15	Pancreaticojejunostomy
16	Hepaticojejunostomy (optional t-tube)
17	Duodenojejunostomy
18	Drainage and closure of the abdominal wound

1.1.2 Surgical Technique

Our preferred procedure for pancreatic head resection is the pylorus-preserving pancreatoduodenectomy described by Longmire and Traverso (Table 1.2). Long-term survival has not been influenced by pyloric preservation in several studies. The Longmire/Traverso procedure is faster than the classic Kausch-Whipple procedure and perhaps more physiologic because of the preservation of the pylorus.

The pylorus-preserving pancreatoduodenectomy is started with a bilateral subcostal incision with an extension more to the right side of the upper abdomen (Fig. 1.1). We regularly use a self-retaining retraction system for the costal margin (Fig. 1.1). The liver and the peritoneal cavity are first inspected and palpated to exclude the presence of metastases. A wide Kocher maneuver is performed to confirm that the tumor does not invade the vena cava, the retroperitoneum, or the superior mesenteric artery (Fig. 1.2). Using bimanual palpation anterior and posterior to the SMA, it is possible to exclude a gross

tumor involving the SMA. Direct tumor invasion of the ligament of Treitz is a strong indicator for involvement of the SMA and a contraindication for resection.

Next the right half of the gastrocolic ligament is divided between ligatures or with the harmonic scalpel. The greater omentum is preserved on the transverse colon. Usually, the vascular supply to the greater omentum remains excellent after this procedure. At the end of the operation, the greater omentum is placed in the subhepatic space in front of the pancreatic anastomosis to cover this area to “protect” the pancreaticojejunostomy. Overall, well-vascularized omentum helps to control postoperative complications of the pancreatic anastomosis; omentum with a poor blood supply should be resected. The right colon flexure is then mobilized from the liver, the duodenum, and the anterior surface of the pancreatic head. The transverse mesocolon is detached from the pancreatic head down to the right lateral aspect of the superior mesenteric vein (Fig. 1.3). For optimal exposure of the infrapancreatic superior mesenteric vein and the anterior surface of the pancreatic





Fig. 1.1 Position of the patient (a), initiated incision for pylorus preserving pancreatoduodenectomy (b), and using a self-retaining retraction system after opening the abdomen (c)

head, it is necessary to divide the gastroepiploic artery and vein (gastrocolic trunk) (Fig. 1.3). The gastrohepatic ligament is then divided and the common hepatic artery, the gastroduodenal artery, and the suprapancreatic portal vein are identified (Fig. 1.4). Dividing the gastroduodenal artery is often necessary for a complete exposure and dissection of the suprapancreatic portal vein. The SMV is mobilized infrapancreatically by

following the venous branches of the transverse mesocolon that drain into the SMV.

At this point, it is very important to confirm that the confluence of the superior mesenteric vein and the splenic vein with the portal vein is not invaded by tumor. Using a blunt clamp, the pancreatic tissue can be mobilized carefully from the anterior surface of the confluence of the veins (Fig. 1.5). If this mobilization is possible, this



- 1 gastroduodenal and hepatic ln
- 2 superior pancreaticoduodenal anterior and posterior ln
- 3 inferior pancreaticoduodenal anterior and posterior ln
- 4 mesenteric ln (right side of the SMA)
- 5 suprapancreatic ln (hepatic artery and right side of celiac trunk)
- 6 aortoinferocaval ln

Fig. 1.2 (a) Mobilization by Kocher (not yet completed). A retropancreatic lymph node is marked by forceps (*c* colon, *vc* vena cava inferior, *gb* gallbladder, *s* stomach, *d* duodenum). (b) Relevant lymph node stations for carcinomas of the pancreatic head. Gray nodes are located behind the pancreas (1 gastroduodenal and hepatic lymph nodes, 2 superior pancreaticoduodenal anterior and posterior lymph nodes, 3 inferior pancreaticoduodenal anterior and posterior lymph nodes, 4 mesenteric lymph nodes, 5 suprapancreatic lymph nodes, 6 aortoinferocaval lymph nodes) (From O'Morchoe 1997). (Illustration by Reinhold Henkel, Heidelberg)




Fig. 1.3 The mesocolon transversum is detached from the pancreatic head to the right lateral aspect of the superior mesenteric vein and dividing the gastroduodenal venous trunk (vena gastroepiploica dextra). The right gastroepiploic artery and vein are divided at the anterior surface of the pancreatic head (*g* right gastroepiploic artery and vein, *gv* divided gastroduodenal venous trunk (vena gastroepiploica dextra), *SMV* superior mesenteric vein, *d* duodenum). (Illustration by Reinhold Henkel, Heidelberg)





Fig. 1.4 (a, b) Identification of the common hepatic artery, the gastroduodenal artery and the suprapancreatic portal vein (*gb* gallbladder, *s* stomach, *d* duodenum, *p* pancreas over the venous confluence, *pv* portal vein suprapancreatic, *ch* common hepatic artery, *g* divided gastroduodenal artery, *b* bile duct)




Fig. 1.5 Using a blunt clamp the pancreatic tissue can be carefully mobilized from the confluence of the superior mesenteric vein and the portal vein (SMV superior mesenteric vein, *p* pancreas, *d* duodenum, *u* uncinate process, *g* gallbladder, *m* mesocolon transversum, *c* vena cava inferior)

maneuver confirms resectability of the pancreatic head. In the absence of gross tumor infiltration of the duodenum, and especially in the postpyloric portion of the duodenum, a pylorus-preserving pancreatoduodenectomy is usually possible. The duodenum is then divided about 2–3 cm distal to the pylorus using a linear stapler (Proximate 75 mm Linear Cutter®, Ethicon Endo-Surgery, Johnson & Johnson, Somerville, USA, blue magazine). The stomach is then mobilized into the left upper abdomen. The gallbladder is removed and the common hepatic duct is transected superior to the junction with the cystic duct. To avoid continuous leakage of bile into the abdomen, the hepatic duct is occluded using a nontraumatic vascular clamp. The jejunum is divided about 8–12 cm distal to the ligament of Treitz using a linear stapler (Proximate 75 mm Linear Cutter® Ethicon Endo-Surgery, blue magazine) and the mesentery divided using the harmonic scalpel (Generator 300® Ethicon Endo-Surgery, Johnson & Johnson); the harmonic scalpel offers excellent control of bleeding and saves time. After mobilizing the ligament of Treitz and the fourth portion of the duodenum, there is free communication between the left and the right side of the abdomen posterior to the superior mesenteric vessels. The fourth portion of the duodenum and the short segment of devascularized proximal jejunum are then drawn to the right side of the abdomen posterior




Fig. 1.6 (a, b) A special nonmetallic probe is placed in front of the venous confluence between the confluence and the pancreas and the pancreas is divided with a scalpel (using an electrocautery scalpel, cutting mode, low energy, a thermal necrotic zone on the specimen can complicate the diagnosis of a tumor free pancreatic margin by the pathologist) (SMV superior mesenteric vein, *b* bile duct, PV portal vein, *ha* hepatic artery)

to the superior mesenteric vessels through the bed of the duodenum. Before dividing the neck of the pancreas, we place single sutures on the superior and inferior rim of the pancreas (Prolene® 4/0, Ethicon) to control small vessels that often bleed during transection of the pancreas. A nonmetallic, special probe or a blunt clamp is placed anterior to the venous confluence but posterior to the neck of the pancreas, and the pancreas is divided with a regular scalpel (Fig. 1.6). When using an electrocautery instrument (cutting mode, low energy) to transect the pancreatic neck, the pathologist can encounter problems determining a tumor-free resection margin if the tumor has reached the thermal necrotic zone. Usually several bleeding points are evident after transection of the pan-




Fig. 1.7 The venous confluence will be exposed by a tension of the pancreas head with the left hand of the surgeon to the right. Small veins from the uncinate process to the superior mesenteric vein or in the portal vein were ligated selectively (*pv* portal vein, *p* pancreas corpus, *ph* pancreas head in the left hand)

creas. We use bipolar electrocautery for hemostasis of these bleeding points on both sides of the transected pancreatic tissue.

The venous confluence is now exposed. Rightward traction applied to the pancreatic head using the left hand is very helpful in this situation. Several small veins drain the uncinate process directly into the superior mesenteric vein or into the portal vein. Such branches are isolated and ligated selectively (Fig. 1.7). After this maneuver, the portal vein can be retracted medially. Sometimes, vascular involvement of the SMV or the portal vein is able to be seen only at the time of operation (a point of no return because the pancreas has been fully transected). Tumors in the uncinate process can be especially adherent to these vessels. Segmental vein resection and reconstruction of the portal and/or the superior mesenteric vein is possible if complete resection (RO) can be achieved with this procedure. Of course, those patients with vascular involvement also have a high rate of lymph node metastases and retropancreatic perineural infiltration. These facts limit the long-term survival independently of the RO vein resection. The long-term survival after RO resection including vein resection is, however, better than palliative surgery in several studies. Vein resection should be an individual decision in every patient.

Several types of vein resection are possible. The lateral wedge resection is the simplest procedure (Fig. 1.8). The vein is clamped using a “side-biting” vascular clamp laterally such that venous flow persists. The defect is closed over the vascular clamp with a continuous nonabsorbable 5/0 monofilament suture (Prolene®, Ethicon). The functional diameter of the SMV or portal vein should not be decreased significantly with this type of resection provided the lateral defect does not involve much of the circumference of the vein. For greater tumor involvement of the vein, a circumferential resection or venous patch reconstruction of the vein is a better oncologic procedure (Fig. 1.8). Venous reconstruction is possible with a primary anastomosis, an autologous vein graft (superficial femoral vein), or synthetic graft. We prefer to use an autologous vein graft, or a direct suture of the vein which is often possible after a wide mobilization. The junction of the splenic vein and the SMV can be preserved by tangential excision of the SMV (Fig. 1.8). After dealing with the SMV and portal vein, attention is turned to resecting the uncinate process from the superior mesenteric artery. The superior mesenteric artery can and should be identified easily by palpation and visualization. The specimen is now only fixed by the retroperitoneal tissue around the SMA. Usually, many small lymphatic vessels are located in this tissue. This retroperitoneal margin of the specimen often shows invasion of tumor cells into the lymphatic vessels and perineural tissues which is the cause for the relatively high rate of local recurrence after resection of carcinomas in the pancreatic head. This tissue is divided on the right side of the SMA using individual ligatures or the harmonic scalpel. The inferior pancreaticoduodenal artery should be sought, isolated, and ligated selectively (Fig. 1.9). We send the complete specimen to the pathology department for frozen section analysis of the bile duct, pancreatic transection margin and, if necessary, the postpyloric duodenum. Positive resection margins need a further resection of the bile duct, pancreas, or stomach. Further resection in the area of the SMA (retroperitoneal margin) is usually not indicated or not possible, which is why we do not obtain a frozen section in this area.




Fig. 1.8 (a–d) Portal vein reconstruction. The lateral wedge resection is the simplest procedure. The vein is clamped out laterally and the venous flow is still existent. The defect is closed over the vascular clamp with a continuous non absorbable 5/0 monofilament suture (Prolene®, Ethicon, Johnson and Johnson, Somerville, USA). The open diameter of the SMV or portal vein should not be reduced significantly (a, b). The drawing shows a tangential

technique for interposition of a superficial femoral vein (SFV) segment with the intent to save the junction with the splenic vein (c). The reconstruction with a superficial femoral vein patch is also possible (d). (cSMV laterally clamped SMV, p pancreas, pv portal vein, sSMV sutured SMV, ha hepatic artery, pd pancreatic duct). (Illustration by Reinhold Henkel, Heidelberg)




Fig. 1.9 Several small veins from the uncinate process or the pancreatic head which drain directly into the superior mesenteric vein or in the portal vein are ligated. The specimen is only fixed by the retroperitoneal tissue around the SMA. This tissue is divided on the right side of the SMA using sutures. The inferior pancreaticoduodenal artery should be isolated and ligated selectively (SMV superior mesenteric vein, *p* pancreas, *d* duodenum, SMA superior mesenteric artery, *pda* inferior pancreaticoduodenal artery). (Illustration by Reinhold Henkel, Heidelberg)

Lymph node metastases are common in periampullary carcinoma and are critically relevant for prognosis. Extended lymphadenectomy is the standard procedure in our opinion. The extent of lymphadenectomy remains a matter of debate. During the pathologic investigation of the margins of the specimen, we complete the lymphadenectomy. The anterior and posterior pancreaticoduodenal lymph nodes are usually located in the specimen. For this reason, it is important to start the Kocher maneuver at the level of the right anterior wall of the inferior vena cava (Fig. 1.2). Usually, the right renal vein is identified at this time. Lymph nodes are removed from the hepatoduodenal ligament (caudal to the former cystic duct junction). Typically, lymph nodes from the cystic duct or from the corner between bile duct and duodenum are removed en bloc with the specimen. Other lymph nodes around the portal vein or hepatic artery are dissected separately (Fig. 1.2). We dissect the hepatic artery from its lymphatic




Fig. 1.10 Operation field after resection of the specimen at the retroperitoneal level of the anterior surface of the vena cava (a) and after dissection of the lymph nodes in the aortocaval sulcus (b) (*h* hepatic artery, *p* portal vein, *smv* superior mesenteric vein, *vc* vena cava, *a* aorta, *r* right renal vein, *g* gonadal vein)

tissue up to the level of the celiac trunk (level of the left gastric artery). We prefer to use bipolar cautery for this procedure. The lymph nodes on the right side of the SMA have been resected with the specimen. We avoid routine dissection of lymph nodes on the anterior and left side of the SMA because of the high morbidity (diarrhea, malnutrition). Other lymph nodes that appear to be malignant, besides the fourth portion of the duodenum and the ligament of Treitz, are also removed with the specimen. Lymph nodes in the aortocaval groove are removed separately (Fig. 1.10).

Reconstruction: The transected jejunum is pulled through a vascular window in the transverse mesocolon and into the subhepatic space. We prefer this pathway for the jejunal limb rather




Fig. 1.11 Performing the outer posterior row using a low absorbable monofilament single sutures (PDS® 4/0 Ethicon, Johnson and Johnson, Somerville, USA), end-to-side anastomosis, pancreatic duct intubated with a flexible Simon-Weidner probe (*j* jejunal loop, *p* pancreas, *pd* pancreatic duct, *h* hepatic artery, *d* dorsal layer with single sutures, *m* transverse mesocolon)

than posterior to the mesenteric vessels (in the bed of the duodenum) because of the risk of tumor recurrence in the area of the SMA and possible obstruction of the jejunal limb. The closed end of the jejunum is oversewn using a 4/0 absorbable suture (Vicryl®, Ethicon). The jejunal limb placed subhepatic has no tension when positioned near the cut edge of the pancreatic remnant. The posterior surface of the pancreatic remnant is mobilized carefully from the venous confluence and the splenic vein using bipolar forceps cautery.

We perform a modified, Cattell-Warren, duct-to-mucosa anastomosis in an end-to-side fashion without stenting. We start with the outer, posterior row using absorbable, 4/0 monofilament interrupted sutures (PDS®, Ethicon) (Figs. 1.11 and 1.12). Depending on the size of the pancreatic duct, we use one or two sutures on each side (posterior, anterior, cranial, and caudal, minimum four sutures, maximum eight sutures). In some cases with a very small pancreatic duct, an anastomosis of the pancreatic duct to the jejunal mucosa with only two sutures still works. The outer anterior row between the pancreas and the jejunal wall is also done using an interrupted suturing technique using PDS® 4/0 (Ethicon) placed 3–5 mm between stitches (Fig. 1.13). The sutures between the pancreatic tissue and the jejunal limb are sometimes difficult to place




Fig. 1.12 (a, b) The pancreatic duct is fixed to a small incision in the corresponding jejunal limb using low-absorbable monofilament single sutures (PDS® 5/0, Ethicon, Johnson and Johnson, Somerville, USA). Depending on the size of the pancreatic duct we use 1 or 2 sutures on each side (posterior, anterior, cranial and caudal) (*j* jejunal loop, *p* pancreas, *h*- hepatic artery, *d* not sutured 5/0 PDS sutures – duct-to-mucosa anastomosis with five single interrupted sutures, *c* completed duct-to-mucosa anastomosis)




Fig. 1.13 Completed outer anterior row of the pancreatico-jejunojejunostomy, using PDS® 4/0 (Ethicon, Johnson and Johnson, Somerville, USA) in an interrupted single suture technique (*j* jejunal loop, *p* pancreas, *h* hepatic artery, *c* completed anterior row)

especially in nonchronic pancreatitis patients because of the softness of the pancreatic tissue. The amount of pancreatic tissue we include in the suture depends on the texture of the pancreas. In a very soft pancreas, the amount of needed pancreatic tissue is greater than in patients with a more fibrotic pancreas like in chronic pancreatitis. Sometimes, it is very helpful to use a U stitch technique to incorporate more tissue in the stitch. The technique of tying the knots itself is important, too. It is crucial to avoid any sawing movements with the suture. The suture should be tied very gently with mild compression of the two tissues with a distance of 3–5 mm between sutures.

Next, the end-to-side hepaticojejunostomy is performed 8–10 cm distal to the pancreatic anastomosis. This point is chosen to avoid kinking of the jejunal limb. The antimesenteric jejunum is opened using electrocautery. The jejunal enterotomy should be a little smaller than the lumen of the hepatic duct, because it will be dilated during manipulation. The posterior part of the anastomosis is performed with a single layer of continuous absorbable monofilament suture (PDS® 5/0, Ethicon) (Fig. 1.14), while the anterior part of the anastomosis is performed with a single layer of the same suture material. To allow better visualization, the sutures are not tied until all have been placed. When the lumen of the hepatic duct is very small, an anastomosis should be performed as described by Goetze-Guetgemann (Fig. 1.15). We do not routinely use t-tubes or stenting jejunal tubes. The jejunum is then fixed with two, single, absorbable sutures (Vicryl® 4/0, Ethicon) to the transverse mesocolon. The defect at the ligament of Treitz is obliterated to avoid hernias. Next, the attention is directed at restoring gastrointestinal continuity. We first inspect the postpyloric duodenum to assure sufficient vascular perfusion of the proximal duodenum. Sometimes the vascular inflow or outflow is compromised and the postpyloric area takes on a purplish, ischemic hue; because we know that is ischemia of the duodenal cuff results in impaired gastric emptying in the postoperative course, we prefer a distal resection of the stomach and reconstruction as described by Kausch-Whipple. If the blood perfusion to the postpyloric duodenum is sufficient, an antecolic, end-to-side duo-




Fig. 1.14 A end-to-side hepaticojejunostomy is performed using a single-layer continuous absorbable monofilament suture (PDS® 5/0, Ethicon, Johnson and Johnson, Somerville, USA) in the back and a single layer interrupted absorbable monofilament suture (PDS® 5/0) in the front. (a) Continuous posterior row completed (*j* jejunal loop, *h* hepatic duct, *ha* hepatic artery, *l* liver, *a* anastomosis), (b) anterior row not yet sutured, (c) anastomosis completed

denojejunostomy is performed in a double layer continuous technique with an absorbable suture (Vicryl® 4/0, Ethicon) (Figure 1.16).

The biliary anastomosis and the posterior part of the pancreaticojejunostomy is drained by