


Adolescent Urology

and Long-Term
Outcomes

Christopher
Woodhouse



WILEY Blackwell

CONTENTS

[Cover](#)

[Dedication](#)

[Title page](#)

[Copyright page](#)

[Contributors](#)

[Preface](#)

[CHAPTER 1: Adolescent and transitional urology—an introduction](#)

[Definition](#)

[Background](#)

[The process of transition](#)

[Training requirements](#)

[The adolescent clinic](#)

[Who can be discharged without long term follow-up?](#)

[Acquisition of patients in adolescence](#)

[Outcome measures](#)

[Commentary](#)

[PART I: The kidney](#)

[CHAPTER 2: Anatomical and cystic anomalies of the kidney](#)

[Pelvi-ureteric junction \(PUJ\) obstruction](#)

[Horseshoe kidney](#)

[Unilateral renal agenesis](#)

[Cystic anomalies](#)

[Commentary](#)

[CHAPTER 3: Stones](#)

[Introduction](#)

[Primary stones in adolescents](#)

[Treatment outcomes](#)

[Complications](#)

[Commentary](#)

[CHAPTER 4: Prune-belly syndrome](#)

[Introduction](#)

[The urinary tract](#)

[Commentary](#)

[PART II: The ureter](#)

[CHAPTER 5: Renal function in adolescents](#)

[Mechanisms of renal damage](#)

[Long term consequences of reduction in nephron mass; subtotal nephrectomy and renal hyperfiltration](#)

[Practical management](#)

[Nephrologic clinic practice](#)

[End-stage renal disease and transplantation](#)

[CHAPTER 6: Structural anomalies and reconstruction of the ureter](#)

[Duplication of the ureter](#)

[Megaureters](#)

[Ureterocele](#)

[Ectopic ureters](#)

[Ureteric reconstruction](#)

[Commentary](#)

[CHAPTER 7: Vesico-ureteric reflux](#)

[Introduction](#)

Reflux found after fetal ultrasound, spontaneous resolution with normal kidneys and bladder

Reflux with abnormal kidneys

Reflux corrected by surgery of one sort or another

Reflux unresolved by adolescence with normal or abnormal kidneys

Reflux in adults

Reflux with other bladder/urinary reservoir abnormalities

Reflux after renal transplantation

Reflux and pregnancy

Inheritance and screening

Commentary

PART III: The bladder

CHAPTER 8: The neuropathic bladder and spina bifida

Introduction

Survival

Renal function

Hypertension

Renal stones

The bladder

Surgery in adulthood

Latex allergy

Preparation for adulthood

Mobility

Sexual development

Fertility and pregnancy in females

Fertility in males

Partnerships and other social outcomes

Commentary

CHAPTER 9: Lower urinary tract replacement and reconstruction

Introduction

Reservoirs

Ileum

Colon

Rectum

Autoaugmentation

Artificial bladders

The conduit

General consequences of lower urinary tract reconstruction

Continence

Renal function

Stones

Histological changes

Perforation

Infection

Neoplasia

Urine testing for pregnancy

Pregnancy and delivery

Commentary

CHAPTER 10: Urinary diversion and undiversion

Introduction

Ileal conduit

Colon conduits

Cutaneous ureterostomy

[Urinary undiversion](#)

[Commentary](#)

[CHAPTER 11: Other bladder conditions](#)

[Non-neuropathic neuropathic bladder](#)

[Enuresis](#)

[Ketamine](#)

[Commentary](#)

[CHAPTER 12: Exstrophy and epispadias](#)

[Introduction](#)

[Mortality](#)

[The bladder](#)

[Bladder cancer](#)

[The kidneys](#)

[The Penis](#)

[Male sexual function](#)

[Females born with exstrophy](#)

[Monsplasty \(both sexes\)](#)

[Sexual health and contraception](#)

[Inheritance of exstrophy](#)

[Psychological issues](#)

[Commentary](#)

[CHAPTER 13: Cloacal exstrophy](#)

[Introduction](#)

[Mortality and long-term outcome](#)

[The kidneys](#)

[The bladder](#)

[Gender assignment](#)

[Penile reconstruction and male sexuality](#)

[Vaginal construction and sexuality in 46XY females](#)

[Genital reconstruction and sexuality in females](#)

[Quality of life](#)

[Commentary](#)

[PART IV: The urethra](#)

[CHAPTER 14: Posterior urethral valves](#)

[Introduction](#)

[The bladder](#)

[Renal function](#)

[Sexuality, fertility and quality of life](#)

[Commentary](#)

[CHAPTER 15: Hypospadias](#)

[Introduction](#)

[Measures of success](#)

[Surgical outcomes](#)

[Micturition](#)

[Sexuality](#)

[Hypospadias in adults](#)

[Fertility](#)

[Inheritance](#)

[Psychological outcomes](#)

[Commentary](#)

[PART V: Disorders of sex development](#)

[CHAPTER 16: Scrotal abnormalities](#)

[Undescended testes](#)

[Varicocele](#)

[CHAPTER 17: The overvirilized female](#)

[Introduction](#)

[Congenital adrenal hyperplasia](#)

[Feminizing genitoplasty](#)

[Cliteroplasty](#)

[Psychological function](#)

[Sexuality](#)

[Fertility](#)

[Screening](#)

[Vaginal reconstruction](#)

[Other overvirilizing \(or under oestrogenizing\) conditions](#)

[Cloaca](#)

[Commentary](#)

[CHAPTER 18: The undervirilized male](#)

[Introduction](#)

[The penis](#)

[Surgical enlargement of the penis](#)

[Absent or very inadequate penis](#)

[Androgen insensitivity syndrome](#)

[5-alpha reductase deficiency \(5 \$\alpha\$ RD\)](#)

[Testicular neoplasia](#)

[Fertility](#)

[Congenital adrenal hyperplasia in males](#)

[Other aspects of undervirilization](#)

[Commentary](#)

[CHAPTER 19: Adolescent sexuality and counseling](#)

[Introduction](#)

[Age of puberty](#)

[Dating](#)

[Sexually transmitted disease](#)

[Teenage pregnancy and paternity](#)

[Pregnancy testing](#)

[Inheritance](#)

[Drug abuse](#)

[Further learning](#)

[PART VI: Neoplasia](#)

[CHAPTER 20: Neoplastic disease: outcomes in survivors](#)

[Introduction](#)

[Wilms' tumor](#)

[Rhabdomyosarcoma](#)

[New cancer in adolescents](#)

[The consequences of cure](#)

[Commentary](#)

[Index](#)

[End User License Agreement](#)

List of Tables

Chapter 01

[Table 1.1 An estimate of the number of adolescent urologists required in the United Kingdom. Births per year are based on published incidences.](#)

[Table 1.2 Details required for successful transfer of children to the adolescent urology clinic. Not all are required for all patients.](#)

[Table 1.3 Associated specialists required for support in an adolescent clinic by urological diagnosis. This list is not exclusive. There is a universal need for](#)

specialist radiologists and nuclear medicine specialists. Psychologists are often required for any of these diagnoses.

Table 1.4 (a) The results of a questionnaire undertaken in the adolescent clinic at University College London Hospital over an 8 week period in 2010. All patients were asked to name their current occupation, which was recorded against their principal diagnosis. No patient refused to answer. (b) Examples.^a

Chapter 04

Table 4.1 Occupations of 27 patients with PBS who have survived into adulthood (N = 41), excluding students and those for whom there are inadequate data (n = 14).

Chapter 05

Table 5.1 Diagnoses seen in the joint nephro-urological adolescent clinic at University College London and forming the basis for our data on CAKUT. Vesico-urteric reflux (VUR) is used for those presenting in childhood who have reflux confirmed by MCU; otherwise the term “renal dysplasia” is used.

Chapter 07

Table 7.1 Pregnancy outcomes in women with significantly impaired renal function following vesico-ureteric reflux.

Chapter 08

Table 8.1 Long-term outcome of 156 children born with spina bifida 1978-1990.

Table 8.2 Family and household tasks undertaken by 102 children with spina bifida (40% ambulant).

Chapter 09

Table 9.1 The components of the lower urinary tract and tissues available to replace it.

Table 9.2 Venous blood biochemical and hematological changes in patients with intestinal urinary reservoirs. Data are very unreliable and the incidences may be taken with a pinch of salt. The denominators are usually unknown, follow-up variable and definitions poorly described. The more the abnormalities are sought, the more often they are found. Arterial blood gas analyses give higher incidences for pH changes.

Table 9.3 Guidance on prevention of stones in intestinal urinary reservoirs. Based on theoretical data. There is no evidence that stone incidence is reduced with these regimes.

Chapter 11

Table 11.1 Summary of the effects of drug abuse in adolescents and young adults.

Chapter 12

Table 12.1 Outcomes of childhood surgery for exstrophy with special reference to continence. Data are best estimates from recent literature from specialist centers. Approximations have been taken as reporting systems are not consistent between papers.

Table 12.2 Cases of cancer in 61 patients with exstrophy with a minimum follow up of 35 years.^a

Table 12.3 Renal abnormalities recorded in 462 patients with exstrophy at birth.

[Table 12.4 Results of a literature search for the results of semen analysis in adults born with exstrophy.](#)

Chapter 13

[Table 13.1 Gender assignments of babies born with cloacal exstrophy in the literature up to 2003.](#)

[Table 13.2 Outcome of psychosexual assessment of male \(46XY\) children and adolescents born with cloacal exstrophy and raised as females.](#)

[Table 13.3 Gender identity outcome of 46XY adults born with cloacal exstrophy, according to gender assignment at birth.](#)

Chapter 14

[Table 14.1 Minimum annual follow-up for adolescents and adults born with posterior urethral valves.](#)

[Table 14.2 Incidence of end stage renal failure in patients born with posterior urethral valves.](#)

Chapter 15

[Table 15.1 The features of the Genital Perception Score described by Mureau *et al.*^a They have been divided into those are surgically correctable and those which are not. Overall, there is discordance between surgeon and patient. However, when recalculated in this table there is better concordance for the surgical result.](#)

[Table 15.2 Development of sexual mile stones in adolescents operated for hypospadias.](#)

[Table 15.3 Sexual activity in men operated for hypospadias.](#)

Chapter 16

Table 16.1 Paternity rates in co-habiting men operated for UDT in childhood.

Table 16.2 Relative risk of the development of a neoplasm in boys operated for UDT according to the performance of a biopsy.

Table 16.3 Prevalence of varicocele and associated testicular hypotrophy by age.

Chapter 17

Table 17.1 The number of women with CAH who have a patent vagina after primary surgery (1^y patency) and after revision surgery (2^y patency). In two of the papers it is possible to divide the patients into those with a low confluence (easy cases) and a high confluence (difficult cases). In the paper by Alizai *et al.* the mean age of the patients was 13.1; although all were post pubertal all were thought to be too young for intercourse to be considered.

Table 17.2 Ages at which women with CAH thought that feminizing genitoplasty should be done.

Chapter 18

Table 18.1 Incidence of neoplasia in the gonads of undervirilized males (46XY).

Table 18.2 Types of neoplasm found in the gonads of patients with DSD.

Chapter 19

Table 19.1 Summary of the effects of drug abuse on the genito-urinary system. Prices are approximate, researched in 2012.

Table 19.2 Incidence of components of sexual dysfunction in regular users of illegal drugs.

List of Illustrations

Chapter 01

Figure 1.1 Graph with logarithmic scale to show the number of papers listed in PubMed with the search terms “adolescent” (solid line) and “adolescent urology” (broken line), by year. The data are inaccurate in about the first 50 years as there were no key words in most papers. On checking a random sample from the early years, most had no relevance to urology.

Figure 1.2 Papers listed in PubMed under “transitional care” (dark line) and “transitional care, urology” (light line) from 1974, when the first paper appears, to 2012.

Chapter 02

Figure 2.1 (a) Intravenous urogram of a middle-aged female with uncomplicated recurrent cystitis and no upper tract symptoms. (b) CT scan of the same woman about 6 years later when she presented with a 2-year history of recurrent left flank pain associated with fluid intake.

Chapter 04

Figure 4.1 (a) Voiding cystogram of a patient with prune-belly syndrome before and (b) after full-length Otis urethrotomy, showing that the urethra fills with contrast and voiding occurs.

Figure 4.2 (a) Cystogram of the same patient before and (b) after micturition, showing that the bladder, although huge, empties virtually to completion.

Figure 4.3 (a-c) Clinical photographs of a boy with prune-belly syndrome showing the process of sitting.

up from the lying position. He has to roll to one side first to allow the lateral abdominal muscles to complete the process.

Figure 4.4 Clinical photograph of the same boy showing that hip flexion is normal.

Figure 4.5 Clinical photograph of a man with prune-belly syndrome showing the smooth but “pot bellied,” appearance of the abdominal wall that may be seen without abdominal-wall reconstruction. Note, also, the excess lumbar lordosis.

Figure 4.6 Clinical photograph of a man with prune-belly syndrome who has not had reconstruction, showing that the wrinkled appearance of the abdominal wall may remain in adult life.

Chapter 06

Figure 6.1 Intravenous urogram of an adult with a minor degree of mega ureter on the right. The kidney was normal. Although the IVU is out of favor as an imaging modality, it remains a good way of seeing the whole ureter on one film.

Figure 6.2 Water color of bilateral megaureters. On the left, the abnormality is minimal. On the right, the characteristic narrow area just above the bladder is seen (arrowed).

Figure 6.3 Ultra sound of the pelvis in an adult showing the characteristic appearance of a mega ureter with gross dilatation down to a point just above the bladder with a narrow area distally as it enters the bladder (arrowed—(a)). (b) The kidney of the same patient with arrows indicating the thinness of the parenchyma. This kidney had a normal GFR.

Figure 6.4 (a) IVU of an adult woman with a symptomless ureterocele (arrowed). It shows the characteristic “cobra-head” appearance. The kidney was normal. (b) Ureteroceles are also visualized on ultrasound as in this case.

Figure 6.5 Coronal (a) and axial (b) T2W MRI images showing an ectopic ureter entering the urethra just below the bladder neck in a woman with a didelphic uterus.

Figure 6.6 Early postoperative cystogram (ureteric stents are still *in situ*) showing filling of the right ileal ureter with contrast. Bladder and ileal ureter emptied completely after voiding, but the ileum would still act as a storage organ between voids.

Chapter 07

Figure 7.1 (a) Centile charts for systolic (left chart) and diastolic (right) blood pressure in mm. Hg by age in years for boys on the 50th centile for height. (b) Centile charts for systolic (left chart) and diastolic (right) blood pressure in mm. Hg by age in years for girls on the 50th centile for height.

Figure 7.2 Incidence of urinary infection complicating pregnancy. Data from Mansfield *et al.* Incidences for the index patients (those with reflux) were obtained by telephone interview by the authors. Other data are normative values quoted by the authors.^a*Note:*

^aMansfield, J. T., Snow, B. W., Cartwright, P. C. and Wadsworth, K. (1995) Complications of pregnancy in women after childhood reimplantation for vesicoureteral reflux: an update with 25 years of followup. *Journal of Urology* 154(2 Pt. 2), 787-790.

Chapter 08

Figure 8.1 Survival of an unselected series of patients born with spina bifida. Data from Hunt and Oakeshott 1990-2003 and Oakeshott 2010.^{a,b}

Figure 8.2 Histogram to show the causes of death in an unselected series of 695 adults born with spina bifida.

Figure 8.3 (a) A patient prepared for surgery. His ileal conduit stoma in the left iliac fossa is too low and invisible to him because of the horizon created by his upper abdomen. X marks the new site for the stoma. Note also that, although he is lying 'flat' on the table, he has a fixed flexion deformity of the hips which has to be supported during surgery and in bed postoperatively. (b) The patient sitting in his wheelchair postoperatively with a stoma that is visible to him.

Figure 8.4 Clinical photograph of an adult patient with spina bifida. She is lying on her right side, head to the right of the picture. It illustrates a kyphosis of 90° at the top of the sacrum marked in red.

Figure 8.5 A CT scan of a patient aged 35 years with spina bifida to show the distorted anatomy that can occur due to progressive kypo-scoliosis. The chest is supine on the x-ray table but the abdomen has been rotated by the spinal deformity and appears to be in the left-side-up position. The left kidney, with a calculus in it, is completely covered by the rib cage and was approached for PCLN above the left ribs.

Figure 8.6 Percentage of children who start ambulation (with or without aids) and percentage who continue to walk up to 9 years 1 month, by spinal level.

Figure 8.7 (a) Sexual activity in women with spina bifida; (b) sexual activity in men with spina bifida.

Figure 8.8 Graph showing the incidence of births of children with neural tube defects (per 100 000 live births) by year. The hatched line shows the sales of folic acid.

Chapter 09

Figure 9.1 Pathological specimen from a ureterosigmoidostomy showing a carcinoma arising from the anastomosis between the ureter on the left and the colon on the right.

Figure 9.2 MRI scan of a carcinoma in a right colon reservoir (arrowed). This patient had multiple dysplastic polyps elsewhere in the colon and the tumor in the reservoir was assumed to be a carcinoma of the colon rather than a result of urine in a reservoir.

Figure 9.3 Endoscopic view of an 8 mm anastomotic neoplasm that was malignant on histology.

Figure 9.4 (a) Clinical photograph of a Monti catheterizable channel formed from ileum. (b) Diagram of the same. Note that the pedicle is at one end allowing a good length for implantation into the urinary reservoir.

Figure 9.5 Cystogram (lateral view) showing an hour-glass deformity of a neuropathic bladder, which had been augmented with ileum several years earlier and developed late incontinence.

Figure 9.6 Clinical photograph showing a mixture of small stone fragments mixed with mucus and larger stones, removed from an intestinal reservoir.

Figure 9.7 (a) Plain x-ray to show a stone in a urinary reservoir. (b) The stone being removed from the reservoir.

Figure 9.8 Plain abdominal x-ray of a patient with a Mainz II showing air in the right ureter (arrowed).

Chapter 10

Figure 10.1 CT loopogram showing bilateral reflux into both ureters indicating that, although the ureters and renal pelvises are dilated, the system is not obstructed.

Figure 10.2 Clinical photograph showing a stenosed conduit stoma.

Figure 10.3 Loopogram to show a long conduit with a redundant loop in the pelvis that drains poorly.

Figure 10.4 Clinical photograph to show a prolapsed ileal conduit stoma.

Chapter 11

Figure 11.1 The prevalence of monosymptomatic enuresis in European children by age in years.

Figure 11.2 Cystogram in a ketamine abuser showing a small bladder and bilateral reflux.

Figure 11.3 CT scan to show papillary necrosis (arrowed) in a patient who abuses ketamine.

Figure 11.4 Cystoscopic findings of two daily ketamine abusers who had been inhaling ketamine nasally for 4 years (left) and 7 years (right), respectively.

Chapter 12

Figure 12.1 (a) Clinical photograph of a boy; (b) of an adolescent with reconstructed exstrophy from

Trendelenberg's paper of 1906.

Figure 12.2 (a) Clinical photograph of an adult man presenting with previously untreated exstrophy. The epispadiac penis had been closed. The whole pelvic floor has prolapsed. (b) The effect of pushing the bladder plate back into the pelvis, which reduces the prolapse.

Figure 12.3 Graph of deaths in patients born with exstrophy by year of series publication. The broken line is for children under 10 years old. The solid line is for premature deaths over the age of 10.

Figure 12.4 The percentage continence rate in children born with exstrophy after reconstruction of the bladder by year of publication. The mean calculated for the previous 60 years from 1970 is shown by * and is 5%.

Figure 12.5 (a) Plain x-ray of an exstrophy pelvis showing a stone in an augmented bladder and the penis seen "end-on." (b) Same patient after removal of the stone.

Figure 12.6 (a) Clinical photograph of a case of bladder cancer ulcerating through the scar of the bladder closure in exstrophy. (b) Computerized tomography of the same patient showing tendrils of the tumor mass extending out to the periprostatic tissues (ringed).

Figure 12.7 Diagram of the exstrophy pelvis showing the downward rotation with the inferior pubic ramus parallel to the floor when standing (right). Normal orientation is shown on the left.

Figure 12.8 Clinical photograph of a typical penis in the flaccid state in a man with exstrophy. The arrow

indicates the recession in the mons area, which may emphasize the penile abnormality.

Figure 12.9 Clinical photograph to show the dorsal chordee that is a natural feature of the exstrophy/epispadias complex. In exstrophy there is no cross circulation between the corpora so both sides have to be infused separately.

Figure 12.10 Clinical photograph to show the distribution of pubic hair with the older, "W," type of closure or with excessive midline scarring.

Figure 12.11 Cavernosogram showing the site of curvature (arrowed) causing dorsal chordee.

Figure 12.12 (a) Clinical photograph showing both dorsal chordee and sharp deviation to the left. This is caused by partial fibrosis of the left corpus seen in the cavernosogram in (b).

Figure 12.13 (a) Clinical photograph showing the rudimentary penis that occurs when both corpora are damaged. (b) Cavernosogram and cystogram of the same case showing small corpora with no proximal portion and no attachment to the pubic rami. Note that the penis lies higher up the supra pubic area as a result.

Figure 12.14 Photograph of the natural erection of the patient in Figure 12.9 after surgical correction of chordee.

Figure 12.15 (a) Diagram of the sagittal view female pelvis in exstrophy to illustrate the anterior displacement of the pelvic organs. Reprinted with permission from Woodhouse, C. R. J. (2001) *The genitalia in exstrophy and epispadias*, in *Pediatric Urology* (eds. J. P. Gearhart, R. C. Rink and P. D. E. Mouriquand). W. B. Saunders & Co., Philadelphia, PA.

pp. 557-564. (b) Clinical photograph of a female pelvis with exstrophy. Note the anterior position of the labia and the small introitus. She had an intestinal reservoir and the urethra had been removed.

Figure 12.16 Clinical photograph of the perineum to show the appearance of the vagina after surgical opening of the introitus.

Figure 12.17 (a) Clinical photograph of a woman with exstrophy after reconstruction of the genitalia, pubic hair and umbilicus. (b) Clinical photograph of the same patient showing that the vagina has not been moved, but its anterior position has been disguised by the reconstruction of the supra-pubic escutcheon.

Figure 12.18 Clinical photograph to show the procedentia that can be associated with exstrophy.

Chapter 13

Figure 13.1 (a) The external anatomy of the perineum in cloacal exstrophy to show the relationship of the two portions of the penis to the pubic bones. (b) Internal pelvic anatomy of the same specimen.

Figure 13.2 (a) Clinical photograph of a man with cloacal exstrophy to show the rudimentary penis. (b) Clinical photograph of the same patient to show the result of a forearm phalloplasty. The native penis is arrowed.

Figure 13.3 Clinical photograph of a 46XY adolescent raised as a female to show the external appearance of the perineum. The two halves of the penis have been joined to form a clitoris. The scrotal skin was used to form labia, but despite the absence of testes, it does not give a convincing female genital appearance.

Figure 13.4 Clinical photograph of the perineum of a woman with cloacal exstrophy. She had a didelphic uterus but the right side was rudimentary. The cervices had descended to the introitus (black arrows). The left half of the clitoris is visible (C).

Chapter 14

Figure 14.1 Histogram to show the percentage incidence of detrusor instability and postvoid residual urine by age in years in boys treated for posterior urethral valves.

Figure 14.2 (a) Micturating cystogram of a boy after valve resection showing a dilated prostatic urethra (PU). The arrow indicates the bladder neck.

(b) Transrectal ultrasound scan of an adult complaining of slow ejaculation showing a dilated prostatic urethra (PU) and a prostatic duct (D).

Figure 14.3 Cystoscopic photograph of the distal prostatic urethra in an adult. The vertical arrow (VR) indicates an insignificant portion of valve remnant. The area between the horizontal arrows is an area of scarring, presumably from the resection of the valve.

Figure 14.4 Cystoscopic photograph of the distal prostatic urethra in another adult. The veru montanum is normal. The upper arrow indicates a slightly dilated duct. The lower arrow indicates a very large dilated duct.

Chapter 15

Figure 15.1 Clinical photograph of BXO in an adult with untreated distal hypospadias. Note the white thickening around the stenotic meatus.

Figure 15.2 Clinical photograph of an adult with untreated hypospadias showing an external meatus

almost at the tip of the penis. However, the spongiosum is missing from the distal 2 cm (as far as the white arrow) shown by the sound in the urethra. This fragile tissue would easily be damaged by instrumentation.

Figure 15.3 Clinical photograph to show chordee in an adult with untreated coronal hypospadias.

Figure 15.4 Urethrogram to show a poor result. N = normal urethra, S = stricture, D = anterior diverticulum.

Figure 15.5 A hard hair ball removed from a reconstructed urethra.

Figure 15.6 Clinical photograph of a penis in an adult with a comprehensively failed reconstruction. The meatus is not terminal, the skin is heavily scarred and the urethra is strictured.

Chapter 16

Figure 16.1 MRI scan of a man with bilateral intra-abdominal testes (arrowed).

Chapter 17

Figure 17.1 (a) Clinical photograph of the clitoris in an unoperated woman with CAH. There is moderate enlargement but it can be at least partly covered by the labia. (b) Clinical photograph of a Prader V clitoris in a woman with untreated CAH. Note that despite the gross androgenization, she still has a female body shape and some breast development.

Figure 17.2 (a) Clinical photograph showing a poor outcome from feminizing genitoplasty. The clitoris has been buried and the urogenital sinus has not been formed into a urethra and vagina. (b) Clinical

photograph to show the excess buried corporeal tissue that was removed.

Figure 17.3 Diagram of a perineum to show the mean level of sexual sensitivity and intensity of orgasm when that area is stimulated. The depth of color indicates the degree of sensitivity.

Figure 17.4 Sagittal view of an MRI in a woman with an obliterated lower vagina to show the dilated vagina. B is the bladder.

Figure 17.5 Diagram to show the formation of a lower vagina by folding downwards a flap of the dilated vagina and tubularizing it, as used in the woman in Figure 17.4.

Figure 17.6 (a) Diagram of the formation of a neovagina from ileum using the Monti principle A—a segment of ileum has been isolated on a pedicle. B—The isolated segment is opened longitudinally close to the pedicle. C—it is reformed as a tube by closing transversely. The pedicle will then be at the top of the neovagina which makes it easier to reach the perineum. (Courtesy of Dan Wood, with permission.) (b) Clinical photograph of the operation at stage C of the diagram: P = pedicle, NV = neovagina.

Chapter 18

Figure 18.1 Clinical photograph of the penis of an adult 46XX patient raised as a male.

Figure 18.2 90th, 50th and 10th centile lines for the stretched penis from birth to maturity. Data from 1500 boys and men in the United States, mainly Caucasian.

Figure 18.3 Clinical photographs of patients with micropenis (a) in childhood and (b) after puberty.

Figure 18.4 Clinical photograph of a patient with PAIS raised as a female and treated with oestrogens. The upper arrow indicates the clitoris; the middle arrow is the urethra; the lower arrow is the vaginal open. There is scarring from surgery in childhood and, as so often is the case, the vaginal surgery needs revision, but the other external genitalia are well oestrogenized. Note the scanty pubic hair and absence of the anterior fourchette.

Figure 18.5 Clinical photograph of an adult with 5-alpha reductase deficiency. Note the scrotal scars from orchiopexy, bifid scrotum, and poorly developed penis.

Figure 18.6 (a) Sagittal MRI scan of a phenotypic 46XY male presenting with a large abdominal mass. He had PMDS. (b) The excised specimen consisting of bilateral testicular seminomas along with a uterus and Fallopian tubes. (Ut = uterus. Blad = bladder).

Figure 18.7 Lateral view of a micturating cystogram showing a large Mullerian remnant in an otherwise normal man presenting with pelvic infection. (MU = Mullerian remnant).

Chapter 20

Figure 20.1 Histogram comparing the actual and the expected number of deaths by length of follow up in childhood cancer patients who survived at least 5 years.

To Anna, my wife, who has supported this project and my
entire career.

Hers was the inspiration for the cover design

Adolescent urology and long-term outcomes

Christopher Woodhouse MB, BS, FRCS, FEBU
Emeritus Professor of Adolescent Urology,
University College, London, UK

With a chapter by Dr Kate Hillman and
Professor Guy Neild

WILEY Blackwell

This edition first published 2015 © 2015 by John Wiley & Sons, Ltd.

Registered Office

John Wiley & Sons, Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Offices

9600 Garsington Road, Oxford, OX4 2DQ, UK

The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

111 River Street, Hoboken, NJ 07030-5774, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell

The right of the author to be identified as the author of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting a specific method, diagnosis, or treatment by health science practitioners for any particular patient. The publisher and the author make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of fitness for a particular purpose. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. Readers should consult with a specialist where appropriate. The fact that an organization or Website is referred to in this work as a citation and/or a potential source of further information does not mean that the author or the publisher endorses the information the organization or Website may provide or recommendations it may make. Further, readers should

be aware that Internet Websites listed in this work may have changed or disappeared between when this work was written and when it is read. No warranty may be created or extended by any promotional statements for this work. Neither the publisher nor the author shall be liable for any damages arising herefrom.

Library of Congress Cataloging-in-Publication Data applied for.

A catalogue record for this book is available from the British Library.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Cover credit: sand-in-sahrara-vertical image: ©iStock.com/ermingut