



MUSCULOSKELETAL EXAMINATION

4TH EDITION



JEFFREY M. GROSS
JOSEPH FETTO
ELAINE ROSEN



WILEY Blackwell

This title is also available as an e-book.
For more details, please see
www.wiley.com/buy/9781118962763
or scan this QR code:



Musculoskeletal Examination

Fourth Edition

Jeffrey M. Gross, MD

Clinical Assistant Professor of Rehabilitation Medicine
Weill-Cornell Medical College
Adjunct Clinical Associate Professor of Rehabilitation
Medicine
New York University School of Medicine
Medical Director
Union Square Rehabilitation and Sports Medicine
New York

Joseph Fetto, MD

Associate Professor of Orthopedic Surgery
New York University School of Medicine
Associate Professor and Consultant
Manhattan V.A. Medical Center
New York

Elaine Rosen, PT, DHSc, OCS

Professor of Physical Therapy
Hunter College

City University of New York
Partner
Queens Physical Therapy Associates
Forest Hills, New York

WILEY Blackwell

This edition first published 2016 © 2016 by John Wiley & Sons, Ltd
Previous editions: 1996, 2009 by Jeffrey Gross, Joseph Fetto, Elaine Rosen

Registered office: John Wiley & Sons, Ltd, The Atrium, Southern Gate,
Chichester, West Sussex, PO19 8SQ, UK

Editorial offices: 9600 Garsington Road, Oxford, OX4 2DQ, UK
The Atrium, Southern Gate, Chichester, West Sussex,
PO19 8SQ, UK

111 River Street, Hoboken, NJ 07030-5774, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell

The right of the author to be identified as the author of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting a specific method, diagnosis, or treatment by health science practitioners for any particular patient. The publisher and the author make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of fitness for a particular purpose. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. Readers should consult with a specialist where appropriate. The fact that an organization or Website is referred to in this work as a citation and/or a potential source of further information does not mean that the author or the publisher endorses the information the organization or Website may provide or recommendations it may make. Further, readers should

be aware that Internet Websites listed in this work may have changed or disappeared between when this work was written and when it is read. No warranty may be created or extended by any promotional statements for this work. Neither the publisher nor the author shall be liable for any damages arising herefrom.

Library of Congress Cataloging-in-Publication Data

Gross, Jeffrey M., 1957–, author.

Musculoskeletal examination / Jeffrey M. Gross, Joseph Fetto, Elaine Rosen. – 4th edition.

p. ; cm.

Includes bibliographical references and index.

ISBN 978-1-118-96276-3 (pbk.)

I. Fetto, Joseph, author. II. Rosen, Elaine, author. III. Title.

[DNLM: 1. Musculoskeletal Diseases–diagnosis. 2. Musculoskeletal Physiological Phenomena. 3. Musculoskeletal System–anatomy & histology. 4. Physical Examination–methods. WE 141]

RC925.7

616.70076–dc23

2015000697

A catalogue record for this book is available from the British Library.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Cover image: istock (photo/muscles-of-the-arm-34748034) 02-09-14 (c) Eraxion

Contents

[How to Use This Book](#)

[Acknowledgments](#)

[About the Companion Website](#)

[Chapter 1 Introduction](#)

[What Is a Physical Examination?](#)

[What Is the Purpose of the Physical Examination?](#)

[How Is the Physical Examination Useful?](#)

[How Does the Musculoskeletal System Work?](#)

[What Are Paradigms?](#)

[What Are the Components of the Musculoskeletal System?](#)

[Bone](#)

[Cartilage](#)

[Ligaments](#)

[Muscle](#)

[Tendons](#)

[Synovium and Bursae](#)

[Fascia](#)

[The Interaction of Connective Tissues](#)

[Chapter 2 Basic Concepts of Physical Examination](#)

[Introduction](#)

[Observation](#)

[Subjective Examination \(History\)](#)

[Objective Examination](#)

[Anterior View](#)

[Correlation](#)

Chapter 3 Overview of the Spine and Pelvis

Chapter 4 The Cervical Spine and Thoracic Spine

Observation

Subjective Examination

Gentle Palpation

Trigger Points of the Cervical Spine

Active Movement Testing

Passive Movement Testing

Resistive Testing

Neurological Examination of the Cervical Spine and Upper Extremity

Special Tests

Upper Cervical Instability Testing

Alar Ligament Stress Tests

Referred Pain Patterns

Radiological Views

Chapter 5 The Temporomandibular Joint

Functional Anatomy of the Temporomandibular Joint

Observation

Subjective Examination

Special Questions

Gentle Palpation

Trigger Points of the TMJ Region

Active Movement Testing

Passive Movement Testing

Resistive Testing

Reflex Testing

Chapter 6 The Lumbosacral Spine

Observation

Subjective Examination

Gentle Palpation

Trigger Points of the Lumbosacral Region

Active Movement Testing

Forward Bending

Passive Movement Testing

Resistive Testing

Neurological Testing

Special Tests

Radiological Views

Chapter 7 Overview of the Upper Extremity

Chapter 8 The Shoulder

Functional Anatomy

Observation

Subjective Examination

Gentle Palpation

Trigger Points of the Shoulder Region

Active Movement Testing

Passive Movement Testing

Resistive Testing

Neurological Examination

Special Tests

Referred Pain Patterns

Radiological Views

Chapter 9 The Elbow

Functional Anatomy

Observation

Subjective Examination

Gentle Palpation

Resistive Testing

Elbow Flexion

Elbow Extension

Forearm Pronation

Forearm Supination

Neurological Examination

Special Tests

Referred Pain Patterns

Radiological Views

Chapter 10 The Wrist and Hand

Functional Anatomy

Observation

Subjective Examination

Gentle Palpation

Active Movement Testing

Passive Movement Testing

Resistive Testing

Neurological Examination

Special Tests

Referred Pain Patterns

Radiological Views

Chapter 11 The Hip

Functional Anatomy

Observation

Subjective Examination

Gentle Palpation

Anterior Aspect—The Patient Is Positioned in Supine

Posterior Aspect—The Patient Is Positioned in Prone

Side-Lying Position

Trigger Points

Active Movement Testing

Flexion

Extension

Abduction

Adduction

Medial (Internal) Rotation

Lateral (External) Rotation

Passive Movement Testing

Physiological Movements

Resistive Testing

Neurological Examination

Special Tests

Tests for Integrity of the Hip Joint for Degenerative
Joint Disease (DJD) or Labral Tear

Radiological Views

Chapter 12 The Knee

Functional Anatomy

Observation

Subjective Examination

Gentle Palpation

Trigger Points

Active Movement Testing

Passive Movement Testing

Resistive Testing

[Neurological Examination](#)
[Special Tests](#)
[Radiological Views](#)
[Chapter 13 The Ankle and Foot](#)
[Functional Anatomy](#)
[Observation](#)
[Subjective Examination](#)
[Gentle Palpation](#)
[Active Movement Testing](#)
[Passive Movement Testing](#)
[Resistive Testing](#)
[Neurological Examination](#)
[Referred Pain Patterns](#)
[Special Tests](#)
[Tests for Alignment](#)
[Radiological Views](#)
[Chapter 14 Gait](#)
[The Lower Extremity](#)
[What Is Gait?](#)
[What Is Normal Gait?](#)
[The Examination of Abnormal Gait](#)
[Appendices](#)
[Bibliography](#)
[Index](#)
[EULA](#)

List of Tables

[Chapter 1](#)

Table 1.1

Table 1.2

Chapter 6

Table 6.1

Chapter 8

Table 8.1

Chapter 9

Table 9.1

Chapter 10

Table 10.1

Table 10.2

Chapter 11

Table 11.1

Chapter 12

Table 12.1

Chapter 13

Table 13.1

Chapter 14

Table 14.1

Appendices

Appendix A

Appendix B

List of Illustrations

Chapter 1

Figure 1.1 Biological systems, like inorganic systems, can fail under one of two modes: an acute single supramaximal stress or repetitive submaximal chronic loading.

Figure 1.2 The “vicious cycle of injury” results from the reinjury of a vulnerable, recently traumatized system. This increased vulnerability occurs due to a diminishing of a system's tolerance limit as a result of adaptation to a lower level of demand during the period of rest necessitated by pain.

Figure 1.3 Conditioning is the adaptation of a biological system to the controlled application of increasing stress at a frequency, intensity, and duration within the system's tolerance limit, with a resultant increase in the system's tolerance limit.

Figure 1.4 (a) Collagen is a linear protein made of α chains that wind into a triple-helix. (b) Collagen fibrils are formed by the cross-linking of collagen monomer proteins. (c) The different types of collagen are determined by the number of $\alpha 1$ and $\alpha 2$ collagen monomers that join to form a triple-helix collagen molecule. For example, two $\alpha 1$ chains and one $\alpha 2$ chain that join to form a triple-helix make type I collagen, which is found in bone, tendon, ligament, fascia, skin, arteries, and the uterus. Type II collagen, which is found in articular cartilage, contains three $\alpha 1$ chains. There are at least 12 different collagen types.

Figure 1.5 The proteoglycan aggregate is formed on a backbone of hyaluronic acid and has the appearance of a bristle brush.

Figure 1.6 The structure of a typical long bone.

Figure 1.7 The composition and structure of articular hyaline cartilage. Water moves in and out of the cartilage due to the pressure of the joint surfaces on one another and attraction of the water by the ground substance. Note the orientation of the collagen fibers.

Figure 1.8 The ligaments of the knee. Because of the inherent instability of the joint, ligaments are necessary to prevent motion in all planes. They act as the primary stabilizers of the joint and are assisted by the muscles and other connective tissues.

Figure 1.9 The mechanical response of stress and strain on the anterior longitudinal ligament and the ligamentum flavum. The anterior cruciate ligament, having more collagen than elastin, can handle a larger load but will only stretch a short amount before breaking. The ligamentum flava, having more elastin than collagen, cannot tolerate a very large load but can stretch a lot before breaking.

Figure 1.10 A microscopic view of muscle shows the repeated patterns of the sarcomeres and the fibrils.

Figure 1.11 The organization of skeletal muscle tissue.

Figure 1.12 Different types of muscle-fascicle arrangements.

Figure 1.13 A tendon.

Figure 1.14 The olecranon bursa is between the skin and the olecranon process at the elbow.

Chapter 2

Figure 2.1 Body chart.

Figure 2.2 Normal posterior view.

Figure 2.3 Calcaneal valgus deformity.

Figure 2.4 Genu varum (a) and valgum (b).
deformities.

Figure 2.5 Scoliosis.

Figure 2.6 Rounded thoracic kyphosis.

Figure 2.7 Abducted scapula.

Figure 2.8 Winged scapula.

Figure 2.9 Sprengel's deformity.

Figure 2.10 Normal anterior view.

Figure 2.11 Pes planus deformity.

Figure 2.12 Hammertoe deformity.

Figure 2.13 Hallux valgus deformity.

Figure 2.14 Squinting patellae. The patellae face
each other.

Figure 2.15 Genu varum (a) and valgum (b).
deformities.

Figure 2.16 Barrel chest deformity.

Figure 2.17 Torticollis.

Figure 2.18 Normal lateral view.

Figure 2.19 Normal medial longitudinal arch.

Figure 2.20 Genu recurvatum deformity.

Figure 2.21 Flat back deformity.

Figure 2.22 Sway back deformity.

Figure 2.23 Dowager's hump deformity.

Figure 2.24 Rounded shoulders.

Figure 2.25 Forward head posture.

Figure 2.26 Movement diagram.

Chapter 3

Figure 3.1 A generalized vertebral segment is composed of a large, solid cylindrical body. Posteriorly, there is a bony ring through which the spinal cord and its coverings pass. Transverse and spinous processes project from the ring.

Figure 3.2 A transverse view through the ring of C1 (atlas) shows the spinal canal to be divided into thirds. The anterior third is occupied by the bony odontoid process of C2; the posterior third is filled by the spinal cord; the transverse ligament prevents migration of C1 on C2, not allowing the odontoid to invade the empty central third of the spinal canal.

Figure 3.3 The skull rests on the atlas (C1); the head rotates about the odontoid process as if it were a gatepost.

Chapter 4

Figure 4.1 Overview of the neck with anterior-posterior relationships.

Figure 4.2 Overview of the posterior thorax.

Figure 4.3 Overview of the skull.

Figure 4.4 The sternocleidomastoid muscle acts both as a cervical flexor and lateral rotator of the cervical spine. The scaleni muscles act to bend the cervical spine laterally and also assist in flexion.

Figure 4.5 Palpation of the inion.

Figure 4.6 Palpation of the superior nuchal line.

Figure 4.7 Palpation of the occiput.

Figure 4.8 Palpation of the mastoid process.

Figure 4.9 Palpation of the transverse process of C1.

Figure 4.10 Palpation of the spinous process of C2.

Figure 4.11 Palpation of the spinous processes.

Figure 4.12 Palpation of the spinous process of C7.

Figure 4.13 Palpation of the articular pillar.

Figure 4.14 Palpation of the transverse processes of the cervical spine.

Figure 4.15 Palpation of the spinous processes of the thoracic spine.

Figure 4.16 Palpation of the transverse processes of the thoracic spine.

Figure 4.17 Palpation of the spine of the scapula.

Figure 4.18 Palpation of the medial border of the scapula.

Figure 4.19 Palpation of the trapezius muscle.

Figure 4.20 Palpation of the suboccipital muscles and the greater occipital nerves.

Figure 4.21 Palpation of the ligamentum nuchae.

Figure 4.22 Palpation of the hyoid bone.

Figure 4.23 Palpation of the thyroid cartilage and gland.

Figure 4.24 Palpation of the first cricoid ring.

Figure 4.25 Palpation of the carotid tubercle.

Figure 4.26 Palpation of the suprasternal notch.

Figure 4.27 Palpation of the sternal angle.

Figure 4.28 Palpation of the sternoclavicular joint.

Figure 4.29 Palpation of the clavicle.

Figure 4.30 Palpation of the first rib.

Figure 4.31 Palpation of the ribs.

Figure 4.32 Palpation of the sternocleidomastoid muscle and the scaleni muscles.

Figure 4.33 Palpation of the lymph node chain.

Figure 4.34 Palpation of the carotid pulse.

Figure 4.35 Palpation of the parotid gland.

Figure 4.36 A trigger point in the upper trapezius muscle may cause headaches. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 4.37 Trigger points in the middle and lower trapezius may cause pain in the occipital region and along the paraspinal region. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 4.38 Additional trigger points in the left lower and right middle trapezius are shown with their referred pain patterns. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 4.39 Trigger points in the sternocleidomastoid muscle may cause referred pain in the face and head and also symptoms of watery eyes and runny nose. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 4.40 Trigger points within the scalene muscles may refer pain all the way to the hand. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 4.41 Trigger points in the suboccipital muscles radiate pain in the region of the greater occipital nerve. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 4.42 A trigger point in the splenius capitis muscle may cause referred pain on the top of the head. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 4.43 Active movement testing. (a) Cervical forward bending. (b) Cervical backward bending. (c) Cervical sidebending. (d) Cervical rotation.

Figure 4.44 Mobility testing of cervical spine flexion.

Figure 4.45 Mobility testing of cervical spine lateral (side) bending.

Figure 4.46 Mobility testing of cervical spine rotation.

Figure 4.47 Cervical spine traction.

Figure 4.48 (a, b) Mobility testing of central posteroanterior pressure on the spinous processes.

Figure 4.49 (a, b) Mobility testing of the posteroanterior pressure on the articular pillar.

Figure 4.50 (a, b) Mobility testing of transverse pressure on the spinous process.

Figure 4.51 Mobility testing of first rib ventral-caudal glide.

Figure 4.52 Testing cervical flexion.

Figure 4.53 The cervical extensors.

Figure 4.54 Testing cervical extension.

Figure 4.55 Testing lateral rotation. Resisting rotation of the head to the left tests the right sternocleidomastoid muscle.

Figure 4.56 Testing lateral bending.

Figure 4.57 The brachial plexus. The anterior divisions of the upper and middle trunks form the lateral cord and the anterior division of the lower trunk forms the medial cord. The three posterior divisions of the trunk form the posterior cord.

Figure 4.58 The median nerve stretch test. (Adapted from Butler, 1991.)

Figure 4.59 The radial nerve stretch test.

Figure 4.60 The ulnar nerve stretch test. (Adapted from Butler, 1991.)

Figure 4.61 The C5 root level.

Figure 4.62 The C6 root level.

Figure 4.63 The C7 root level.

Figure 4.64 The C8 root level.

Figure 4.65 The T1 root level.

Figure 4.66 The thoracic dermatomes and their key sensory areas.

Figure 4.67 The Spurling test. The patient's head is flexed laterally. Compression causes the neural foramina on the same side to narrow in diameter.

Figure 4.68 The distraction test. Distracting the cervical spine increases the diameter of the neural foramina.

Figure 4.69 Sharp-Purser test. Stabilize at C2 and translate the head posteriorly.

Figure 4.70 Aspinall's transverse ligament test. Direct force gradually from posterior to anterior.

Figure 4.71 Lateral flexion alar ligament stress test. Attempt passive sidebending of the head in neutral, flexion, and extension.

Figure 4.72 Rotational alar ligament stress test. Stabilize at C2. Passively rotate the head, first to the asymptomatic side and then to the other side.

Figure 4.73 Lhermitte's sign. Do not apply any pressure with your right hand. Gently bend the head forward to stretch the dura.

Figure 4.74 Vertebral artery test. This test should be performed if cervical manipulation is being contemplated.

Figure 4.75 Shoulder abduction (relief) test (Bakody's sign). Radicular pain is relieved by this position.

Figure 4.76 The scalp, ears, face, jaw, teeth, and throat may all refer pain to the cervical spine.

Figure 4.77 Anteroposterior view of the cervical spine.

Figure 4.78 Lateral view of the cervical spine.

Figure 4.79 Oblique view of the cervical spine.

Figure 4.80 Magnetic resonance image of the cervical spine, sagittal view.

Chapter 5

Figure 5.1 Palpation of the mandible.

Figure 5.2 Palpation of the temporalis muscle.

Figure 5.3 Palpation of the pterygoid muscles.

Figure 5.4 Palpation of the masseter muscle.

Figure 5.5 Palpation of the suprahyoid (a) and infrahyoid muscles (b).

Figure 5.6 Trigger points of the lateral pterygoid, shown with common areas of referred pain.

Figure 5.7 Trigger points of the masseter, shown with common areas of referred pain.

Figure 5.8 Trigger points of the temporalis, shown with common areas of referred pain.

Figure 5.9 Trigger points of the medial pterygoid, shown with common areas of referred pain.

Figure 5.10 Observe as the patient opens their mouth as far as they can. Both TMJs should be working simultaneously and synchronously allowing the mandible to open evenly without deviation to one side. A quick functional test is performed by asking the patient to place two to three flexed fingers, at their knuckles, between the upper and lower teeth.

Figure 5.11 Observe as the patient juts the jaw anteriorly so that it protrudes out from the upper teeth.

Figure 5.12 Observe as the patient disengages his or her bite and then moves the mandible first to one side, back to the midline, and then to the other side.

Figure 5.13 The freeway space is the point within the open pack position where the soft tissues of the temporomandibular joints are the most relaxed.

Figure 5.14 Overbite is the point where the maxillary teeth overlap the mandibular teeth. Overjet is the distance that the maxillary teeth protrude anteriorly over the mandibular teeth.

Figure 5.15 Measurement of the mandible is taken from the back of the TMJ to the notch of the chin. Compare the measurement of both sides.

Figure 5.16 The normal position of the tongue is at the top of the palate.

Figure 5.17 Mobility testing of distraction of the temporomandibular joint.

Figure 5.18 Lateral and medial pterygoid muscles.

Figure 5.19 Masseter.

Figure 5.20 Temporalis.

Figure 5.21 Jaw jerk. Make sure the patient is relaxed.

Chapter 6

Figure 6.1 Palpation of the iliac crest.

Figure 6.2 Palpation of the spinous processes.

Figure 6.3 Palpation of the transverse processes.

Figure 6.4 Palpation of the posterior superior iliac spine (PSIS).

Figure 6.5 Palpation of the sacroiliac joint.

Figure 6.6 Palpation of the sacral base.

Figure 6.7 Palpation of the inferior lateral angle.

Figure 6.8 Palpation of the ischial tuberosity.

Figure 6.9 Palpation of the coccyx.

Figure 6.10 Palpation of the supraspinous ligament.

Figure 6.11 Palpation of the erector spinae muscles.

Figure 6.12 Palpation of the quadratus lumborum muscle.

Figure 6.13 Palpation of the sacrotuberous ligament.

Figure 6.14 Palpation of the piriformis muscle.

Figure 6.15 Palpation of the sciatic nerve.

Figure 6.16 Palpation of the anterior superior iliac spine (ASIS).

Figure 6.17 Palpation of the pubic tubercles.

Figure 6.18 Palpation of the abdominal muscles.

Figure 6.19 Palpation of the psoas muscle.

Figure 6.20 Trigger points in the rectus abdominis may simulate the pain of dysmenorrhea. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 6.21 Trigger points in the rectus abdominis may also radiate pain into the posterior lower part of the thorax and lower back. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 6.22 Trigger points within the multifidi muscles may cause referred pain in the paraspinal region. Pain may also radiate anteriorly or inferiorly. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 6.23 Trigger points in the iliocostalis thoracis muscle may radiate pain superiorly and inferiorly as well as anteriorly. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 6.24 Trigger points in the iliocostalis lumborum and longissimus thoracis muscles radiate pain inferiorly. (Adapted with permission from Travell and Rinzler, 1952.)

Figure 6.25 Pelvic, abdominal, and retroperitoneal organs may radiate pain to the lumbar spine. The hip may also cause low-back pain.

Figure 6.26 Active movement testing. (a) Lumbar forward bending. (b) Lumbar backward bending. (c) Lumbar side bending. (d) Lumbar rotation.

Figure 6.27 Mobility testing of lumbar spine flexion.

Figure 6.28 Mobility testing of lumbar spine side bending.

Figure 6.29 Mobility testing of lumbar spine rotation.

Figure 6.30 (a, b) Mobility testing of central posteroanterior spring on the spinous process.

Figure 6.31 (a, b) Mobility testing of central posteroanterior spring on the transverse process.

Figure 6.32 (a, b) Mobility testing of transverse pressure on the spinous process.

Figure 6.33 Mobility testing of the sacroiliac joint: standing forward-bending test.

Figure 6.34 Mobility testing of the sacroiliac joint: stork test.

Figure 6.35 Mobility testing of the sacroiliac joint: backward-bending test.

Figure 6.36 Mobility testing of the sacroiliac joint: sitting forward-bending test.

Figure 6.37 (a, b) Mobility testing of the sacroiliac joint: posteroanterior spring of the sacrum.

Figure 6.38 The trunk flexors.

Figure 6.39 Testing trunk flexion.

Figure 6.40 The trunk rotators.

Figure 6.41 Testing trunk rotation.

Figure 6.42 The trunk extensors.

Figure 6.43 Testing trunk extension.

Figure 6.44 The lumbar plexus. The lumbar plexus is formed by the ventral primary rami of L1, L2, L3, and L4 and possibly T12. Note that the peripheral nerves

from the anterior divisions innervate the adductor muscles of the hip, and the peripheral nerves from the posterior divisions innervate the hip flexors and knee extensors.

Figure 6.45 The lumbosacral plexus. This plexus is formed by the ventral primary rami of L4, L5, S1, S2, and S3.

Figure 6.46 The dermatomes and key sensory areas of the lower extremity.

Figure 6.47 The cutaneous innervation of a lower limb.

Figure 6.48 The L1 and L2 root levels.

Figure 6.49 The L3 root level.

Figure 6.50 The L4 root level.

Figure 6.51 The L5 root level.

Figure 6.52 The S1 root level.

Figure 6.53 The upper abdominal skin reflex (T5–T8).

Figure 6.54 The mid-abdominal skin reflex (T9–T11).

Figure 6.55 The lower abdominal skin reflex (T11, T12).

Figure 6.56 The cremasteric reflex. Note that immediate movement of the scrotum upward is a positive test result.

Figure 6.57 A posterolateral herniation of the L4–L5 disc can cause pressure and injury to the L5 nerve root.

Figure 6.58 A posterolateral herniation of the L5–S1 disc can cause pressure and injury to the L5 and S1 nerve roots.

Figure 6.59 Straight-leg-raising test. Between 35 and 70 degrees, the L5 and S1 nerve roots may be stretched against an intervertebral disc. Flexing the hip more than 70 degrees causes stress on the lumbar spine.

Figure 6.60 By lowering the leg slightly to the point where the patient stops feeling pain or paresthesias in the leg, and then dorsiflexing the ankle, you can determine whether the pain in the leg is due to tight hamstrings or has a neurogenic origin. If the pain is reproduced on dorsiflexion of the ankle after the hamstrings have been relaxed by lowering the leg slightly, the pain has a neurogenic origin.

Figure 6.61 The slump test. (Adapted from Butler, 1991.)

Figure 6.62 The femoral stretch test. The test leg is extended at the hip first, and then the knee is flexed.

Figure 6.63 Valsalva maneuver.

Figure 6.64 Gaenslen's sign. Bring the patient to the edge of the table with the test-side buttock over the edge.

Figure 6.65 Allow the patient's thigh and leg to move downward to stress the sacroiliac joint on that side. Pain on this maneuver reflects sacroiliac joint pathology.

Figure 6.66 Patrick's or Fabere test. The hip is flexed, abducted, externally rotated, and extended.

Figure 6.67 The sacroiliac distraction test. By compressing the pelvis medially and distracting the sacroiliac joints, this test determines whether sacroiliac pathology is present.

Figure 6.68 Test for spondylolysis (extension in one-leg standing).

Figure 6.69 Anteroposterior view of the lumbosacral spine.

Figure 6.70 Lateral view of the lumbosacral spine.

Figure 6.71 Oblique view of the lumbosacral spine.

Figure 6.72 Magnetic resonance image of the lumbosacral spine, sagittal view.

Figure 6.73 Magnetic resonance image of the lumbosacral spine, transverse view.

Figure 6.74 Magnetic resonance view of the lumbosacral spine, sagittal view. DG, dorsal root ganglion.

Chapter 7

Figure 7.1 Overview of the upper extremity.

Chapter 8

Figure 8.1 Overview of the shoulder showing the importance of the soft tissues in maintaining the round humeral head in the flattened glenoid process of the scapula. The other joints of the shoulder are also shown.

Figure 8.2 The inferior glenohumeral ligament prevents further inferior movement of the humeral head as the arm abducts.

Figure 8.3 Inferior dislocation of the glenohumeral articulation, with attenuation of the inferior glenohumeral ligament.

Figure 8.4 Superior view of the shoulder, showing the acromioclavicular joint and the coracoacromial ligament overlying the humeral head.

Figure 8.5 The subacromial space is defined superiorly by the acromioclavicular bony arch and the coracoacromial ligament. Inferiorly, it is defined by the humeral head. Within this space lies the subacromial bursa and the tendons of the supraspinatus and long head of the biceps muscles.

Figure 8.6 Palpation of the suprasternal notch.

Figure 8.7 Palpation of the sternoclavicular joint.

Figure 8.8 Palpation of the clavicle.

Figure 8.9 Palpation of the acromioclavicular joint.

Figure 8.10 Palpation of the acromion process.

Figure 8.11 Palpation of the greater tubercle of the humerus.

Figure 8.12 Palpation of the coracoid process.

Figure 8.13 Palpation of the bicipital groove.

Figure 8.14 Palpation of the sternocleidomastoid muscle.

Figure 8.15 Palpation of the trapezius muscle.

Figure 8.16 Palpation of the pectoralis major muscle.

Figure 8.17 Palpation of the deltoid muscle.

Figure 8.18 Palpation of the biceps muscle.

Figure 8.19 Palpation of the spine of the scapula.

Figure 8.20 Palpation of the medial border of the scapula.

Figure 8.21 Palpation of the lateral border of the scapula.