

The Mediterranean Diet Health and Science

Richard Hoffman and Mariette Gerber



 WILEY-BLACKWELL

The Mediterranean Diet

Health and Science

The Mediterranean Diet

Health and Science

Richard Hoffman PhD, FRSA

Senior Lecturer, University of Hertfordshire, Hatfield,
Herts, UK

and

Mariette Gerber MD, PhD

Former INSERM Senior Scientist, Cancer Research Institute,
Montpellier, France

 **WILEY-BLACKWELL**

A John Wiley & Sons, Ltd., Publication

This edition first published 2012
© 2012 by Richard Hoffman and Mariette Gerber

Blackwell Publishing was acquired by John Wiley & Sons in February 2007. Blackwell's publishing program has been merged with Wiley's global Scientific, Technical and Medical business to form Wiley-Blackwell.

Registered Office

John Wiley & Sons, Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Offices

9600 Garsington Road, Oxford, OX4 2DQ, UK

The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

2121 State Avenue, Ames, Iowa 50014-8300, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell.

The right of the author to be identified as the author of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. This publication is designed to provide accurate and authoritative information in regard to the subject matter covered. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

Library of Congress Cataloging-in-Publication Data

Hoffman, Richard, 1957-

The Mediterranean diet : health and science / Richard Hoffman and Mariette Gerber.
p. cm.

Includes bibliographical references and index.

ISBN-13: 978-1-4443-3002-1 (pbk.)

ISBN-10: 1-4443-3002-0 (pbk.)

1. Nutrition. 2. Cooking, Mediterranean--Health aspects. 3. Biochemistry.

I. Gerber, Mariette. II. Title.

RM216.H685 2012

613.2--dc23

2011021992

A catalogue record for this book is available from the British Library.

Set in 10/12 pt Sabon by SPi Publisher Services, Pondicherry, India



Contents

Preface	xi
1 Overview	1
1.1 Development of the MedDiet	1
1.1.1 A brief history of the MedDiet	2
1.1.2 The traditional MedDiet and present day MedDiets	2
1.1.3 International differences	6
1.1.4 National representations of the MedDiet	7
1.1.5 Sources of information for the general public	13
1.2 Lifestyle factors	13
1.2.1 Meal patterns	14
1.2.2 Siestas	16
1.2.3 Physical activity	16
1.2.4 Sunshine	16
1.3 Health benefits	16
1.3.1 Mediterranean dietary patterns	17
1.3.2 Endorsements	19
1.4 The MedDiet, past, present and future	19
1.4.1 Current trends	19
1.4.2 The Greek experience	23
1.4.3 Not all bad news	24
1.4.4 Future prospects	25
References	25

SECTION 1: CONSTITUENTS

2 Constituents and Physiological Effects of Mediterranean Plant Foods	31
2.1 Introduction	31
2.2 Carbohydrates	32
2.2.1 Glycaemic index	32
2.2.2 Honey	34
2.2.3 Physiological effects of carbohydrates	35
2.2.4 Fibre	36
2.3 Fats	38
2.4 Organic acids	39
2.5 Minerals	40
2.6 Vitamins	40
2.6.1 Water soluble vitamins	40
2.6.2 Fat soluble vitamins	41
2.7 Phytochemicals	44

2.7.1	Classes of phytochemicals	46
2.7.2	Sensory properties of phytochemicals	46
2.7.3	Phenolics	50
2.7.4	Terpenes	56
2.7.5	Sulphur-containing compounds	62
2.7.6	Polyacetylenes	62
2.7.7	Nitrogen-containing compounds	63
2.8	Mediterranean plants as medicines	63
	References	64
3	Influences of the Supply Chain on the Composition of Mediterranean Plant Foods	67
3.1	Significance of the supply chain	67
3.2	Growth conditions	68
3.2.1	A brief overview of plant metabolism	68
3.2.2	Soil	70
3.2.3	Sun	71
3.2.4	Water	71
3.2.5	Other environmental factors	71
3.3	Plant cultivar	72
3.4	Food retailers and food processing	74
3.4.1	Anatomical distribution of nutrients	74
3.4.2	Freshness	75
	References	76
4	Influences of Food Preparation and Bioavailability on Nutritional Value	77
4.1	Introduction	77
4.2	Food preparation	78
4.2.1	Chopping	78
4.2.2	Cooking	79
4.3	Nutrient bioavailability	82
4.3.1	Pharmacokinetics of phytochemicals	82
4.3.2	Liberation	83
4.3.3	Absorption	83
4.3.4	Distribution	86
4.3.5	Metabolism	87
4.3.6	Excretion	90
4.3.7	Inter-individual variations in phytochemical pharmacokinetics	90
	References	92
5	Guide to the Composition of Mediterranean Plant Foods	94
5.1	Types of plant foods consumed as part of a MedDiet	95
5.1.1	Consumption	95
5.1.2	Diversity of consumption	95
5.2	Vegetables	96
5.2.1	Green leafy vegetables	96
5.2.2	Other green vegetables	102
5.2.3	Root vegetables	103
5.2.4	Alliums	104
5.2.5	Solanaceous vegetables	107
5.2.6	Legumes	109
5.3	Wheat products	112

5.4	Fruits	114
5.4.1	Citrus fruits	114
5.4.2	Apples and related fruits	118
5.4.3	Stone fruits	120
5.4.4	Grapes	121
5.4.5	Other berries	122
5.4.6	Pomegranates	122
5.4.7	Figs	123
5.4.8	Dates	123
5.4.9	Olives	123
5.5	Herbs and spices	127
5.6	Nuts and seeds	129
5.6.1	Nuts	129
5.6.2	Seeds	132
	References	132
6	Olive Oil and Other Fats	137
6.1	Overview	137
6.2	Olive oil	138
6.2.1	Consumption and production	138
6.2.2	Origin and varieties of olive trees	139
6.2.3	Olive oil production	139
6.2.4	Traceability and European regulations	143
6.2.5	Biochemical composition	144
6.3	Other fat sources	147
6.3.1	Nuts and seeds	147
6.3.2	Herbs and spices	148
6.3.3	Meat and dairy	148
6.3.4	Eggs	150
6.3.5	Fish and shellfish	151
	References	151
7	Wine and Other Drinks	154
7.1	Wine	154
7.1.1	Introduction	154
7.1.2	Production	155
7.1.3	Composition	156
7.1.4	Metabolism of wine	158
7.1.5	Consumption patterns and health	159
7.1.6	The resveratrol controversy	162
7.2	Aniseed-flavoured spirits	164
7.2.1	Consumption	164
7.2.2	Composition	165
7.2.3	Physiological effects	165
7.3	Tea	165
7.3.1	Consumption	165
7.3.2	Composition	166
7.3.3	Physiological effects	167
7.4	Coffee	168
7.4.1	Consumption	168
7.4.2	Composition and physiological effects	168
	References	169

SECTION 2: HEALTH EFFECTS

8	Epidemiological Methods	175
8.1	Introduction	175
8.2	Study designs	176
8.2.1	Descriptive epidemiology	176
8.2.2	Ecological studies	178
8.3	Analytical epidemiology	178
8.3.1	Case-control and prospective studies	179
8.4	Intervention studies	180
8.5	Expression and interpretation of data from epidemiological studies	180
8.6	Dietary patterns	181
8.6.1	<i>A priori</i> dietary patterns	181
8.6.2	<i>A posteriori</i> dietary patterns	183
8.7	Criteria for judging epidemiological data	184
	References	185
9	General Mechanisms for Disease Prevention	187
9.1	Introduction	188
9.2	Methods for studying the effects of nutrients on disease mechanisms	189
9.2.1	General considerations	189
9.2.2	<i>In vitro</i> models	190
9.2.3	Animal models	191
9.2.4	Human intervention studies	191
9.3	Oxidative stress	192
9.3.1	What are free radicals?	192
9.3.2	Production of reactive species	192
9.3.3	Effects of reactive species	193
9.3.4	Antioxidant defences	195
9.3.5	Measuring antioxidant activity	198
9.3.6	Antioxidant activity in the Mediterranean diet	199
9.3.7	Limitations of the dietary antioxidant theory	204
9.4	Inflammation	205
9.4.1	The inflammatory response	205
9.4.2	The MedDiet and inflammation	206
9.4.3	<i>n</i> -3 Fatty acids	207
9.4.4	Phenolics	210
9.4.5	Pro-inflammatory foods	212
9.5	Modulation of cell signalling pathways by phenolics	212
9.5.1	Cell signalling pathways in disease	212
9.6	Gene interactions	213
9.6.1	Genetic predisposition to diet – nutrigenetics	213
9.6.2	Effects of diet on gene expression	216
9.7	Increased stress resistance	222
9.7.1	The general concept	222
9.7.2	Mechanistic basis of hormesis	225
9.8	Nutrient interactions and the Mediterranean dietary pattern	226
9.8.1	Minimising spikes in plasma concentrations	227
9.8.2	Synergistic interactions	228
	References	229

10	Metabolic Disorders	234
10.1	Introduction	234
10.2	Obesity	235
10.2.1	Introduction	235
10.2.2	Epidemiology	237
10.2.3	Mechanisms	239
10.3	Metabolic syndrome	241
10.3.1	Introduction	241
10.3.2	Epidemiology	242
10.3.3	Mechanisms	245
10.4	Type 2 diabetes	249
10.4.1	Introduction	249
10.4.2	Epidemiology	250
10.4.3	Mechanisms	252
	References	254
11	Cardiovascular Diseases	258
11.1	Introduction	259
11.2	Nutrition and the biology of CVD	260
11.2.1	Atherogenesis	260
11.2.2	Cardiac arrhythmias	265
11.3	Epidemiological evidence and mechanisms	265
11.3.1	Fat and fatty acids	265
11.3.2	Plant food constituents	269
11.3.3	Whole foods	273
11.3.4	Dietary patterns	283
	References	286
12	Cancers	293
12.1	Introduction	293
12.2	Nutritional factors and the biology of cancer	294
12.2.1	Introduction	294
12.2.2	Carcinogenesis	294
12.2.3	The issue of selectivity	309
12.2.4	Mediterranean dietary pattern and cancer prevention	311
12.3	Epidemiological evidence and mechanisms	316
12.3.1	Fats	316
12.3.2	Plant food constituents	318
12.3.3	Tea	322
12.3.4	Coffee	323
12.3.5	Wine and alcohol	323
12.3.6	Plant foods	327
12.3.7	Dietary patterns	330
	References	334
13	Neurological and Other Disorders	343
13.1	Introduction	343
13.2	Dementias	344
13.2.1	Introduction	344
13.2.2	Epidemiology	344
13.2.3	Mechanisms	346

13.3	Parkinson's disease	349
13.3.1	Introduction	349
13.3.2	Epidemiology	349
13.3.3	Mechanisms	349
13.4	Depression	350
13.4.1	Introduction	350
13.4.2	Epidemiology	350
13.4.3	Mechanisms	350
13.5	Rheumatoid arthritis	351
13.5.1	Introduction	351
13.5.2	Epidemiology	351
13.5.3	Mechanisms	352
13.6	Age-related macular degeneration	352
13.6.1	Introduction	352
13.6.2	Epidemiology	353
13.6.3	Mechanisms	354
13.7	All cause mortality	354
13.7.1	Fruit and vegetables	355
13.7.2	Alcohol	355
13.7.3	Wine	355
13.7.4	Dietary pattern	356
13.8	General conclusions	357
	References	357
14	Public Health Issues	361
14.1	Introduction	361
14.1.1	Public health and its objectives	362
14.1.2	The challenges of public health in Europe	362
14.2	Which MedDiet?	366
14.3	Which constituents are important in the MedDiet?	367
14.3.1	Olive oil	367
14.3.2	Cereals and legumes	368
14.3.3	Fruit and vegetables	368
14.3.4	Herbs and spices	369
14.3.5	Meat and dairy products	370
14.3.6	Wine and tea	370
14.3.7	How can the Mediterranean dietary pattern be implemented?	371
14.4	Transferring the Mediterranean dietary pattern	372
14.4.1	Public health recommendations and education	374
14.4.2	Governments' and communities' food policies	375
14.4.3	Individual choices and attitudes	377
	References	378
 SECTION 3: APPENDICES		
Appendix 1	Abbreviations	383
Appendix 2	Epidemiological Studies	385
Index		389



Preface

This book is an introduction to the health benefits of the Mediterranean diet (MedDiet), and the epidemiological and experimental evidence that underpins these benefits. The book is mainly intended for dietitians and nutritionists, and will also be of interest to other health workers, food scientists, and students taking courses in biomedical sciences.

The MedDiet is best known for helping prevent cardiovascular diseases (CVDs) and, to a lesser extent, cancers, and these are discussed in detail. In addition, we discuss the increasing evidence that the MedDiet can help prevent other chronic degenerative diseases such as neurodegenerative disorders and metabolic syndrome. With the ageing of populations, these diseases have become the major cause of morbidity and mortality in both developed countries and many less-developed countries. This highlights the growing importance of nutrition in helping prevent these diseases, and the book concludes with a discussion of public health strategies that can facilitate implementing a MedDiet in the community, and so help increase the number of years that people live in good health.

Several chapters in the book discuss experimental evidence that contributes to understanding how the MedDiet works. In particular, we have discussed common disease mechanisms that underlie many chronic diseases, and how, by targeting these mechanisms, key nutrients in the MedDiet can help prevent a broad range of diseases. These key nutrients probably include various plant chemicals (phytochemicals), some vitamins, fibre and fatty acids. The levels of these nutrients – in Mediterranean foods as elsewhere – are influenced by many steps of the food chain, from food production to metabolic effects in the body. Hence, we have highlighted features of the food chain that can have a significant impact on levels of these nutrients and hence impact on the overall health benefits of a MedDiet. For example, phytochemical levels are influenced all along the food chain from decisions regarding plant cultivar selection, to the growth conditions of the plant, food preparation and metabolic effects in the body. Similarly, the feed of land animals and fish has a major impact on their fatty acid composition and hence this impacts on human nutrition as well.

Throughout the book we have emphasised the importance of considering the composition of the MedDiet in its totality, i.e. as a Mediterranean “dietary pattern”, rather than as a collection of individual foodstuffs. This is because both epidemiological evidence and – increasingly – underlying experimental studies, support the notion that the food groups (such as plant foods, fish and olive oil) comprising the totality of a MedDiet interact for optimum health. In fact, even this perspective is probably insufficient, since many other aspects associated with a traditional Mediterranean lifestyle can influence health. Best recognised of these is physical activity, but other features such as social eating, regular meal structure, sunlight, and even taking a

siesta, may contribute to the benefits of a MedDiet. By discussing these aspects, we hope to provide something of a counter-balance to some current nutritional trends which are promoting individual “superfoods” or “nutraceuticals” as strategies for reducing chronic degenerative disorders.

Research on the MedDiet is proceeding at a very rapid pace, and we have highlighted promising new areas of experimental research, some of which may in the future lead to a deeper understanding of the scientific basis of the MedDiet. What is evident even now is that the MedDiet is very much a diet for our times, both through its potential for improving the health and well-being of an ageing society, and its low environmental impact by eschewing the consumption of large amounts of meat from intensively raised animals and instead favouring foods, both plant and animal, produced under more natural conditions. These features promote a healthy attitude towards food, and help make the MedDiet very appealing to a large number of people. And its delicious tastes help ensure high levels of compliance amongst consumers.

1

Overview

Summary

- Olive oil is the main dietary fat in the Mediterranean diet (MedDiet), and there is high consumption of fruits, vegetables, unrefined cereals and legumes, moderate consumption of fish, dairy products and wine, and low consumption of meat.
- There are many variations of the MedDiet reflecting influences of culture and landscape.
- Factors such as social eating and physical activity are also important, so the MedDiet represents a whole lifestyle.
- There is strong epidemiological evidence for a protective effect by the MedDiet against cardiovascular diseases and also protection against cancers and neurological disorders. Many biologically plausible mechanisms have been developed to explain these health benefits.
- There is increasing evidence that the health benefits are best explained by considering the overall dietary pattern.
- Consumption of the MedDiet is decreasing in some Mediterranean countries, whereas consumption of Mediterranean foods in some North European countries is increasing.

1.1 Development of the MedDiet

The Mediterranean diet (MedDiet) is tasty, easy to prepare and extremely healthy. The ability of the MedDiet to help prevent a wide range of today's most common ailments has been confirmed in numerous studies, and more widespread implementation of a MedDiet would undoubtedly result in significant public health benefits. How has this cuisine become one of the healthiest in the world? Probably part of the answer lies in the enlightened understanding in the Mediterranean of the link between food and health. From the Ancient Greeks to the Moors onwards, the MedDiet has been developed not only for its gastronomic virtues, but also as a synonym for a complete system of life. Another important aspect of the MedDiet is related to climate. The long growing season in the Mediterranean reduces the need for complex preservation techniques, and this has permitted an emphasis on natural, seasonal ingredients. The reliance on fresh ingredients cooked from raw is probably the single most important factor in Mediterranean cuisine that helps to ensure the consumption of a balanced intake of nutrients. Many traditional cooking pots, such as the *tagine*, *paella* and *plancha*, are still widely used in Mediterranean countries

and have given their names to classic Mediterranean dishes. Although originally developed to enhance flavour, these vessels are employed in cooking techniques that help enhance the health benefits of the food. Indeed, flavour and health are closely interlinked in any cuisine based on natural ingredients. A third factor is the dominance of the olive and the eschewing of animal products rich in saturated fat. Cooking vegetables with olive oil is the quintessential Mediterranean way of obtaining the benefits of both taste and health. Hence, the MedDiet represents a legacy of the link between Man and the Mediterranean environment that has existed since the time of the last Ice Age.

1.1.1 A brief history of the MedDiet

The land surrounding the Mediterranean sea has been the cradle for many civilisations and cultures, and the MedDiet represents one of the most significant achievements of these civilisations. Agriculture itself began with the cultivation of cereals and pulses in the Levant, a region which comprises the Eastern Mediterranean countries of Lebanon, Israel, Palestine, Syria, Jordan and Iraq. Later, the Phoenicians, Greeks and Romans cultivated the three basic elements of the MedDiet: olive trees for producing olives and olive oil, wheat for making bread, and grapes for fermenting into wine. These colonisers of the Mediterranean basin then spread olives and grapes to the Western Mediterranean. A wide range of vegetables were already being consumed by the time of the Romans including onions, leeks, lettuce, carrots, asparagus, turnips, cabbage, celery and artichokes. Among the fruits consumed by this time were figs, apples, pears, cherries, plums, peaches, apricots and citrons (a type of citrus fruit). Chestnuts, almonds and walnuts were also eaten [1, 2].

Important developments in the MedDiet occurred from the 8th century when the Moors occupied much of the Iberian peninsula (calling the region al-Andalus). The Moors introduced rice, lemons, aubergines (American: egg plants), saffron and other spices, and these products then spread, to varying extents, throughout the Mediterranean basin. The Moors had a particularly enlightened awareness of the importance of diet for general health. For them, ‘diet was a synonym for a system of life. It included the practice of eating correctly, of choosing the best places for staying healthy and lengthening one’s life, of bathing and washing correctly, of sleeping and staying awake, of expelling useless substances from one’s body and of dealing with the ups and downs of the spirit’ [3]. The occupation by the Moors ended in 1492, and this was the same year that Christopher Columbus arrived in the New World. Columbus returned to Spain with tomatoes and bell peppers, and these are now an integral part of the MedDiet.

1.1.2 The traditional MedDiet and present day MedDiets

The term ‘Mediterranean diet’ was originally coined in the 1950s by Ancel Keys, the epidemiologist who first recognised the health benefits of this way of eating. Since then, the MedDiet has undergone many changes, and it is now convention to use the term ‘traditional’ MedDiet to indicate the type of diet that could be found in rural communities in the 1950s and early 1960s, especially in Southern Italy and Greece (and rural Crete in particular), and before the impacts of migrations to the towns, rising wealth and modern food technologies. The traditional MedDiet was shaped by terrain and climate. Cereals and vegetables were grown in the flatter,

low-lying areas, vines and olive trees on the slopes, and higher ground was left for grazing sheep and goats. The Cretan diet up until the 1960s has been described as ‘olives, cereals, grains, pulses, wild greens, herbs and fruits, together with limited quantities of goat meat and milk and fish ... no meal was complete without bread ... olives and olive oil contributed heavily to the energy intake ... food seemed to be “swimming’ in oil” [4].

There are various formulations of what is meant by the term ‘traditional MedDiet’, and the following list is taken from a statement issued by a working group at the MedDiet 2004 International Conference [5]:

- Olive oil as added lipid
- Daily consumption of vegetables
- Daily consumption of fruits
- Daily consumption of unrefined cereals
- Bi-weekly consumption of legumes
- Nuts and olives as snacks (generally eaten just before a meal)
- Bi-weekly consumption of fish
- Daily consumption of cheese or yogurt
- Monthly or weekly consumption of meat or meat products
- Daily moderate consumption of wine, if it is accepted by religion and social grounds

Although this list sums up most of the important aspects of a traditional MedDiet, other versions include consumption of herbs and spices, herbal teas, and wild greens gathered from the countryside, and also the importance of significant levels of physical activity.

The traditional MedDiet is a rich source of macronutrients and micronutrients. It is not possible to define the precise amounts of various beneficial nutrients in the MedDiet because of significant variations between countries (see below). However, one set of figures, based on an analysis of a traditional Greek diet, is shown in Table 1.1.

Fats are an important component of the traditional MedDiet, and account for about 30% of total calories in Spain, and up to about 40% of total calories in Greece [7]. This compares with about 34% in the American diet [8]. Hence, the traditional MedDiet is not a low fat diet. There are, however, significant differences in the fatty acid composition compared to a North European or a North American diet. This is mainly due to the fairly low level of saturated fats in the MedDiet (7–8% of total calories), and relatively high consumption of monounsaturated fatty acids (MUFAs) (>20% of total calories), which is mostly oleic acid derived from olive oil. Not only are total saturated fats relatively low in a traditional MedDiet, but the types of saturated fatty acids (SFAs) consumed are quantitatively different to those in a North European diet. This is partly because consumption of SFAs from meat and cow’s milk is relatively low, and consumption of SFAs from cheese and yogurt made from goat and sheep milk can be quite high. Goat and sheep milk contain a relatively high percentage of medium chain fatty acids (MCFAs) compared to cow’s milk, and these are not as strongly associated with adverse effects on plasma cholesterol levels as some longer chain SFAs. The Greeks have one of the highest consumption of cheese in the world – at 26 kg per person per year (2005 figures) it is even higher than for the French! But about half of this is feta, a cheese traditionally made with ewe’s milk and up to 30% goat milk.

Table 1.1 Estimated daily intake of macro- and micronutrients in a Greek MedDiet [6]. Reproduced with permission. © 2006 Elsevier.

Component	Daily intake
Macronutrients	
Protein	74.5 g
Carbohydrates	255.8 g
Dietary fibre	29.8 g
Ethanol	14 g
Total lipids	110.7 g
SFA	29.8 g
MUFA	63.8 g
PUFA	9.9 g
TFA	1.4 g
Phytochemicals	
Flavonoids	118.6 mg
Carotenoids	65.7 mg
Sterols	256.8 mg
α -tocopherol	4.3 mg
Inorganic constituents	
K	1774 mg
Fe	14.9 mg
Na	2632 mg
Ca	696 mg
Mg	234 mg
Zn	10.3 mg
Cu	3.8 mg
Mn	3.5 mg
Total energy value	2473 Kcal

SFA, saturated fatty acids; MUFA, monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; TFA, trans fatty acids.

Note: There were no data on vitamins in this analysis.

The traditional MedDiet is also a good source of polyunsaturated fats (PUFAs). Fish is the main contributor of the long chain (LC) *n*-3 FAs eicosapentaenoic acid (EPA) (20:5 *n*-3) and docosahexaenoic acid (DHA) (22:6 *n*-3). There is a modest intake of the *n*-6 fatty acid linoleic acid from nuts and sunflower seeds and pumpkin seeds, and these are popular aperitif foods in some Mediterranean countries. Linoleic acid is the predominant fatty acid in many seeds such as sunflower seeds and corn, and hence in oils made from these seeds. Seed oils are not a significant part of the traditional MedDiet and, as a consequence, *n*-6 fatty acid consumption is lower than in North Europe and North America. However, it should be mentioned that corn oil and sunflower oil are now increasingly replacing olive oil for cooking in some Mediterranean countries due to their lower cost.

Fats in the MedDiet

- High consumption of MUFAs, particularly oleic acid from olive oil
- High consumption of LC *n*-3 PUFAs (α -linolenic acid – ALA, EPA and DHA)
- Relatively low consumption of *n*-6 PUFAs
- Relatively high consumption of SCFAs and MCFAs from goat and sheep milk

Table 1.2 Estimate of the macronutrient composition of a typical MedDiet and a typical western diet (data from [9]).

Macronutrients	Mediterranean diet (%)	Western diet (%)
Carbohydrates	47	42
Proteins	15	20
Saturated fats	10	17
Monounsaturated fats	22	14
Polyunsaturated fats	6	7

Besides its typical fat composition, the MedDiet is also a rich source of a variety of carbohydrates, and these are discussed in Chapter 2. One estimate of how the overall proportions of macronutrients in a ‘typical’ western diet compare with those in a ‘typical’ MedDiet is shown in Table 1.2 [9]. This estimates MUFA intake in the MedDiet at 22% compared to 14% in a typical ‘Western’ diet. Although this analysis estimates that total PUFA intake between the two diets is similar, it should be noted that this analysis did not distinguish between *n*-6 and *n*-3 PUFAs.

Due to the high consumption of plants foods, the traditional MedDiet is a particularly rich source of plant chemicals (phytochemicals) and some vitamins and minerals (see Table 1.1). One aspect of particular relevance here is that plasma folate levels have been found to be a good biomarker for adherence to the MedDiet [10]. This mainly reflects the high consumption of green leafy vegetables in the MedDiet. Folate consumption is linked to a wide range of beneficial effects in the body, including prevention of neural tube defects in early pregnancy, and protection against cancers of the pancreas, oesophagus and colon-rectum [11]. By contrast, folic acid given in supplements has been shown to promote the progression of pre-malignant colorectal lesions [12]. This illustrates the increasing evidence that micronutrient supplements may not always afford the same protective effects attributed to dietary sources. This is an important point in the debate between whole diets and the use of supplements, and is discussed further in later chapters.

Although a traditional MedDiet is still widely consumed, especially by more elderly people, the diet is now increasingly under threat. Protecting healthy traditional diets against the encroaching uniformity of food, particularly the influence of fast food, is now recognised as a high priority [6]. Consequently, there was an initiative by Spain, Italy, Greece and Morocco to help protect the traditional MedDiet by applying for it to be adopted by UNESCO’s Intergovernmental Committee for the Safeguarding of the Intangible Cultural Heritage of Humanity [13]. The MedDiet achieved this recognition in November 2010. The box below gives the statement issued by UNESCO at the time of this recognition, and emphasises how much the MedDiet represents an overall lifestyle rather than just the consumption of food.

UNESCO DECLARATION ON THE MEDITERRANEAN DIET AS AN INTANGIBLE CULTURAL HERITAGE OF HUMANITY

<http://www.unesco.org/culture/ich/index.php?lg=en&pg=00011&RL=00394>

The Mediterranean diet constitutes a set of skills, knowledge, practices and traditions ranging from the landscape to the table, including the crops, harvesting, fishing, conservation, processing, preparation and, particularly, consumption

of food. The Mediterranean diet is characterized by a nutritional model that has remained constant over time and space, consisting mainly of olive oil, cereals, fresh or dried fruit and vegetables, a moderate amount of fish, dairy and meat, and many condiments and spices, all accompanied by wine or infusions, always respecting beliefs of each community. However, the Mediterranean diet (from the Greek *diaita*, or way of life) encompasses more than just food. It promotes social interaction, since communal meals are the cornerstone of social customs and festive events. It has given rise to a considerable body of knowledge, songs, maxims, tales and legends. The system is rooted in respect for the territory and biodiversity, and ensures the conservation and development of traditional activities and crafts linked to fishing and farming in the Mediterranean communities which Soria in Spain, Koroni in Greece, Cilento in Italy and Chefchaouen in Morocco are examples. Women play a particularly vital role in the transmission of expertise, as well as knowledge of rituals, traditional gestures and celebrations, and the safeguarding of techniques.

The current widespread interest in the MedDiet has necessitated the development of various definitions of what constitutes a ‘modern’ MedDiet. Such a definition is particularly important for epidemiologists in order to be able to assess the adherence of individuals to a MedDiet [14]. Many of these epidemiological studies have been conducted in European Mediterranean countries, and hence the definitions of the MedDiet tend to reflect the traditional MedDiet of these countries [10]. One widely-used definition of the relative consumption of nine key food groups is as follows:

1. high consumption of olive oil and low consumption of lipids of animal origin (resulting in a high ratio of monounsaturated to saturated fat)
2. high consumption of vegetables
3. high consumption of fruit
4. high consumption of legumes
5. high consumption of cereals (including bread)
6. moderate to high consumption of fish
7. low to moderate consumption of milk and dairy products (mainly cheese and yogurt from goats and sheep milk)
8. low consumption of meat and meat products
9. moderate consumption of wine

The use of current definitions of the MedDiet is discussed in Chapter 8, and the application of these assessments to disease prevention is considered in Chapters 10–13.

It is important to recognise that the MedDiet not only defines foods whose consumption *is* desirable, but should also encompass foods whose consumption is *not* desirable. This includes the absence of industrial processed foods, and is one reason for the relatively low levels of salt, saturated fat, trans fats and sugar in the MedDiet compared to the standard Western diet.

1.1.3 International differences

Despite some similarities between MedDiets, it is generally agreed that there is no one MedDiet. This is not surprising in view of the fact that Mediterranean countries are located in three different continents, namely Europe, Asia and Africa. These continents

have major cultural differences, not least of which is religion: European Mediterranean countries are Christian and those in Asia and Africa are Muslim. Hence wine consumption, a cornerstone of the European MedDiet, is absent in Muslim countries where alcohol consumption is prohibited. There are many other differences throughout the Mediterranean basin that can influence dietary habits, and these range from climate and geography, to socio-economic factors, culture and history. These can be regional as well as international. For example, consumption of fish within a country tends to vary depending on proximity to the sea, and, at the international level, low consumption in some countries is also due to the relative scarcity of fish in some parts of the Mediterranean. Even olive oil consumption – considered another cornerstone of the MedDiet – can vary widely between Mediterranean countries. Hence, the overall types of foods can vary quite widely between various Mediterranean regions as is illustrated in Table 1.3.

The geographical boundaries within which a ‘MedDiet’ is eaten are not precisely defined. Twenty-one countries border the Mediterranean sea (although this number varies according to the definition of a national state) (Figure 1.1). The climates of these countries can vary widely from region to region, and parts of many countries that border the Mediterranean sea do not have a ‘Mediterranean’ climate in its precise climatological definition. (This is defined by climatologists as the Cs climatic region, i.e. having warm to hot, dry summers and cool, wet winters.) Even bordering the Mediterranean sea itself does not guarantee a MedDiet since parts of the coasts of Libya and Egypt do not have a Mediterranean climate. Northern Italy is another region that does not have a Mediterranean climate and correspondingly the traditional diet here is quite different to that found in Mediterranean Southern Italy. In France, only the regions of Provence, Languedoc and part of Roussillon have a Mediterranean climate, and the cuisines of other regions are quite different. By contrast, many parts of Portugal do have a Mediterranean climate, although this country is on the Atlantic seaboard and has no border with the Mediterranean sea.

A poetic, yet very useful, definition of the Mediterranean is that of the French writer Georges Duhamel who wrote: ‘The Mediterranean ends where the olive tree no longer grows’. This can be used to delineate the northern limits of ‘Mediterranean’ cuisine and is shown in Figure 1.2.

The Mediterranean climate, as defined by climatologists, is not restricted to countries of the Mediterranean region but includes parts of California, the Western Cape in South Africa, central Chile, southern Western Australia and the coastal areas of central and south-east Australia (Figure 1.3). For cultural and historical reasons, these countries did not develop a traditional MedDiet. However, their climates have allowed the production of traditional Mediterranean foods, especially grapes, and, increasingly, olives, and levels of production of traditional Mediterranean foods are increasing in these countries.

1.1.4 National representations of the MedDiet

Efforts to promote the MedDiet have led to the development of various pictorial representations. These graphics aim to convey the essentials of the MedDiet in a single glance without recourse to lengthy text, as it is considered that this could be off-putting to some people. The design of the graphic is often angled in order to appeal to the populace of a particular country.

Perhaps the best known of these pictorial representations is the food pyramid developed by the American organisation Oldways (Figure 1.4). This structure is

Table 1.3 Important food groups in Mediterranean regions (from [15]). With permission from Elsevier.

Mediterranean region	Cereals	Dairy	Olive oil consumption	Meat	Other
<i>Western</i> (Spain, France, Italy, Malta)	Bread, rice, pasta	Cheeses	High in Italy and Spain	Pork	Potato
<i>Adriatic</i> (Croatia, Bosnia, Albania)	White wheat flour as bread and pitta	High (butter, buttermilk, ricotta, cheese, sour cream)	Low to moderate	Beef	
<i>Eastern</i> (Greece, Lebanon, Cyprus, Turkey, Egypt)	White flour products	Various cheeses	Very high in Greece, negligible in Egypt	Chicken*	Okra in summer, herbs (dill, parsley, oregano)
<i>North Africa</i> (Libya, Algeria, Morocco, Tunisia)	Bread made from whole meal flour and barley flour Couscous		Wide range	Lamb	Potato, pumpkin, chickpeas, dates, date molasses

*Lamb is also popular, especially in Greece.



Figure 1.1 Countries of the Mediterranean Basin (Wikipedia). The countries bordering the Mediterranean sea are (with semi-autonomous countries): Gibraltar, Spain, France, Monaco, Italy, Slovenia, Croatia, Bosnia & Herzegovina, Albania, Greece, Turkey, Cyprus, Syria, Israel, Lebanon, Egypt, Libya, Malta, Tunisia, Algeria, Morocco.




Figure 1.2 The Mediterranean region as defined by the northern limits of olive cultivation (black line) and climate (hashed areas) [1]. With permission from John Wiley & Sons.




Figure 1.3 Regions of the world with a Mediterranean climate (Wikipedia).

similar to the US Food Guide Pyramid developed by the USDA. The Oldways pyramid has been refined over several versions. The base of the pyramid depicts physical activity and enjoying food with others. The remainder of the pyramid depicts the relative proportions of various foodstuffs that should be consumed, with the largest portion of the pyramid being devoted to plant foods. In addition, this 2009 version includes herbs and spices for the first time, although wild greens are not included, perhaps because their collection is not a traditional part of food culture in industrialised countries such as the US. Fish and shellfish have a prominent




Figure 1.4 The Oldways' pyramide depiction of a MedDiet [16]. Reproduced with permission. © 2009 Oldways Preservation & Exchange Trust, www.oldwayspt.org

position, with lesser amounts of poultry and dairy produce. No distinction is made between dairy produce from cow milk (uncommon in the traditional MedDiet) and milk from goats and sheep, although cheese and yogurt are depicted and milk is excluded, and this is consistent with a traditional MedDiet. Meats and sweets are simply advised to be eaten 'less often'. Wine in moderation and water are also shown.

Nutritionists in Mediterranean countries may not necessarily consider (Egyptian) pyramids to be the most appealing depiction of the MedDiet for their populaces, so they have developed images more representative of their own countries. Greek nutritionists have come up with the idea of seven Greek columns, each column showing the food to be consumed – in words rather than images – on one day of the




Figure 1.5 The Food Standards Agency Eatwell plate (<http://www.eatwell.gov.uk/healthydiet/eatwellplate/>). © Crown copyright material 2007, with permission from the Controller of HMSO and Queen's Printer for Scotland.

week [17]. The basic principles of a healthy way of life, 'moderation, variety and proportionality', are also mentioned, together with having a correct energy balance. Italian nutritionists have developed the historical theme by turning the Greek column into a Greco-Roman temple, an apparent attempt to capitalise on the idea that a temple symbolises 'healthiness, spirituality and self-improvement' [18]. In this Italian model, the steps of the temple convey a healthy lifestyle, the need for adequate exercise, and to use virgin olive oil and to drink wine in moderation. The columns of the temple represent desirable food groups, and foods that should only be consumed in limited amounts are shown at the top of the temple. A practical weekly guide has also been developed for French consumers [19] and this is discussed fully in Chapter 14.

It is interesting to compare these representations of the MedDiet with a current pictorial representation of UK dietary guidelines as promoted by the UK Food Standards Agency (FSA) – the so-called 'Eatwell Plate' (Figure 1.5). Although there are some similarities, there are also significant differences. Fruits, vegetables and carbohydrates all feature prominently on the Eatwell Plate, and this is similar to the representations of the MedDiet. Dairy products (from cow milk) are a more prominent feature on the Eatwell Plate than on the MedDiet depictions, reflecting a long tradition of dairy farming and dairy consumption in the UK. No distinction is made on the Eatwell Plate between the relative proportions of meat and fish to be consumed. Cakes and biscuits represent a relatively large slice of the Eatwell plate. Although these are also depicted on some of the MedDiet graphics, in the MedDiet these products are likely to be based on fruit, honey, nuts and olive oil, whereas in

the UK there will be a far higher content of refined sugar and saturated fat. Olive oil and wine are absent from the Eatwell Plate.

1.1.5 Sources of information for the general public

Mediterranean restaurants and cookery books

Tourism has had a major impact on the appreciation by North Europeans of Mediterranean cuisine, not to mention Mediterranean wines. There are now a plethora of Mediterranean restaurants in most towns. Elizabeth David pioneered home cooking of Mediterranean food in the UK with the publication of her book *A Book of Mediterranean Food* in 1950. It remains in print to this day. Many other Mediterranean cookery books are now available and some, such as those by Claudia Roden on Middle Eastern cookery and those by Clifford A. Wright, are scholarly works and include authentic recipes. There is, however, very little information on whether or not cookery books promote a healthier lifestyle. In the case of Mediterranean cookery books, most do not give a clear indication of the relative proportions of different foods that should be eaten – a fundamental aspect of healthy eating. Some books emphasise recipes using ingredients readily available to the non-Mediterranean cook rather than being truly representative of local Mediterranean cooking. In particular, MedDiet cookery books often include a relatively high proportion of recipes that contain meat, and this is not representative of a traditional MedDiet.

MedDiet organisations

There are several organisations that promote the MedDiet, and these have useful websites. The Mediterranean Diet Foundation (La Fundación Dieta Mediterránea, FDM) is a non-profit organisation based in Barcelona, Spain which promotes the investigation and dissemination of the MedDiet and the Mediterranean lifestyle. It organises conferences, runs courses, and organises a range of workshops with a particular focus on children and the elderly. Its website is at <http://www.fdmmed.org>. Oldways is an American non-profit organisation that was responsible for developing the best-known of the Mediterranean diet food pyramids (see above). It promotes the MedDiet through conferences, events and has introduced the ‘Med Mark’ in the US to help guide consumers in choosing traditional Mediterranean foods, drinks and other products. They have a website at <http://www.oldwayspt.org>.

1.2 Lifestyle factors

“We do not sit at table only to eat, but to eat together.”

(Plutarch)

The traditional MedDiet, and its health benefits, cannot be fully understood without considering its cultural context. The ‘Mediterranean food culture’ (or ‘Mediterranean lifestyle’ as it is sometimes known) includes factors such as fixed meal times, eating as part of a social gathering and – in some countries – taking a siesta after the midday meal. Other lifestyle factors such as physical activity, not smoking, and low

Table 1.4 Breakfast ingredients in Mediterranean countries (adapted from [23]). With permission from Elsevier.

Country	Typical breakfast ingredients
Greece (rural)	bread, cheese, coffee, goat's milk, fruits, olives
Greece (urban)	coffee, milk, pastries, fruit juices, preserved fruits, eggs with ham and sausages, cheese, butter
Spain	coffee, pastries (<i>churros</i>)
Egypt (rural)	coffee or tea, bread, onion, salt
Egypt (Cairo)	coffee or tea, goat cheese, bread, broad beans (in a stew accompanied with eggs or smoked meat)
Italy	cappuccino, wheat biscuits, fruit
Turkey	Turkish coffee or tea, cheese, raw vegetables, olives

levels of stress have also been found to contribute to the overall health benefits of a MedDiet [20–22]. If one considers ‘diet’ in terms of its original meaning of ‘a mode of living’ from the Greek *diata*, then the term ‘Mediterranean diet’ is completely congruous when implying the overall Mediterranean lifestyle.

The central part played by food in daily Mediterranean life is reflected in the care with which local, seasonal ingredients are sourced from respected producers and the care given to preparation and cooking. Producers themselves often sell their produce at local Mediterranean markets, and this promotes a close relationship between the consumer and the producer. This relationship helps maintain the quality of the produce and ensures that good agricultural practices are used, practices that can have a major impact on the nutritional value of the food (see Chapter 3).

1.2.1 Meal patterns

Meals are still an integral part of daily society in most Mediterranean countries, and the main meal is an important opportunity for bringing family members together. Lunch is still the main meal for many Mediterranean people, and many go home for lunch whenever possible, especially in Southern and Eastern countries [23]. However, the evening meal is increasingly becoming the main meal in regions where people work through the day or who work too far away from home to return there for lunch [23]. Breakfasts tend to be light compared to North European countries (Table 1.4), both in terms of quantity and in the low consumption of produce with high levels of saturated fats such as butter, sausages and bacon that typify the traditional English breakfast.

There is evidence that eating at fixed times during the day, and the associated periods when meals are not consumed, is important for inducing satiety which discourages excessive calorie intake [24]. This does not exclude the occasional consumption of snacks in traditional MedDiets (see Table 1.5), but whereas snacks in Northern Europe are a major source of salt, sugars, and saturated and trans fats [25], snacks in Mediterranean countries mostly consist of either fruit, nuts or home-made delicacies.

Table 1.5 shows a weekly food pattern collated from Cretans eating a traditional diet. During the week, the meal that included the main daily source of protein was mostly lunch, and this may particularly benefit physically active people, since studies have shown that optimal muscle building occurs when protein is eaten immediately after exercise [26]. By contrast, dinners were mainly based on easily digested cooked vegetables.

Table 1.5 Typical foods in a traditional Cretan Mediterranean diet consumed over a week [27]. With permission from Elsevier.

	Breakfast	Mid-morning	Lunch	Mid-afternoon	Dinner
Monday	ksinohontros ¹ , rusk, orange	pear	broad beans, onion, salad (cucumber, tomato, purslane, olives, olive oil), whole-wheat bread, apple, red wine	walnuts, dry figs	boiled vegetables, potatoes, olive oil, boiled egg, melon, red wine
Tuesday	rusk, cheese, apple	orange	snails, potatoes and vegetables, salad (tomato, cucumber, onion, olive oil), whole-wheat bread, red wine, longan	halva ² (home-made)	rice with spinach, yogurt, whole-wheat bread, longan
Wednesday	doughnuts (homemade) with honey, apple, herbal tea	pear	chickpeas, herring, salad (tomato, cucumber, onion, olive oil), whole wheat bread, red wine	walnuts, figs, <i>raki</i>	stuffed tomatoes, whole wheat bread, salad (tomato, cucumber, onion), melon
Thursday	fresh whole milk boiled with ground wheat	melon	fish, broad beans (puree), oil, lemon juice, whole-wheat rusk, salad (tomato, cucumber, onion, olives, olive oil), pear, red wine	halva (home-made)	lentils, salad (tomato, cucumber, onion, olives, olive oil), apple, red wine, cheese, whole wheat bread
Friday	rusk, olives, herbal tea, apple	apple	beans, potatoes, whole-wheat bread, olives, orange	walnuts, dry figs, <i>raki</i>	broad beans, artichoke, olive oil, rusk, red wine, melon
Saturday	milk and whole wheat, melon	apple	chicken, okra, potatoes, salad (lettuce, cucumber, olives, olive oil)	home-made cheese pie, honey, coffee	boiled vegetables with olive oil, rusk, red wine, melon
Sunday	homemade cheese pie with honey, melon		rabbit, pasta, salad (tomato, cucumber, onion, olives, olive oil), rusk, wine, orange	coffee, halva	fish, fish soup with vegetables, rusk, red wine, apple

¹Ksinohontros = yogurt, wheat; ²halva = semolina, olive oil, sugar, walnuts.

1.2.2 Siestas

A siesta after a midday meal is still common in many Mediterranean countries. In Spain, most siestas were found to last for less than an hour. In a large study of healthy Greek men and women, working men who took a siesta were found to have fewer coronary deaths than those who did not nap [28].¹ The authors of this study suggested that regular siestas acted as a stress-reducing habit, which lowered the risk for CVD [28]. Not all studies, however, have found an inverse association between siestas and coronary deaths; indeed some studies have found a positive association. Taking a siesta is not conducted in a vacuum, and interpreting the data from some of these studies is subject to possible confounding factors. For example, people taking a siesta may take less physical exercise, which is a protective factor against CVD, and hence could give rise to a positive association between taking a siesta and CVD. Also, some individuals sleep during the day due to nightly sleep disturbances, which could be associated with underlying health problems. The Greek study did attempt to control for these possible confounding factors, although more studies examining the possible benefits of a siesta would be useful.

1.2.3 Physical activity

The high level of physical activity of the traditional Mediterranean peasant contrasts with the far more sedentary lifestyle of most present-day Mediterranean people. Despite this modern trend, the clement Mediterranean climate does still favour an outdoor lifestyle and more physical activity than is the case for many people living in more northern climates. In the HALE project, which evaluated the effects of a Mediterranean diet and lifestyle factors on mortality in elderly European men and women, physical activity was associated with a lower risk of all-cause mortality [22]. Increasing evidence is demonstrating that the health benefits of diet interact with other lifestyle factors, and physical exercise is now widely accepted to be an important factor that reduces the risk of age-related diseases such as CVD and some cancers [29]. A detailed study of Spanish children found that physical fitness is very important to reduce the risk of CVD and other diseases in later life, a point emphasised in the title of the paper: ‘A Mediterranean diet is not enough for health’ [30].

1.2.4 Sunshine

That Mediterranean countries are sunny is accepted almost without thought. But it has been argued that by inducing endogenous synthesis of vitamin D, sunlight is an important contributor to the health of Mediterranean people [31]. This proposal is based on recent evidence that the role of vitamin D in the body extends well beyond its role in bone health by preventing rickets and osteoporosis, and may also include a reduction in the risk for some cancers, hypertension and some immunological disorders (see Chapter 2).

1.3 Health benefits

There is a substantial body of epidemiological evidence for the health benefits of the MedDiet, and this is discussed fully in Chapters 10–13. A useful meta-analysis of a number of epidemiological studies generated the following estimates for the protective value of the MedDiet [32]:

¹ This analysis was not possible for women due to their lower incidence of coronary deaths.