







































# List of Figures

Fig. 1.1	The brown bear, which is also called the grizzly bear, is a large bear that ranges across most of North America (west of the Mississippi River) and in northern Europe and Asia. Infanticide, or the killing of young, has been widely studied in humans, but it also occurs in brown bear and other animals ranging from rotifers to other mammals .....	2
Fig. 1.2	An American psychologist, B.F. Skinner, invented an operant conditioning chamber, which measured the response of organisms (typically, rats) to a lever that provided food to the organism .....	3
Fig. 2.1	The woodchuck, or groundhog, is the only marmot in eastern North America. Unlike its western counterparts, the woodchuck is solitary; western marmots tend to be colonial.....	7
Fig. 2.2	The sandwich tern is a seabird, in which males plunge for fish and offer the fish caught to females as part of the courtship. Young sandwich terns learn food items from both parents via observational learning .....	12
Fig. 3.1	The yellow warbler is very wide-spread, extending its range throughout North America and as far south as central Mexico .....	16
Fig. 3.2	Fur seals consist of several species, with the northern fur seals occurring in the North Pacific and most other species occurring south of the equator. Typically, fur seals form assemblages each year during summer months to give birth and breed. All species are polygynous (a dominant male reproducing with more than one female) .....	18

Fig. 4.1 The spotted hyena, or laughing hyena, occurs today only of the south of the Sahara, except in the basin of the Congo. Hyenas live in large matriarch groups, or clan, which may comprise 80 individuals. Hyenas are not scavengers, but actively hunt large mammals..... 26

Fig. 4.2 The shiny cowbird of South America, although it may breed in southern Florida. Like other cowbirds, the shiny cowbird is a brood parasite..... 29

Fig. 5.1 The gray wolf, or timber wolf, is very social, living in packs, which often are closely related. The gray wolf shares a common ancestry with the domestic dog ..... 36

Fig. 5.2 The gypsy moth is a forest pest, from Europe and Asia, having escaped into the USA about 1860. Egg masses of gypsy moths are typically placed on branches and trunks of trees. Egg masses are buff colored when first laid but may become bleached over winter..... 37

Fig. 6.1 The great horned owl has females that are larger than males. Its “horns” are not ears or horns, but tufts of feathers. It is believed to occur from Nova Scotia to east Texas and to Minnesota and southward to South America. Great horned owls breed early in late January or early February ..... 44

Fig. 6.2 The wood stork, or sometimes called the wood ibis, is the only stork that breeds in North America. It extends its range throughout South America, in tropical and subtropical regions ..... 46

Fig. 7.1 The monarch is perhaps the best known of the butterflies in North America; it is often called the milkweed butterfly. Besides occurring in North America, monarchs are found in Europe and New Zealand. It is famous for its migration in the Americas, especially to Mexico, which may span 3–4 generations ..... 56

Fig. 7.2 The fox squirrel is closely related to gray squirrels, but fox squirrels are considered the most polymorphic mammal. In the southeastern USA, for example, the color of fox squirrels has different percentages of gray and black on dorsal surfaces..... 57

Fig. 8.1 The spotted salamander of eastern North America. It is considered as a mole salamander..... 71

Fig. 8.2 The northern river otter, or North American river otter, is found only in (endemic) North America. It is a weasel, which is well-adapted to aquatic habitats ..... 73