

Coastal Research Library 11

Gautam Kumar Das

Estuarine Morphodynamics of the Sunderbans

 Springer

Coastal Research Library

Volume 11

Series Editor

Charles W. Finkl
Department of Geosciences
Florida Atlantic University
Boca Raton, FL 33431
USA

The aim of this book series is to disseminate information to the coastal research community. The Series covers all aspects of coastal research including but not limited to relevant aspects of geological sciences, biology (incl. ecology and coastal marine ecosystems), geomorphology (physical geography), climate, littoral oceanography, coastal hydraulics, environmental (resource) management, engineering, and remote sensing. Policy, coastal law, and relevant issues such as conflict resolution and risk management would also be covered by the Series. The scope of the Series is broad and with a unique crossdisciplinary nature. The Series would tend to focus on topics that are of current interest and which carry some import as opposed to traditional titles that are esoteric and non-controversial. Monographs as well as contributed volumes are welcomed.

More information about this series at <http://www.springer.com/series/8795>

Gautam Kumar Das

Estuarine Morphodynamics of the Sunderbans

 Springer

Gautam Kumar Das
Department of Chemical Engineering
Jadavpur University
Kolkata, India

ISSN 2211-0577

ISBN 978-3-319-11342-5

DOI 10.1007/978-3-319-11343-2

Springer Cham Heidelberg New York Dordrecht London

ISSN 2211-0585 (electronic)

ISBN 978-3-319-11343-2 (eBook)

Library of Congress Control Number: 2014954024

© Springer International Publishing Switzerland 2015

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media (www.springer.com)

*... All things rush on, they stop not, they
look not behind, no power can hold them
back, they rush on.*

Rabindranath Tagore (1912)
Gitanjali (Hymn No. LXX)

Dedicated to the People of Sunderbans

Foreword

The Sunderbans, a UNESCO World Heritage Site covering parts of Bangladesh and the southern tip of Indian state of West Bengal, is a part of the world's largest deltaic plain of fluvio-marine deposit formed by the Ganges and Brahmaputra at the confluence of the Bay of Bengal. It is the largest single block of tidal halophytic mangrove forest in the world, conspicuous for its great size and biodiversity. With an anastomosing network of channels and creeks, and tidal inundation twice daily, Sunderbans mangroves wetland is a dynamic and complex ecosystem, which undergoes continuous processes of erosion and accretion. Natural processes like changes in local hydrology, sediment motion under wind, wave and tidal action, beach dynamics, regional and global processes like sea level rise as well as the impact of human interference in the form of reclamation of forest land, changes in land use patterns, coastal urbanizations etc. are the lead factors for the changes in the environmental scenario of Sunderbans. Only a century ago, Indian Sunderbans with a total area of 9,630 km² was covered with lush green mangrove forests, which at present is left with only 4,266.6 km² area with mangroves and the rest has been converted to various land use patterns.

Global warming has left its imprint in the form of sea level rise in the Sunderbans. Sagar Island, the biggest deltaic island of the Sunderbans, has recorded 3.1 mm/year of sea level rise (IPCC 2007). As a result of this there has been flooding of low-lying deltas, change of shoreline, and salinisations have been decreasing gradually due to mixing of seawater with melt water of ice. Again, the incursion of flooded coastal saline water into the freshwater region increases the salinity of the latter. Silt particles get mixed with water and the resultant heat of mixing as a result of the kinetic energy from those vortexes may cause the water temperature to be warmer than that of the ambient temperature during winter.

Evidences of erosion followed by differential subsidence and sediment filling in several parts of the Sunderbans can be sited from the terraced estuarine bank pattern, undercutting and collapsing of river banks etc. The estimated rate of erosion, as evidenced from the relics of the sea wall exposed in the intertidal beach zone at Bakkhali and Fraserganj in coastal Sunderbans, is 9.8 m/year during the period

1930–1970 and 8.6 m/year during 1971–1995 which is very high. The northeastern, southeastern and southwestern sides of Sagar Island are facing vigorous erosion due to the concerted acts of various natural processes and anthropogenic activities. The erosion rate from 1996 to 1999 was calculated to be 5.47 m/year. The part of the destabilized coastal dunes have been advancing inland at the rate of 17 m/year. Beach erosion has also been computed from the presence of gastropod species *Amalda ampla* which is considered as a biological indicator.

The highly specialized mangrove ecosystem of Indian Sunderbans basically can be treated as an estuarine ecosystem supporting a large number of aquatic organisms living either entire life cycles within the mangroves or visiting the mangrove swamps and waters for food or to breed. The occurrence of both the flora and fauna in this fragile ecosystem has become vulnerable as a result of changing morphodynamics of this estuarine environment. Expansion of mangrove swamp to the salt marsh is the clear indicator of changing trends of the mangrove habitat. Mangrove extension into the marshy areas and the gradual disappearance of islands emerged in the coastal areas because of the increased tidal amplitudes for climatic changes and sea level rise are highly alarming for the Sunderbans.

A large number of endangered species of fin fishes and shell fishes are suffering a severe depletion in population because of random exploitation in addition to the damage caused for the capture of prawn seeds. Continuous prawn seed collection can decrease the density of pneumatophores and the biomass of epiphytic algae as trampling is a perennial problem which results the habitat structure of mangrove plants as well as aquatic organisms. This, in turn, adds to a tremendous threat to the balance of the coastal ecosystem.

Estuarine morphodynamics are further influenced by the changes in river course, impeded fresh water flow, climate variability and sea level changes. All these deleterious operations are currently continuing in most areas of Sunderbans, which, in turn, during the recent decades, have been bringing about a serious change in the geomorphology and hydrodynamics of the water courses together with an obvious change in the nature of nutrient recycling. As a result of these morphodynamic processes and anthropogenic interferences, exploitations of mangrove vegetations have led to the large scale degradation of mangrove areas. Further, mangroves have also been lost through natural causes including erosion and severe siltation as a result of sedimentation. At present, tiger straying is a common phenomenon. The present study encompasses the areas of the trend of salinity decline, increase of temperature of river waters, non-uniformity of bottom topography, erosion, change of the tidal courses and shapes of islands, increasing rate of sedimentation in the river bed and mid channel bar formations; collapsing of bank materials; and random collection of prawn seeds that results in the ecological imbalance in this unique ecosystem. It is expected that the findings of this study would be a potential contribution to enlist the morphodynamic factors responsible for the changing environmental scenario and to formulate a comprehensive and sustainable management plan for the long-term conservation and protection of the mangrove ecosystem in the Indian Sunderbans.

Acknowledgement

I would like to extend my sincere thanks to Sri S. Rakshit, Former Director, Geological Survey of India; Prof. Asokkumar Bhattacharya, Former Head, Department of Marine Science, University of Calcutta; Prof. A. Mazumder, Director and Head, Department of Water Resource Engineering, Jadavpur University; Prof. S. R. Basu, Former Head, Department of Geography, Calcutta University; Prof. S. Dutta, Pro-VC, Jadavpur University; and Prof. H. R. Betal, Former Head, Department of Geography, Calcutta University, for their constant inspiration, guidance and advice.

The author gratefully acknowledges various helps including laboratory facilities provided by the Director, Marine Wing, Geological Survey of India and Head, Department of Chemical Engineering, Jadavpur University.

I also gratefully acknowledge the helps rendered by the Directors of Zoological Survey of India, Sunderbans Biosphere Reserves and Indian Meteorological Department. Most cordially I acknowledge the co-operation and assistance extended by the various departments of the Government of India and West Bengal, Forest Directorate, Sunderbans Tiger Reserves and Sunderbans Development Board, Government of West Bengal.

I gratefully remember the generous helps and hospitality rendered to me during field work and on-board field surveys in the forest of Sunderbans by many forest rangers, guards, fishermen, honey collectors, tour operators and many islanders, without which it would have been impossible to conduct this work. Rokeya and Titas helped in drawings and corrected the proofs.

Contents

1	Sunderbans: Physical Aspects and Configurations	1
1.1	Physical Aspects	2
1.2	Location and General Boundaries	5
1.3	Reclamation of Sunderbans	5
1.4	Zonation of Sunderbans	5
1.5	Climate	6
	1.5.1 Cyclones	7
	1.5.2 Tides	8
1.6	Waves	8
1.7	Physiography and Configuration	9
1.8	Biogenous Coast	10
1.9	Pattern of Sedimentation	10
1.10	Stratigraphy	11
1.11	Geotectonic Settings	11
	1.11.1 Bengal Basin	13
	1.11.2 Bengal Delta	14
	1.11.3 Deltaic Flood Plain	14
	1.11.4 Lower Delta Plain	15
1.12	Geomorphic Environments	15
	1.12.1 Coastal Zone	16
	1.12.2 Sandy Beaches	16
	1.12.3 Islands	17
1.13	Sunderbans: Its Uniqueness	17
1.14	Summary	19
	References	19
2	River Systems and Tidal Courses	23
2.1	River Networks	24
	2.1.1 Ichamati	25
	2.1.2 Kalindi- Raimangal -Haribhanga	27
	2.1.3 Bidyadhari	27

2.1.4	Piyali	28
2.1.5	Matla	29
2.1.6	Bidya	29
2.1.7	Guasuba	30
2.1.8	Bangaduni	30
2.1.9	Thakuran	30
2.1.10	Mridangabhanga	31
2.1.11	Gobadia	32
2.1.12	Saptamukhi	32
2.2	Morphodynamics of the Tidal Rivers	32
2.3	Geomorphic Sandbodies	33
2.4	Bedforms: Indicators of Hydrodynamics	33
2.5	Riverine Geomorphic Environments	34
2.5.1	River Banks (Natural Levees)	35
2.5.2	Point Bar	36
2.5.3	Mid-channel Bar	37
2.5.4	Swash Platform	38
2.5.5	River Mouth Bar	38
2.6	Tidal Creek Systems	39
2.6.1	Mud Ridges	40
2.6.2	Mud Microdelta	41
2.6.3	Mud Pellets and Mud Lumps	41
2.6.4	Wood Clumps on Mudflats	41
2.7	Embankments	41
2.8	Summary	43
	References	44
3	Brackish Water	47
3.1	Physico-chemical Properties	48
3.1.1	pH	48
3.1.2	Temperature	49
3.1.3	Dissolved Oxygen (DO)	51
3.1.4	Salinity	51
3.1.5	Alkalinity	55
3.1.6	Turbidity	55
3.1.7	Carbon Dioxide	56
3.2	Dissolved Nutrients	56
3.3	Water Quality of Coastal Areas of Sunderbans	57
3.4	Summary	57
	References	58
4	Substrate Soils	61
4.1	Soil Formation	62
4.2	Soil Characteristics	63
4.3	Soil Classifications	63
4.4	Saline Soils	64

4.5	Characteristics of the Saline Soils	64
4.5.1	Saline Alkali Soils	64
4.5.2	Non-saline Alkali Soils	65
4.5.3	Degraded Alkali Soils	65
4.5.4	Acid Sulphate Soils	65
4.6	Physico-chemical Properties of Soil	66
4.6.1	Soil Reaction (pH)	66
4.6.2	Organic Carbon	68
4.7	Soil Nutrients	69
4.7.1	Nitrogen	69
4.7.2	Phosphorus	69
4.7.3	Potassium	70
4.7.4	NPK of Sunderbans	70
4.8	Micronutrients	70
4.9	Metal Content and Metal Oxides of Soils	71
4.10	Soil Distribution	73
4.10.1	Saline Soils	73
4.10.2	Saline: Alkaline Soils	73
4.10.3	Non-saline Alkali Soils	74
4.10.4	Degraded Saline Alkali Soils	74
4.11	Summary	74
	References	75
5	Coastal Morpho-Dynamics	77
5.1	Coastal Environment	77
5.2	Sediment Transport and Shoreline Progradation	78
5.3	Bakkhali and Fraserganj Coastal Environments	78
5.3.1	Beach Morphology	79
5.3.2	Climate	80
5.3.3	Tide	81
5.4	Dunes	81
5.4.1	Roles of Coastal Dunes	82
5.5	Coastal Processes	83
5.6	Tidal Shoals	84
5.6.1	Bedforms	85
5.7	Sediment Texture	85
5.8	Composition of Beach Sediments	85
5.9	Micro-paleontology	85
5.10	Shoreline Recession	86
5.11	Mud Balls	87
5.11.1	Significance of Mud Balls	88
5.12	Trash in the Beach	88
5.13	Coastal Waters	88
5.14	Sedimentation at Ganga Sagar Beach and Chuksar Island	89
5.15	Bedforms of Upper Foreshore Zone	91

5.15.1	Backwash Ripple Marks	91
5.15.2	Rhomboid Ripple Marks	91
5.16	Bedforms of Mid Foreshore Zone	92
5.16.1	Mega Ripple Marks	92
5.16.2	Lingoid Ripple Marks	92
5.16.3	Interference Ripple Marks	93
5.17	Bedforms of Lower Foreshore Zone	94
5.17.1	Double Crested Ripples	94
5.17.2	Straight Crested Small Ripples Marks	95
5.17.3	Capped – Off Ripples	95
5.17.4	Ripple – Rill Complex	95
5.18	Texture of Beach Sediments	95
5.19	Beach Character and General Flow Phenomena	96
5.20	Summary	98
	References	99
6	Granulometry of Mangrove Sediments	103
6.1	Measures of the Grain Size Distribution	104
6.2	Statistical Analysis of Grain Size	104
6.3	Sampling of Sediments	105
6.4	Cumulative Curves	105
6.5	Expression of Sediment Grain Size	105
6.6	Grain Size Characteristics of Different Estuarine Landforms	106
6.7	Grain Size Populations	106
6.8	Texture of Dune Sands	107
6.9	Texture of Beach Sediments	108
6.10	Texture of Sand Flats	109
6.11	Texture of Tidal Shoal Sediments	110
6.12	Texture of Point Bar Sediments	111
6.13	Texture of Mudflat Sediments	112
6.14	Texture of Marsh Sediments	113
6.15	Texture of Mangrove Swamp Sediments	115
6.16	Texture of River Sediments	115
6.17	Textural Sensitiveness	116
6.18	Summary	116
	References	117
7	Estuarine and Coastal Erosions	119
7.1	Controlling Factors for Erosions	120
7.2	Flow Pattern	120
7.3	Geomorphic Sites of Erosion	122
7.4	Characteristics of Estuarine Banks	122
7.5	Characteristics of River Banks	123
7.6	Erosional Processes and Features	123
7.6.1	Undercutting and Collapsing	124
7.6.2	Subsidence and Rapid Sedimentation	125

7.6.3	Piecemeal Mechanism	125
7.6.4	Furrow and Ridge Structures	126
7.6.5	Mud – Mounds	127
7.6.6	Mud Balls	127
7.7	Erosion at Estuarine Mouths	128
7.8	Coastal Erosion	128
7.9	Mechanisms of Coastal Erosion	129
7.10	Biological Evidences of Erosion	129
7.11	<i>Amalda ampla</i> – A Biological Indicator of Erosion	131
7.11.1	Characteristic Features of <i>Amalda ampla</i>	131
7.11.2	Physico – chemical Parameters	132
7.11.3	Sediment Texture	132
7.11.4	Habitat Structure of <i>Amalda ampla</i>	132
7.12	Estimation of Beach Erosion Using Biological Tools	133
7.12.1	Physico-chemical Environment of Sediments	133
7.12.2	Observations	134
7.12.3	Computation	134
7.13	Summary	135
	References	137
8	Floral Diversity	139
8.1	Mangroves	141
8.1.1	Geological History of Mangroves	141
8.2	Forest Management	141
8.3	Mangrove Characteristics	142
8.4	Geomorphic Set Up of Mangrove Swamps	142
8.5	Floral Dynamics	143
8.6	Forest Types	143
8.7	Mangrove Varieties	145
8.8	Mangrove Zonation	146
8.9	Habitat of Occurrence	148
8.10	Chemistry of Mangrove Sediments	148
8.11	Composition of Mangrove Sediments	148
8.12	Salt Marshes	149
8.12.1	Texture and Chemistry of Marsh Sediments	150
8.12.2	Marsh as Sediment Binder	150
8.12.3	Roles of Salt Marsh	150
8.13	Mangrove Afforestation and Deforestation	151
8.13.1	Regeneration	151
8.13.2	Afforestation	151
8.13.3	Deforestation	153
8.14	Summary	155
	References	155

9	Mangrove Ecosystems	159
9.1	Mangrove Community	160
9.2	Mangrove Fauna and Food Web	160
9.3	Primary Food Source	161
9.4	Components of the Mangrove Ecosystems	162
9.5	Detritus Food Web	164
9.6	Mangrove Litter	164
	9.6.1 Break Down of Mangrove Litters	167
	9.6.2 Decomposition Process	167
	9.6.3 Microbial Utilizations	169
9.7	Nutrient Recycling	169
9.8	Nutrient Inputs and Exports	170
9.9	Energy Flow	170
9.10	Functions of Mangrove Ecosystem	171
9.11	Mangrove and Man	174
9.12	Summary	174
	References	175
10	Anthropogenic Interferences	177
10.1	Prawn Captures	178
10.2	Environmental Requirements for Habitat	179
	10.2.1 Sediment Composition	179
	10.2.2 Physico-chemical Control for Sustainability of Prawn Seeds	179
10.3	Geo-hydrological Control Related to Capture	180
10.4	Seed Availability	180
10.5	Seed Composition	181
10.6	Drifting Pattern	182
10.7	Different Phases of Prawn Seeds (<i>P. monodon</i>)	182
	10.7.1 Metamorphosis of Prawn	183
10.8	Collection Method	183
10.9	Encountered Species	183
10.10	Transport of Seeds	184
10.11	Observations	185
10.12	Management Plan	186
10.13	Impact of Seed Capture	187
10.14	Water Quality of Bidyadhari River	188
10.15	The River System	189
	10.15.1 Decline of the Bidyadhari River	189
	10.15.2 Present State of the Bidyadhari River	190
10.16	Environmental Parameters of River Waters	191
10.17	Man-Tiger Conflict and Tiger Straying	194
	10.17.1 Classification of Man Eaters	194
	10.17.2 Occurrences for Man Eating	195
	10.17.3 Measures for Minimization of Confrontation	195

10.18	Trash in the Coastal Sunderbans	196
10.19	Summary	196
	References.....	198
11	Changing Environmental Scenario	199
	References.....	203
Index	205

Abbreviations and Units

BOD	Biochemical oxygen demand
COD	Chemical oxygen demand
DO	Dissolved oxygen
ECw	Electrical conductivity of water
ECs	Electrical conductivity of soil
IFAD	International Fund for Agricultural Development
IMD	Indian Meteorological Department
IUCN	International Union for the Conservation of Nature
LOI	Loss on ignition
MAB	Man and biosphere
ROOM	Readily oxidisable organic matter
SBR	Sunderbans Biosphere Reserve
STR	Sunderbans Tiger Reserve
TDS	Total dissolved solids
TSS	Total suspended solids
UNDP	United Nations Development Programme
UNEP	United Nations Environmental Programme
ppt	Parts per thousand (‰)
ppm	Parts per million
dS/m	desi Siemens per meter
1 ppt	2161 dS/m
1 dS/m	1000 EC = 1000 ppm
100 hectares	1 km ²

