

Gianfranco Butera · Massimo Chessa
Andreas Eicken · John Thomson *Editors*



Cardiac Catheterization for Congenital Heart Disease

From Fetal Life to
Adulthood

Forewords by

Shakeel Qureshi and Mario Carminati

 Springer

Cardiac Catheterization for Congenital Heart Disease

Gianfranco Butera • Massimo Chessa
Andreas Eicken • John Thomson
Editors

Cardiac Catheterization for Congenital Heart Disease

From Fetal Life to Adulthood

Forewords by Shakeel Qureshi
and Mario Carminati



Springer

Editors

Gianfranco Butera
Pediatric and Adult Congenital Heart
Center
IRCCS Policlinico San Donato
Milano
Italy

Andreas Eicken
Department of Pediatric Cardiology
and Congenital Heart Disease
German Heart Center, Technical
University of Munich
Munich
Germany

Massimo Chessa
Pediatric and Adult Congenital Heart
Center
IRCCS Policlinico San Donato
Milano
Italy

John Thomson
Pediatric Cardiology
Yorkshire Heart Centre
Leeds
UK

ISBN 978-88-470-5680-0

ISBN 978-88-470-5681-7 (eBook)

DOI 10.1007/978-88-470-5681-7

Springer Milan Heidelberg New York Dordrecht London

Library of Congress Control Number: 2014948068

© Springer-Verlag Italia 2015

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Printed on acid-free paper

Springer is part of Springer Science+Business Media (www.springer.com)

Foreword

There has been a revolution in cardiac catheterization and interventional techniques in congenital heart disease in the last three decades and, in particular, in the 1990s and the early part of this century. There is always concern amongst clinicians that the pace of developments is likely to slow down. However, fortunately, there does not seem to be any slowing in the progress of developments of newer interventional techniques. Therefore, there has been a need for a book dealing with practical aspects of cardiac catheterization in congenital heart disease. Such books are not only aimed for trainees and interventionists in early parts of their careers, but they are also invaluable to experienced interventionists, helping them to keep abreast of developments. The Editors of this handbook on *Cardiac catheterization for Congenital Heart Disease: From Fetal life to Adulthood* are to be commended for their efforts and for addressing this important task.

This book, composed of 47 chapters, will attract a large audience from all over the world, thanks to the involvement of some of the leading authorities in our specialty. The interventional techniques from the fetus to the adult with congenital heart

disease are covered. All the technical chapters follow a similar format, dealing with important practical steps that each interventionist needs to know. These include the indications for interventions, any pre-procedure assessment, step-by-step technical information, as well as practical tips and tricks. Issues such as informed consent represent important steps in the performance of interventional procedures, and these aspects are not forgotten in this book. There are new interventional balloons and devices appearing regularly, and attempts are made to provide up-to-date information on these. Interventionists need detailed step-by-step information about access into the heart, and so usual and unusual accesses are dealt with. Inevitably with the interventional techniques there will be a spectrum of complications, and so these and many retrieval techniques are addressed.

The editors have produced a much-needed book at the correct time. It will act as a quick practical guide to all the interventionists. I have no doubt whatsoever that it will become a crucial requirement as a stock item for all the departments and libraries.

Evelina Children's Hospital,
London, UK

Shakeel A. Qureshi

Foreword

It is my great pleasure to write a foreword for this book on transcatheter interventions of congenital heart defects. The volume follows “how-to” format, as a result of a worldwide cooperation of many international experts, who put together their knowledge and expertise.

As the chief of Pediatric and Adult Congenital Cardiology Department at Policlinico San Donato IRCCS, I have been involved for many years in the fascinating world of catheter interventions in congenital heart defects. During this period I had the pleasure to be the mentor of Gianfranco Butera and Massimo Chessa, who shared with me the enthusiasm in taking part of continuous and rapidly evolving process of a variety of new techniques in this field. They should be congratulated, together with Andreas Eicken and John Thomson, as Editors of this book, for the excellent work they did. This book can really become a day by day companion of

all young fellows and trainees around the world. I am convinced that also seniors and already expert interventionalists will find it useful in the daily practice.

Pediatric Cardiology Unit,
IRCCS Policlinico San Donato,
Milan, Italy

Mario Carminati

Preface

The population of patients recognized with congenital heart disease has dramatically increased over the past years as a consequence of significant advances in diagnosis and the improvement in surgical and transcatheter techniques.

Actually, the number of adults with congenital heart disease in the United States is almost the same as that of children with these anomalies. Additionally, the evolutions in both device technology as well as non-invasive imaging technology have allowed the safe and effective catheter-based treatment of various congenital or post-surgical residual cardiac defects. Furthermore, these patients have a quicker recovery and a shorter hospital stay, and a better quality of life.

On the other hand, these techniques require not only a specific training, but also the necessity of a tight collaboration between pediatric and adult interventionist as well as with a trained echo-cardiographer and anesthesiologist.

Actually, interventional cardiology of subjects with congenital heart disease is a well-established field on its own and, programmatically, is a fundamental component of any center providing care for these patients.

Excellence in the field of interventional cardiology is obtained with passion, competence, hard study and attention to details.

This book is the result of the collaboration of more than 30 world renowned experts from 4 continents and gives plenty of details, tips and tricks on more than 40 different topics.

It provides a practical guide to the large majority of procedures performed during daily life, and its step-by-step approach will be a precious tool during hard times in the catheterization laboratory.

Almost everything is in the book, to the reader the effort to put in hard work and passion!

Milan, Italy

Milan, Italy

Munich, Germany

Leeds, UK

Gianfranco Butera

Massimo Chessa

Andreas Eicken

John Thomson

Contents

Part I General Issues

| | |
|--|-----------|
| 1 Patient Information and Informed Consent . . . | 3 |
| Maarten Witsenburg | |
| 2 Anaesthesiological Management of the Paediatric Patient in the Catheterisation Laboratory | 7 |
| Giuseppe Isgrò and Marco Ranucci | |
| 3 Antibiotics and Anticoagulation | 17 |
| Luciane Piazza | |
| 4 Angiography: Basics and Contrast Media | 23 |
| Jose Luis Zunzunegui | |
| 5 Angiography: Radiation Exposure and Standard Projections | 37 |
| Federico Gutierrez-Larraya, Angel Sanchez-Recalde, and Enrique Balbacid-Domingo | |

| | | |
|--------------------------------|--|-----|
| 6 | Catheters and Wires | 59 |
| | Adam Koleśnik and Grażyna Brzezińska-Rajszyś | |
| 7 | Balloons | 85 |
| | Caroline Ovaert | |
| 8 | Stents | 105 |
| | Eric Rosenthal and Sebastian Goreczny | |
| 9 | Hemodynamics: Pressures and Flows | 125 |
| | Juan Pablo Sandoval Jones and Lee Benson | |
| 10 | Psychological Aspects in Congenital Heart Disease: A Lifelong Perspective | 149 |
| | Edward Callus, Cristina Farè, and Emilia Quadri | |
| Part II Vascular Access | | |
| 11 | The Usual Vascular Access | 159 |
| | Jochen Weil | |
| 12 | Unusual Access | 171 |
| | Stephan Schubert and Felix Berger | |
| 13 | Hemostasis | 181 |
| | Zakhia S. Saliba, Sami G. Slaba, and Elie B. Sawan | |
| 14 | Access Complications and Management | 193 |
| | Zakhia S. Saliba, Georges S. Tabet, and Chady C. Kallassy | |
| 15 | Transseptal Access | 205 |
| | Chad Kliger and Carlos E. Ruiz | |

Part III Fetal Procedures

- 16 Fetal Interventions** 225
 Carlos A.C. Pedra and Simone Fontes Pedra

**Part IV Step-by-Step Procedures:
Valve Dilatation**

- 17 Aortic Valvular Stenosis** 243
 Xiangbin Pan

- 18 Pulmonary Valve Stenosis** 261
 Tingliang Liu and Wei Gao

- 19 Pulmonary Atresia and Intact
Ventricular Septum** 279
 Zaheer Ahmad and Mazeni Alwi

- 20 Percutaneous Transcatheter Balloon Mitral
Commissurotomy** 299
 Raghavan Subramanyan
 and Anuradha Sridhar

**Part V Step-by-Step Procedures:
Vessel Treatment**

- 21 Pulmonary Artery Branches** 319
 Peter Ewert and Andreas Eicken

- 22 Aortic Coarctation** 335
 Raul Ivo Rossi Filho
 and João Luiz Langer Manica

- 23 Reopening of Peripheral
and Central Arteries and Veins** 355
 Henri Justino and Athar M. Qureshi

| | | |
|--|--|-----|
| 24 | PDA Stenting in Duct-Dependent Pulmonary Circulation | 375 |
| | Kothandam Sivakumar | |
| 25 | Patent Ductus Arteriosus Stenting in Duct-dependent Systemic Circulation | 401 |
| | Dietmar Schranz | |
| Part VI Step-by-Step Procedures: Closing or Creating a Defect | | |
| 26 | Step-by-Step Closure of Atrial Septal Defects (ASDs) | 421 |
| | John D.R. Thomson | |
| 27 | Fontan Fenestration Closure | 449 |
| | Derize E. Boshoff and Marc H. Gewillig | |
| 28 | Ventricular Septal Defects | 465 |
| | Massimo Chessa and Gianfranco Butera | |
| 29 | Patent Ductus Arteriosus Closure | 489 |
| | Ahmed Mohammed Alkamali | |
| 30 | Catheter Closure of Coronary Artery Fistula .. | 507 |
| | Kothandam Sivakumar, Ajit Mallasari, and Bharat Dalvi | |
| 31 | Vessel Embolization: Transcatheter Embolization of Pulmonary Arteriovenous Malformations and Aortopulmonary Collateral Arteries | 527 |
| | Liang Tang, Zhen-fei Fang, and Sheng-hua Zhou | |

| | | |
|---|--|------------|
| 32 | Closure of Residual Postsurgical Defects | 549 |
| | Gerrit Kaleschke and Helmut Baumgartner | |
| 33 | ASD Closure in Special Situations: Elderly, PA-IVS | 561 |
| | Giuseppe Santoro, Luca Giugno, Cristina Capogrosso, Gianpiero Gaio, and Maria Giovanna Russo | |
| 34 | Creating an Interatrial Communication | 571 |
| | Derize E. Boshoff and Marc H. Gewillig | |
| Part VII Step-by-Step Procedures: Valve Implantation | | |
| 35 | Melody Valve Implantation in Pulmonary Position | 593 |
| | Gianfranco Butera, Massimo Chessa, and Philipp Bonhoeffer | |
| 36 | SAPIEN Valve Implantation in the Pulmonary Position | 611 |
| | Noa Holoshitz and Ziyad M. Hijazi | |
| 37 | Percutaneous Tricuspid Valve Implantation . . . | 633 |
| | Andreas Eicken and Peter Ewert | |
| Part VIII Step-by-Step Procedures: Principles of Hybrid Approach | | |
| 38 | Hybrid Approach in Hypoplastic Left Heart Syndrome (HLHS) | 649 |
| | Dietmar Schranz | |

| | |
|--|-----|
| 39 Hybrid Approach: Defect Closure | 667 |
| Gareth Morgan and Eric Rosenthal | |
| 40 Hybrid Approach: Stent Implantation | 679 |
| Ralf J. Holzer | |
| Part IX Step-by-Step Procedures: Miscellanea | |
| 41 Retrieval Techniques | 697 |
| Rui Anjos, Inês C. Mendes, and Duarte S. Martins | |
| 42 Pericardiocentesis | 715 |
| Maarten Witsenburg | |
| 43 Endomyocardial Biopsies | 721 |
| Davide Marini | |
| 44 Evaluations Before Partial and Total Cavopulmonary Connections | 735 |
| Gabriella Agnoletti | |
| 45 Hemodynamics in Pericardial and Myocardial Diseases | 749 |
| Giacomo Pongiglione and Maria Giulia Gagliardi | |
| 46 Imaging and Treating Coronary Arteries in Children | 765 |
| Teiji Akagi | |
| Part X Tables and Charts: Appendix | |
| 47 Hemodynamic Formulae, Calculations, and Charts | 779 |
| Lee N. Benson and Juan Pablo Sandoval Jones | |

Part I
General Issues

Chapter 1

Patient Information and Informed Consent

Maarten Witsenburg

1.1 Introduction

Interventional (and diagnostic) catheterization is an important tool in congenital heart disease. It has evolved from atrial septostomy in the 1970s to a wide range of procedures including device closure of various defects and percutaneous valve implantation nowadays.

As any form of invasive study or treatment, it is not without risks and serious complications may occur. Therefore, it should only be performed after balancing the advantages and risks of the procedure [1]. The risk associated with the use of ionizing radiation for these procedures should be kept in mind, especially because of the young age of many of these patients.

The patient (or his or her legal representative) has to agree on the suggested treatment, but can only do so after having been

M. Witsenburg
Departments of Congenital and Pediatric Cardiology, Thoraxcentre,
Erasmus MC, 2040, 3000 CA Rotterdam, The Netherlands
e-mail: m.witsenburg@erasmusmc.nl

informed appropriately. The combination of the duty to inform and the agreement of the patient with the treatment plan is called informed consent.

Informed consent is an essential step in any diagnostic or interventional cardiac catheterization in a patient with congenital cardiac disease.

1.2 Background

Healthcare ethics is based on the moral concepts of benevolence, autonomy, absence of malice, equity, and responsibility. Autonomy implies that the patient (and/or the legal representatives) can only consent after the provision of adequate information. The major elements in a valid consent process are sufficient understanding, sufficient information, and freeness from duress [2, 3].

In the ESC-EACTS myocardial re-vascularization guidelines, it is stated that information should be “objective and unbiased, patient oriented, evidence based, up-to-date, reliable, understandable, accessible, relevant, and consistent with legal requirements” [3].

1.3 Information and Consent in Clinical Practice

In a non-emergent setting, the indication for a diagnostic or interventional cardiac catheterization should be discussed within a multidisciplinary team including at least the (pediatric) cardiologist, interventional cardiologist, and cardiac surgeon. For non-complex cases a written and locally approved protocol can be an alternative for the discussion within the multidisciplinary team.

In such a heart team, the indication, risks and benefits, possible other treatment options, and timing of procedure are discussed. This team decision is written down in the patient record, as well as the team members who were involved in the discussion.

Once the decision is made, the patient (or legal representative) is informed. It is important to take enough time to discuss the reason for treatment, its timing, risks, and possible treatment complications. One should realize that a lay person as a patient will always have a major lack of knowledge, even after an extensive discussion with the interventional cardiologist. The consent will therefore for a major part be based on the patient's trust in the treating physician. After the patient has consented, this is documented in the patient record. Depending on local rules and practice, the consent can be given orally or in writing.

In emergencies, time may be lacking to fulfill the steps mentioned above. A typical example is severely hypoxic neonate with d-transposition in need of urgent balloon atrial septostomy to improve atrial mixing. In such cases the information needs to be given after the procedure, including explanation of possible complications that may have occurred.

1.4 Conclusion

Recommendations for treatment in congenital heart disease will rarely have a higher than 1C level of evidence. As such expert opinion plays a major role. Even for procedures that have been used extensively for many years, the implications, including complications, have become clearer recently.

In addition the availability of an extending range of devices might sometimes result in using these for questionable indications.

The important point is that any interventional cardiologist should act in a responsible way before, during, and after the intervention. Whenever complications may have happened, he should be able to explain the problem, both to the patient and to colleagues, and how steps were taken to minimize any further harm.

References

1. Feltes TF, Bacha E, Beekman RH 3rd et al (2011) Indications for cardiac catheterization and intervention in pediatric cardiac disease: a scientific statement from the American Heart Association. *Circulation* 123:2607–2652
2. Bonnet C, Greffier A (2013) Paediatric cardiac catheterization: an information sheet. *Arch Cardiovasc Dis* 106:228–237
3. Wijns W, Kolh P, Danchin N, Di Mario C, Falk V, Folliguet T, Garg S, Huber K, James S, Knuuti J, Lopez-Sendon J, Marco J, Menicanti L, Ostojic M, Piepoli MF, Pirllet C, Pomar JL, Reifart N, Ribichini FL, Schalij MJ, Sergeant P, Serruys PW, Silber S, Sousa Uva M, Taggart D (2010) Guidelines on myocardial revascularization. Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS), European Association for Percutaneous Cardiovascular Interventions (EAPCI). *Eur Heart J* 31(20):2501–2555

Chapter 2

Anaesthesiological Management of the Paediatric Patient in the Catheterisation Laboratory

Giuseppe Isgro and Marco Ranucci

2.1 Introduction

The widespread use of therapeutic cardiac catheterisation in the management of congenital heart disease requires the presence of a trained paediatric cardiac anaesthesiologist with the ability to provide both safe and consistent sedation or general anaesthesia to paediatric cardiac patients. Specific knowledge of the pathophysiology of congenital cardiac lesions and the clinical implications of diagnostic and therapeutic procedures are essential.

Sedation is often preferred to general anaesthesia, in particular for diagnostic procedures, because mechanical ventilation can cause haemodynamic disturbance and can alter the results of the study. General anaesthesia is applied mainly for interventional procedures (i.e. percutaneous valve implantation, atrial septal defect or ventricular septal defect closure, patent ductus

G. Isgro, MD • M. Ranucci, MD, FESC (✉)

Department of Cardiothoracic and Vascular Anaesthesia and Intensive Care, IRCCS Policlinico San Donato, San Donato Milanese, Milan, Italy
e-mail: cardioanesthesia@virgilio.it

G. Butera et al. (eds.), *Cardiac Catheterization for Congenital Heart Disease: From Fetal Life to Adulthood*,

7

DOI 10.1007/978-88-470-5681-7_2, © Springer-Verlag Italia 2015

arteriosus closure, or STENT implantation), during which it is essential to keep the patient deeply relaxed to permit the precise deployment of the device.

Other factors influencing the decision to use sedation or general anaesthesia are patient related: age, clinical conditions, and complexity of cardiac lesions.

Diagnostic and interventional procedures in the catheterisation laboratory carry risks for the patient such that continuous patient monitoring is essential.

The anaesthesiologist should contribute to the treatment of complications associated with cardiac catheterisation and, obviously, pre-empt and manage issues arising from sedation and anaesthesia. Finally good anaesthetic practice means that after the procedure the patient is delivered to a post-anaesthesia care unit or directly to the intensive care unit in the best condition possible.

2.2 Anaesthesia

2.2.1 Preoperative Consideration

Preoperative clinical evaluation is mandatory to assess the general condition of the patient and the type of cardiac disease and make plans for post-procedural care. Those patients affected by severe cyanosis should be hydrated prior to cardiac catheterisation, to minimise dehydration.

Fasting should be planned according to the age, clinical condition, and related laboratory investigations.

Routine preoperative tests (ECG, chest X-ray, lab investigations) are required and evaluated by the anaesthesiologist—in some cases, review of echocardiography. An assessment should include scrutiny of previous anaesthetic records and prior premedication.

Certain patients including chronically cyanotic patients are at risk of post-procedural bleeding, so that packed red cell units, fresh frozen plasma and concentrated platelet units are quickly available according to the procedure.

Strict attention to intercurrent illness is required, and if necessary, the catheterisation procedure should be postponed whilst this resolves.

2.2.2 Premedication

Drugs for premedication are administered to reduce anxiety and promote cooperation. Additional benefits include induction of anaesthesia without memory of this stressful time and reduced adrenergic stimulation that can be deleterious, particularly certain anomalies (i.e. tetralogy of Fallot, uncompensated ventricular septal defect with pulmonary hypertension and anomalous origin of left coronary artery arising from the pulmonary artery).

Children under 6 months of age or those that are very sick often can be managed without premedication as this can be deleterious under some circumstances.

Many drugs are available for premedication; the most commonly used are ketamine, midazolam, fentanyl and morphine. Dexmedetomidine, a new centrally acting alpha 2-adrenoceptor agonist, has been used in the setting of cardiac catheterisation laboratory safely with good results.

The choice of the drug alone or in combination must be decided by the anaesthesiologist after assessment of the patient and according to local experience and protocols.

2.2.3 Sedation and Anaesthesia

Sedation and general anaesthesia can be administered according to the preoperative condition, including the risk of developing

Table 2.1 Suggested anaesthetic drugs

| Drug | Induction | Maintenance |
|-----------------|---------------|----------------------|
| Ketamine | 0.5–2 mg/kg | 0.01–0.05 mcg/kg/min |
| Midazolam | 0.1–0.3 mg/kg | 1–3 mcg/kg/min |
| Propofol | 1–2 mg/kg | 3–5 mg/kg/h |
| Sevoflurane | 3–5 % | 1–2 % |
| Fentanyl | 3–5 mcg/kg | 1–2 mcg/kg/min |
| Morphine | 0.1 mg/kg | 1–2 mcg/kg/min |
| Cisatracurium | 0.1–0.2 mg/kg | 1–2 mcg/kg/min |
| Dexmedetomidine | 1 mcg/kg | 0.2–1.4 mcg/kg/h |

post-procedural deleterious effects related to cardiac catheterisation and anaesthesia drugs use (i.e. pulmonary hypertension).

Pathophysiology of any cardiac lesion should be discussed beforehand with the paediatric cardiologist to reduce the risk of anaesthesia delivery, although modern anaesthesia drugs have reduced impact on cardiovascular system (Table 2.1). Sevoflurane, a volatile anaesthetic, has very little effects on systemic pressure and heart rate. Dexmedetomidine is thought to be protective for postoperative atrial fibrillation.

Midazolam is safely used to maintain sedation, usually in combination with fentanyl or morphine.

The use of muscle relaxants that permit to keep the patient ventilated under general anaesthesia is nowadays safe, because the introduction of many newer agent with low rate of adverse effects; the combination of modern volatile anaesthetics and modern muscle relaxants have reduced to very rare event the incidence of malignant hyperthermia.

Cisatracurium, a non-depolarising muscle relaxant, a cis-isomer of atracurium, is indicated in paediatric anaesthesia because of the absence of histamine release; its half-life is 22–29 min and it is eliminated through the Hoffman metabolism, so it can be used safely in patients with poor renal function.

2.3 Monitoring and Anaesthetic Equipment in the Cardiac Catheterisation Laboratory

2.3.1 *Electrocardiogram*

Electrocardiogram is used for continuous monitoring of heart rate, rhythm and ST changes throughout the pre-, intra- and post-procedural phases.

2.3.2 *Blood Pressure*

Systemic blood pressure may be monitored noninvasively during the most common procedures by an automated oscillometric technique.

During risky procedures or in very sick patients, it may be necessary to monitor invasive blood pressure, by cannulation of an artery. Thereby arterial cannulas, transducers, and flushing devices must be present in the laboratory.

2.3.3 *Pulse Oximetry*

It provides a continuous and noninvasive monitor of oxygen saturation, which is mandatory during both sedation and general anaesthesia in paediatric cardiac patients, who are at risk for the development of hypoxia.

2.3.4 *Capnometry*

Is a continuous and noninvasive method of measurement of expired carbon dioxide and is very useful to monitor the

adequacy of ventilation during general anaesthesia or to detect malfunction or failure of the anaesthesia machine. Moreover, it provides a useful information related to the quality of pulmonary perfusion and can reflect haemodynamic changes.

2.3.5 Temperature Monitoring

Temperature monitoring is very important especially in the newborns, who are at risk for hypothermia because of their relatively large surface area and the inefficiency of their thermoregulatory mechanisms. Cutaneous temperature may be monitored by adequate probes. Central temperature, if required, can be measured using a nasogastric probe.

In order to avoid hypothermia in children, it is important to warm the environment and the inhalatory gases by a humidifier. Warming of intravenous fluids may be needed. The use of heating blankets is also recommended especially in the newborn.

2.3.6 ScvO₂ (Continuous Mixed Venous Oxygen Saturation) Monitoring

Paediatric and adult patients with severe cardiac disease, who undergo catheter laboratory interventional procedures, can be monitored with respect to cardiac output. In congenital heart disease patients, it is usually either not possible or desirable to insert a pulmonary artery catheter designed for output measurement. Currently central venous catheters with oximetry are available to continuously monitor venous saturations. These catheters are usually inserted into the right jugular internal vein like a normal central venous line, with the same dimension and length (PediaSat and PreSep catheters—Edwards Lifesciences, Irvine, CO) (Fig. 2.1).




Fig. 2.1 (a, b) PediaSat catheter (paediatric) (a) and PreSep catheter (adult) (b) (Courtesy of Edwards Lifesciences)

The continuous monitoring of venous saturation can help to identify sudden changes in haemodynamic status, rapidly changing when cardiac output decreases or increases.

This parameter is included also in the management of the early goal-directed therapy (EGDT) for critically ill patients.

2.3.7 NIRS (*Near-Infrared Spectroscopy*) *Monitoring*

Another tool of haemodynamic monitoring is near-infrared spectroscopy (NIRS) (Fig. 2.2). NIRS is used in many clinical situations to continuously monitor cerebral and splanchnic perfusion and has the potential to provide information on the adequacy of systemic oxygen delivery. Some authors have demonstrated a good correlation between NIRS and $ScvO_2$, but NIRS cannot precisely predict $ScvO_2$ value, though it can be used for trend monitoring.

Fig. 2.2 NIRS monitoring

2.3.8 Anaesthetic Equipment

- Different sizes of cannulas for venous and arterial cannulation
- Different sizes of central venous catheters
- Different sizes of face masks
- Different sizes of endotracheal tubes
- Airway management equipment and difficult airway management equipment available
- Suction apparatus and different sizes of suction catheters
- Mechanical ventilator with inhalatory anaesthetic agents
- Scavenging setup for waste inhalational agents
- Sedative, analgesic and anaesthetic drugs
- Resuscitation drugs
- Intravenous infusion set – intravenous fluids (crystalloids and colloids)
- Defibrillator
- Stethoscope