



DENTAL EMERGENCIES

EDITED BY
MARK GREENWOOD | IAN CORBETT



 WILEY-BLACKWELL

Contents

Cover

Title Page

Copyright

List of Contributors

Preface

Acknowledgements

Chapter 1: Introduction, Infection Control and Prescribing

Introduction to the dental emergency clinic
Infection and infection control
Prescribing
Conclusions

Chapter 2: History Taking and Clinical Examination of Patients on a Dental Emergency Clinic

Introduction
History taking
Patient examination

Special tests
Record keeping
Consent
Conclusions

Chapter 3: Radiology and the Dental Emergency Clinic

Introduction
Patients in pain
Trauma radiology
Looking at radiographs
Conclusions

Chapter 4: Acute Oral Medical and Surgical Conditions

Introduction
Oro-facial swelling
Blistering disorders of the oral mucosa
Oral ulceration
Disturbed oro-facial sensory or motor function
Haemorrhage
Other acute conditions
Bony pathology
Summary

Chapter 5: Restorative Dental Emergencies

Introduction

General principles

Pain management

Infections and soft tissue problems

Crack, fracture and mobility of teeth and dental restorations

Fractured and loose implants

Fractures and swallowing of removable prostheses

Conclusions

Chapter 6: Acute Presentations of Chronic Oro-Facial Pain Conditions

Introduction

Oro-facial pain history

Examination of patients with oro-facial pain

Special investigations for oro-facial pain

Presentation, investigations and initial management of acute non-odontogenic oro-facial pain

Chapter 7: Traumatic Injuries to the Teeth and Oral Soft Tissues

Assessment of the traumatised patient

Management of traumatic dental injuries

Injuries to the hard dental tissues and the pulp

Injuries to the hard dental tissues, the pulp and the alveolar process

Injuries to the periodontal tissues

Dento-alveolar fractures
Conclusions

Chapter 8: Pain Relief in the Dental Emergency Clinic

Introduction

General mechanism of pain

Distribution of pain fibres in the mouth and jaws

Sources of pain in the mouth and jaws

Control of pain

Psychology of pain

Medication

The ladder of analgesia

Avoiding problems in prescribing analgesics

Chapter 9: Management of the Special Needs Patient

Introduction

Commonly seen conditions

Assessment

Medical history in the patient with special needs

Informed consent

Examination

Factors to consider in treatment

Conclusion

Chapter 10: Making a Referral

Introduction

When to refer

How to refer

Where to refer

The referral letter

Urgency

Cancer referrals

Copies of the referral letter

Summary

Chapter 11: Medical Emergencies in the Dental Emergency Clinic - Principles of Management

Introduction

Contents of the emergency drug box

The 'ABCDE' approach to an emergency patient

Airway (A)

Use of defibrillation

Principles of management after the initial treatment of a medical emergency

Conclusions

Chapter 12: Examples of Specific Medical Emergency Situations

Introduction

Vasovagal syncope (simple faint)

Hyperventilation

Asthma

Cardiac chest pain
Epileptic seizures
Diabetic emergencies
Allergies/hypersensitivity reactions
Adrenal insufficiency
Stroke
Local anaesthetic emergencies
Problems with haemostasis
Conclusions

Appendix 1: Normal Reference Ranges

Appendix 2: Changes in Vital Signs in Patients with Infection

Appendix 3: Warfarin Protocol

Principles
Should the INR be measured before a dental procedure?
Are there any drug interactions that are relevant to this patient group?
Oral surgery patients
Restorative patients
Checklist for patients taking warfarin who require dental treatment

Appendix 4: Aspects Relating to Local Anaesthetics

[Appendix 5: NICE Guidelines for the Removal of Wisdom Teeth](#)

[Appendix 6: Protocol for Surgical Dental Treatment of Patients Taking Bisphosphonates](#)

[Introduction and background](#)

[Management of patients taking bisphosphonates](#)

[Appendix 7: Common ‘Recreational’ Drugs](#)

[Introduction](#)

[Index](#)

Dental Emergencies

Edited by

Mark Greenwood

MDS, PhD, FDS, FRCS, FRCS (OMFS), FHEA

*Consultant and Honorary Clinical Professor
Newcastle University*

Ian Corbett

BDS, BSc, PhD, FDS (OS), RCS

*Lecturer in Oral and Maxillofacial Surgery
Newcastle University*

 **WILEY-BLACKWELL**

A John Wiley & Sons, Ltd., Publication

This edition first published 2012
©2012 by Blackwell Publishing Ltd

Wiley-Blackwell is an imprint of John Wiley & Sons, formed by the merger of Wiley's global Scientific, Technical and Medical business with Blackwell Publishing.

Registered office: John Wiley & Sons, Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial offices: 9600 Garsington Road, Oxford, OX4 2DQ, UK

The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

2121 State Avenue, Ames, Iowa 50014-8300, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell.

The right of the authors to be identified as the authors of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. This publication is designed to provide accurate and authoritative information in regard to the subject matter covered. It is sold on the

understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

Library of Congress Cataloging-in-Publication Data

Dental emergencies / edited by Mark Greenwood.
p. ; cm.

Includes bibliographical references and index.

ISBN 978-0-470-67396-6 (pbk. : alk. paper)

I. Greenwood, M. (Mark)

[DNLM: 1. Dental Care--methods. 2. Emergencies. 3. Emergency Treatment--methods. WU 105]

617.6'026-dc23

2011042659

A catalogue record for this book is available from the British Library.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

List of Contributors

U. Chaudhry, BDS, MFDSRCS, DHyp

Specialist Registrar in Paediatric Dentistry, Manchester Dental Hospital and Royal Manchester Children's Hospital, Manchester

I.P. Corbett, BDS, BSc, PhD, FDS (OS), RCS

Lecturer in Oral and Maxillofacial Surgery, Newcastle University, Newcastle

J. Durham, BDS, PhD, FDS (OS), RCS, FHEA

Walport Lecturer in Oral and Maxillofacial Surgery, Newcastle University, Newcastle

J. Greenley, BDS, MFDSRCS

Senior Dental Officer, Newcastle Dental Hospital, Newcastle

M. Greenwood, MDS, PhD, FDS, FRCS, FRCS (OMFS), FHEA

Consultant and Honorary Clinical Professor, Newcastle University, Newcastle

C.B. Hayward, BDS, Dip Cons Sed

Associate Specialist, Dental Emergency Clinic, Newcastle Dental Hospital, Newcastle

I.C. Mackie, BDS, MSc, PhD, FDSRCPS

Professor and Consultant in Paediatric Dentistry, Manchester Dental School, Manchester

R.I. Macleod, BDS, PhD, FDSRCS, DDR, RCR

Consultant in Dental and Maxillofacial Radiology, Newcastle

Dental Hospital, Newcastle

U. J. Moore, PhD, FDSRCS

Senior Lecturer in Oral Surgery, School of Dental Sciences,
Newcastle University, Newcastle

A. Moufti, DDS, Dip OMFS, PhD, MFDSRCS

Lecturer in Restorative Dentistry, University of Damascus,
Syria

T. Nugent, BDS, MFDSRCS

Senior Dental Officer, Community Dental Service, Newcastle

P.J. Thomson, BDS, MSc, PhD, FFD, FDS, FRCS

Professor in Oral and Maxillofacial Surgery, Newcastle
University, Newcastle

Preface

The dental emergency clinic is an important area for any dental service. Such departments are usually staffed by clinicians with a variety of backgrounds and levels of experience. This book does not attempt to be exhaustive but is a guide to help clinicians with the management of the wide variety of patients that may present. An attempt is made, where appropriate, to place patient management in an academic context.

Acknowledgements

Thanks are due to the clinicians and nurses who work on the dental emergency clinic at Newcastle Dental Hospital. We would also like to acknowledge the help given by Mrs Beryl Leggatt in the typing of the manuscript.

Where figures or photographs have been taken from other sources, due acknowledgement is given. Thanks are also due to Dr Anna Beattie and Dr Helen Stancliffe for some of the photographs seen in Chapter 11.

Chapter 1

Introduction, Infection Control and Prescribing

M. Greenwood

Introduction to the dental emergency clinic

The dental emergency clinic (DEC) is an important part of the service provided to patients. It is a demanding environment in which to work for main two reasons. First, many patients who attend such departments have a general tendency to avoid dental treatment and view attending such a department as a last resort. Second, from the point of view of the clinicians who work in such clinics, the clinical spectrum is wide, and although there is no remit to provide a specialist service, the boundaries of knowledge and experience for clinicians in certain areas are approaching this. Clinical staff working in these departments need a wide skill mix.

This textbook aims to summarise important areas of knowledge with which practitioners working in the DEC should be familiar. Modern clinical working often requires adherence to protocols, and a summary of some of the more important current management protocols, together with supporting evidence, is provided in the appendices.

For maximum efficiency in any department that deals with emergencies, a system of triage is immensely valuable.

Triage is essentially the process of determining the priority of patients' treatment on the basis of severity of their condition. Triage should result in determining the order and priority of a patient's emergency treatment and occasionally their onward transport. In the DEC, emergency situations include those where the airway may be compromised due to infection or trauma. Such patients must be assessed promptly and referred quickly for onward management. Other patients, who may have sustained trauma, need to be assessed expeditiously, particularly from the point of view of airway and vital signs, and also possible head injury and concomitant injuries, which in some cases may take priority over the facial or dental injuries. More detail in relation to the assessment of trauma patients is given in Chapter 7.

Clearly, it is important that the wide variety and, sometimes, the large number of patients that pass through these departments are handled in an appropriate and a safe manner. In no area does this apply more than the area of infection control, the principles of which are discussed in the following sections.

Infection and infection control

Hand care

The most simple and effective method of preventing healthcare-acquired infections is to undertake effective hand hygiene. The World Health Organisation has produced guidelines that have been widely adapted into the '5 moments for hand hygiene'. These are summarised in Box 1.1.

Box 1.1 The 5 moments for hand hygiene at the point of care

- Before patient contact
- Before aseptic task
- After body fluid exposure risk
- After patient contact
- After contact with patient surroundings

Source: Adapted from WHO Alliance for Patient Safety (2006).

Handwashing is clearly important in the prevention of spread of infection in general and has received significant media attention in recent years. This is largely due to the prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA). *S. aureus* is a bacterium that lives on the skin and in the nose of approximately one in three of the population. Usually, people who carry MRSA do not require treatment and it is no more likely to cause infection than 'ordinary' *S. aureus*, but different antibiotics are used to treat these patients. Screening for MRSA is carried out for new appointees to healthcare posts and hospital inpatients – but not for outpatients. Effective handwashing is critical in the prevention of spread of MRSA.

The other bacterium that has received significant attention, particularly in recent years, is *Clostridium difficile*. This is a bacterium living in the bowel of less than 5% of the healthy adult population. Patients can develop problems if they are brought into contact with contaminated surfaces (which include hands). Unlike MRSA, alcohol gels are not effective against *C. difficile* spores, and therefore, effective handwashing is mandatory.

It is important that healthcare workers remove all hand jewellery (with the exception of wedding bands), are bare below the elbows and do not wear a wristwatch. All cuts and abrasions should be covered with a waterproof adhesive dressing. It is important that, after handwashing, gloves are worn, and these should be changed between each patient and the hands washed again after removing the gloves. Non-sterile medical gloves can be used for examination

purposes, but sterile gloves should be worn for operative procedures.

There is significant individual variation in requirements, but the regular use of an emollient hand cream is important to prevent drying of the skin after frequent handwashing. Contact dermatitis can be significant enough in some practitioners to cause real practical problems with clinical practice. Most organisations now routinely use latex-free gloves as standard.

Sterilisation and disinfection

Sterilisation is defined as the killing or removal of all viable organisms. Concern about the transmissible spongiform encephalopathies such as Creutzfeldt-Jakob disease (CJD) and particularly variant-CJD has improved the level of understanding of prion disease. This has led to a necessary redefinition of sterilisation as the inactivation or removal of all self-propagating biological entities.

Disinfection is the reduction in viable organisms to the point where risk of infection is acceptable.

Antisepsis is a related term, defined as the disinfection of skin or wounds. It is not practically possible or even necessary to sterilise absolutely everything in a dental surgery. Clearly, all surgical instruments must be sterile and anything coming into direct contact with the surgical site should also be sterile. Everything else should be disinfected.

Sterilisation and disinfection methods

Before any attempt is made to sterilise or disinfect an instrument, macroscopically evident contamination should be removed. If this is not done, physical access of the sterilising or disinfecting agent to the object being sterilised may be prevented. Therefore, instruments should be pre-

cleaned and, if they have been in contact with infectious material, pre-cleaning should include adequate disinfection as a first step.

Methods of sterilisation and disinfection include dry or moist heat, a wide variety of gaseous or liquid chemicals, filtration and ionising radiation. The choice depends largely on the nature of the material being treated, the degree of inactivation required and the organisms involved.

In contemporary practice, procedures are followed that are known to result in sterility for different batch sizes and materials. The performance of equipment in terms of the temperature and duration is carefully monitored. The Bowie-Dick tape is one method of ensuring that an autoclave has been functioning effectively. The cross-hatchings turn brown when sterilisation has been achieved ([Figure 1.1](#)). [Figure 1.2](#) shows that the sterilisation has been effective within the packaging itself as the coloured area has changed from yellow to blue.

[Figure 1.1](#) A surgical pack after autoclaving. The cross-hatchings on the Bowie-Dick tape have turned brown indicating that the pack has been successfully sterilised.



Figure 1.2 The sticker at the bottom of the paper has changed from yellow to blue, indicating that sterilisation has been effective within the pack (which is where this form should be placed prior to sterilisation).

DDTA00002-12
Basic Minor Oral Surgery
Customer D-OSL
Created 21-04-2011
Expiration date 15-04-2012
Sterilizer program group
Unit 1419900
Sterile Services MP103

Packed D01 SW
Checked *LS*
Wash Checked In

Clinic/Dept/Th. Checked In (Post Surname)
Clinic/Dept/Th. Checked Out (Post Surname)

Item Count SSD Th In Please Check All Screws Are In Retractors & Speculums Out Wash

THIS TRAY LIST MUST BE RETURNED, SIGNED IN & OUT & ANY SHORTAGES NOTED

4319	1	✓	Minor
9940	1	✓	BP Handle No 3
4335	1	✓	Mitchell Trimmer No4
8873	1	✓	Downs Periosteal Elevator
3411	1	✓	Kiner Retractors
8531	1	✓	Bowdler Henry Retractor
0210	2	✓	Couplands Chisel No. 1 & 3
8557	2	✓	Cryers Elevators left/right
8786	3	✓	Warwick James Elevators Right Left & Straight
0271	1	✓	Alveolotomy Shears 5S
8725	1	✓	Glasgow Bone Nibbling Forcep 659932
8558	1	✓	Alveolotomy File
8514	1	✓	Excavator 125/126
0272	1	✓	Flat Plastic PF1 21
4385	3	✓	Mosquito Artery Fcp curved 4"
9922	1	✓	Spencer Wells Artery Fcp
3406	1	✓	Kiner Needle Holder
8257	1	✓	Suture Scissor
769	1	✓	Ball & Socket Towel Clips (Sunlight)
0813	1	✓	Giles Dissecting Fcp 6" Toothed

Total number of Items: 26 Please initial to confirm all SHARPS have been removed.....

Please initial to confirm all SHARPS have been removed.....

Cycle verification: 120°C for 15-20 min
Class 2 Sterilizing Indicator

Basic Minor Oral Surg
DDTA00002-12
Unit 1419900
Use by 15-04-12

ACCEPT WHEN BLUE / PURPLE

BROUKE TST Control 500W
Part 2 - inside inside pack

Autoclaves

The most common method of sterilisation used in dentistry is by moist heat in an autoclave. The method depends on the use of steam under pressure at temperatures between 121°C and 134°C. It is critical that the autoclave is fully saturated with water vapour and that all other gases are excluded. This method of sterilisation is more efficient than

dry heat as it takes less heat to denature fully hydrated proteins and moist heat releases latent heat of vapourisation, which transfers more energy than dry heat.

In some cases, particularly in individuals at high risk from, or those who have known, prion disease, single-use (disposable) equipment should be used wherever possible. Such items include local anaesthetic syringes ([Figure 1.3](#)), scalpels, saliva ejectors and impression trays. There is an ever-increasing array of disposable equipment being manufactured.

[Figure 1.3](#) A disposable local anaesthetic syringe for use in dentistry. The retractable sheath at the end of the syringe can be drawn over the needle and locked into place as a prevention against needlestick injury.



It is important when dealing with the cleaning of handpieces that the manufacturer's instructions are followed closely. Such equipment should never be completely immersed in disinfectant. It is important to lubricate handpieces appropriately.

Impressions should be rinsed under running cold water to remove macroscopic debris. It is important that further disinfection is carried out according to manufacturer's instructions. The request form to the laboratory that accompanies the impression should highlight known

infections or high-risk groups. The same is true of blood samples that are sent for analysis.

Contamination of surgery water supplies

Most water supplies to dental units will be lined with a biofilm, which provides a reservoir of microbial contamination. Amongst the contaminants is the bacterium *Legionella*, responsible for Legionnaire's disease. Some dental units have a bottle-fed water system, in which case disinfectants can be relatively easily supplied. It is important that manufacturer's instructions are followed. Some of the dental units that are supplied with water more remotely must be decontaminated in conjunction with manufacturer's instructions, and some organisations have introduced decontamination units using ethene oxide plant to perform this function. It is important that such lines are run through for the recommended length of time to reduce contamination. This is particularly important when the unit has been out of use.

Healthcare workers in the dental emergency clinic

All those who carry out exposure-prone procedures should be non-infectious or immune to hepatitis B virus, usually via appropriate immunisation. Any worker who has reason to believe that they may be infectious with a blood-borne virus has a professional and ethical duty to report this and to obtain appropriate counselling and testing where relevant. Dependent on the result to such testing, changes to the clinical practice of the individual may be required ranging from modifications of practice to ceasing clinical practice altogether.

All members of the team should have training in the principles of infection prevention and control and this should be updated on a regular (usually, annual) basis. It is important that regular reviews are made of the infection control policies and procedures and that these are updated as required.

Inoculation injuries

Inoculation injuries are defined as incidents where infected objects or substances breach the integrity of the mucous membranes or skin or where a contaminant comes into contact with the eyes.

There are local protocols in place for dealing with such incidents. Clearly, prevention is best and attention to zoning in the surgery ('clean' and 'dirty' areas), and proper disposal of contaminated equipment is vital in this respect. If such an injury does occur, it is important that the wound is allowed or encouraged to bleed and washed thoroughly with running water. The local protocol should then be followed. Usually, such protocols involve taking blood samples from both the donor (usually, but not always, the patient) and the recipient of the injury (usually, but not always, the clinician). The blood samples should then be tested for blood-borne viruses and, depending on the result, appropriate counselling obtained. The occupational health service should be informed at the earliest opportunity.

Waste disposal

The safe disposal of waste is critically important in a clinical environment. The main subdivision is into clinical and non-clinical categories. Local protocols may vary but in general terms, clinical waste is disposed of in different coloured plastic bags depending on its nature. It is important that clinicians are familiar with local policies and procedures.

All sharps should be disposed of in a suitable container designed for the purpose. It is important when filling either waste bags or sharps containers that they are not over-filled as this significantly increases the risk of contamination or needlestick injury. Therefore, clinical waste bags must not be filled to a level more than three-quarters full and they should be tied at the neck.

Blood spillages

Prompt attention is required in the event of a blood spillage. Disposable towels should be placed over the spillage and 10,000 parts per million sodium hypochlorite applied and left for 5 minutes before disposing of the towels as clinical waste. The area should be adequately ventilated, particularly during the cleaning up process. The person clearing such waste should wear appropriate personal protective equipment. In this context, such equipment includes a plastic apron, protective eyewear and heavy-duty gloves. It is advisable to use appropriate protective footwear. Other examples of protective personal equipment are given in Box 1.2.

Box 1.2 Examples of personal protective equipment

- Gloves
- Protective eyewear (spectacles/visor)
- Masks
- For surgical procedures, clothing protection such as a surgical gown
- Appropriate footwear

Prescribing

The British National Formulary (BNF) contains excellent guidance on the principles of prescribing. Clinicians in a DEC will need to write prescriptions for some patients and it is important that basic principles are adhered to. An example of a prescription is given in [Figure 1.4](#).

Figure 1.4 An example of a handwritten prescription. Clearly, such prescriptions should be legible and may be typed, but must be indelible.

Pharmacy Stamp

Age: 68 YRS
D.O.B: 6.3.45

Title, Forename, Surname & Address
MRS ANN OTHER
11 PRESUMPTION STREET,
AMYDLE,
ABI 2LP

Please don't stamp over age box

Number of days' treatment
N.B. Ensure dose is stated
5

Endorsements

HOSPITAL PRESCRIBER

HP

Amoxicillin 250 mg tabs
Three times daily
Supply 15 tabs (5 days)

Prescriber's name and initials in block capitals
CLINICIAN. A

Signature of Prescriber
A. Minn

Date
07/01/11

For dispenser
No. of
Prescs.
on form

DENTAL HOSPITAL
RICHARDSON ROAD
NEWCASTLE UPON TYNE

RTD04

NE2 4AZ

NEWCASTLE HOSPITALS

RTD

NHS

FP10NC0608

Box 1.3 summarises some of the important principles of prescription writing.

Box 1.3 Some basic principles of prescription writing

- Prescriptions should be written or typed legibly and in indelible ink.
- All areas of the prescription should be completed and signed.
- It is a legal requirement to state the age of children less than 12 years in precise terms.
- The unnecessary use of decimal points should be avoided, e.g. 3 mg not 3.0 mg.
- Use of a decimal point is acceptable to express a range, e.g. 0.5–1 g.
- It is important that figures are placed either side of a decimal point, even if one of the figures is zero.
- Micrograms and milligrams should not be abbreviated as they can easily be misread.
- The dose and frequency should be stated.
- If medications are 'as required', a minimum dose interval must be specified.
- Names of drugs should not be abbreviated.
- Directions should be written in English but the use of well-recognised Latin terms is acceptable, e.g. tds (three times per day), bd (twice per day), qds (four times per day), prn (when required).
- Unlicensed medications (those without marketing authorisation) can be prescribed but the patient should be informed of this.

Antibiotic prescribing

Good infection control and prudent antibiotic prescribing are important in reducing the spread of antibiotic resistance. In many cases of acute oral infections, antibiotics will be required but dental surgeons should not forget important local measures such as irrigation with chlorhexidine where appropriate, together with the removal where possible of the source of infection and provision of drainage where this is possible. Where it is likely that antibiotics are required for a protracted period, or the infection is clinically significant, it

is good practice to take a pus swab to send for culture and sensitivity testing. Clearly, if this is done, it is important that the result is checked to ensure that the patient has been prescribed an appropriate antimicrobial (results of culture may take 2–3 days).

In patients who have had previous recent antibiotics or who have a complex medical history that impinges on their overall management, it is important that the clinician working in a DEC discusses the case with a microbiologist, who will provide good advice regarding the best way of managing the patient, particularly from a prescribing viewpoint.

Prescribing for children

Drug dosages in children are usually based on body weight (measured in kilograms). For many preparations used in dentistry, an age range is used. It is vitally important that the drug dosages are carefully worked out to ensure that they are appropriate. In the BNF, unless the age is specified, the term ‘child’ applies to persons aged 12 years or less. Dosages according to the age range may vary but the broad age categories are as follows:

- First month (neonate)
- Up to 1 year (infant)
- 1–5 years
- 6–12 years

Prescribing in the elderly

Older people are often taking multiple medications, which increases the likelihood of drug interactions. Altered pharmacokinetics often result in increased sensitivity of older people to the action of drugs. Even in the absence of known renal disease, renal function deteriorates with age, resulting in reduced drug excretion. Hepatic function also

declines with similar potential effects. Low concentrations of serum albumin in chronic ill health can lead to increased free concentrations of protein-bound drugs. Drug absorption is often relatively normal, and therefore, older people often need lower doses, particularly of those drugs that have a narrow therapeutic window.

In the Dental Practitioners Formulary, most antibiotics can be prescribed for the elderly at standard doses. Older patients may be taking iron or calcium preparations, which can impair absorption of tetracyclines. Non-steroidal anti-inflammatory drugs (NSAIDs) should be used with caution, especially in patients who report dyspepsia, have renal disease or cardiac disease requiring treatment with angiotensin-converting enzyme inhibitors. Older people are particularly prone to side effects from drugs that act on the central nervous system, leading to potential confusion, drowsiness and falls.

Elderly patients with impaired vision, mental function or manual dexterity may lead to problems with compliance. It is important that this is borne in mind when prescribing for elderly patients. A summary of points to bear in mind when prescribing for the elderly is given in Box 1.4.

Prescribing in pregnancy

All drugs should be considered to be potentially teratogenic and should, therefore, be avoided wherever possible. This is particularly the case in the first trimester. The BNF should be checked before any prescription is written for a pregnant patient.

The use of adrenaline-containing local anaesthetics is acceptable in pregnant patients. The optimum time for treatment is the second trimester as the danger is least to the foetus at this time. A hypersensitive gag reflex may be observed in some pregnant patients, which may lead to practical difficulties in treatment.

Box 1.4 Considerations in prescribing for the elderly patient

- Ensure that the patient's current treatment regimen and medical conditions are known.
- A tendency to use lower doses, particularly of drugs that affect the CNS.
- Bear in mind potential difficulties with compliance (child-proof drug bottles with tops that an elderly person cannot remove).
- The British National Formulary has an informative section on prescribing for the elderly, which should be consulted in cases of doubt.

Intravenous sedation must be avoided in the first trimester and the last month of the third trimester and ideally should be avoided altogether in a pregnant patient. Although nitrous oxide does not appear to be teratogenic, it can interfere with vitamin B₁₂ and folate metabolism and should not be used in the first trimester. If it is used at all, the duration of use should be less than 30 minutes, repeated exposure should be avoided and at least 50% oxygen should be administered.

Breast-feeding

Some drugs can be transferred to the baby in breast milk. The BNF contains a section stating which drugs are implicated.

Prescribing in hepatic impairment

Liver disease is an important consideration when prescribing drugs. There are a variety of potential reasons for this as summarised in Box 1.5.