

Rolf Gleiter, Henning Hopf (Eds.)

Modern Cyclophane Chemistry



**WILEY-
VCH**

WILEY-VCH Verlag GmbH & Co. KGaA

Rolf Gleiter, Henning Hopf (Eds.)

Modern Cyclophane Chemistry

Further Titles of Interest:

N. Krause, A. S. K. Hashmi (Eds.)

Modern Allene Chemistry

Two Volumes

2004, ISBN 3-527-30671-4

R. Mahrwald (Ed.)

Modern Aldol Reactions

Two Volumes

2004, ISBN 3-527-30714-1

T. Takeda (Ed.)

Modern Carbonyl Olefination

Methods and Applications

2004, ISBN 3-527-30634-X

D. Astruc (Ed.)

Modern Arene Chemistry

Concepts, Synthesis, and Applications

2002, ISBN 3-527-30489-4

Rolf Gleiter, Henning Hopf (Eds.)

Modern Cyclophane Chemistry



**WILEY-
VCH**

WILEY-VCH Verlag GmbH & Co. KGaA

Professor Dr. Rolf Gleiter

Institute of Organic Chemistry
University of Heidelberg
Im Neuenheimer Feld 270
69120 Heidelberg
Germany

Professor Dr. Henning Hopf

Institute of Organic Chemistry
Technical University of Braunschweig
Hagenring 30
38106 Braunschweig
Germany

This book was carefully produced. Nevertheless, authors, editors and publisher do not warrant the information contained therein to be free of errors. Readers are advised to keep in mind that statements, data, illustrations, procedural details or other items may inadvertently be inaccurate.

Library of Congress Card No.: applied for

A catalogue record for this book is available from the British Library.

**Bibliographic information published
by Die Deutsche Bibliothek**

Die Deutsche Bibliothek lists this publication in the Deutsche Nationalbibliografie; detailed bibliographic data is available in the Internet at <<http://dnb.ddb.de>>.

© 2004 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

All rights reserved (including those of translation in other languages). No part of this book may be reproduced in any form – by photoprinting, microfilm, or any other means – nor transmitted or translated into machine language without written permission from the publishers. Registered names, trademarks, etc. used in this book, even when not specifically marked as such, are not to be considered unprotected by law.

Printed in the Federal Republic of Germany
Printed on acid-free paper

Cover design: The drawings for the cover were provided by Björn Hellbach

Typesetting K+V Fotosatz GmbH, Beerfelden

Printing betz-druck gmbh, Darmstadt

Bookbinding Litges & Dopf Buchbinderei GmbH, Heppenheim

ISBN 3-527-30713-3

Contents

Preface XV

List of Contributors XVII

1	Cyclophynes	1
	<i>Yoshito Tobe and Motohiro Sonoda</i>	
1.1	Introduction	1
1.2	Orthocyclophynes and Related Systems	2
1.2.1	Planar Dehydrobenzoannulenes	3
1.2.2	Nonplanar Orthocyclophynes	13
1.3	Metacyclophynes and Related Systems	17
1.4	Paracyclophynes and 1,3,5-Bridged Cyclophynes	28
1.5	Concluding Remarks	35
1.6	Acknowledgement	36
1.7	References	36
2	Hetera (Cyclo-)phanes	41
	<i>Fritz Vögtle, Gregor Pawlitzki, and Uwe Hahn</i>	
2.1	Introduction	41
2.1.1	Selection of Topics Presented	41
2.1.2	Definitions	42
2.1.3	Why Conduct a Survey on Heteraphanes?	49
2.2	Heteraphanes	51
2.2.1	Planar Chiral and Helical Chiral Phanes	51
2.2.1.1	Design of Phanes with Planar and Helical Chirality	51
2.2.1.2	Hetera [2.2]Metacyclophanes	53
2.2.1.3	Planar Chiral [2.2]Metacyclophanes	55
2.2.1.4	Planar Chiral Hetera [<i>n</i>]Para- and Hetera [<i>n</i>]Metacyclophanes	56
2.2.1.5	Dioxa [2.2]Phanes and Oxaza [2.2]Phanes	57
2.2.1.6	Enantiomer Separations	57
2.2.1.7	Strongly Helical Heteraphanes	58
2.2.2	Catenanes, Rotaxanes, and Knotanes of the Heteraphane Type	59
2.2.2.1	Template Synthesis of Rotaxanes Using Cyclophane Wheels	59

2.2.2.2	Higher Order $[n]$ Rotaxanes via Non-ionic Template Effect	60
2.2.2.3	Combination of Anionic and Non-ionic Template	61
2.2.2.4	Molecular Knots and Similar Macrocycles of the Heteraphane Type	61
2.2.3	Further Heteraphanes, Metallaphanes and Supramolecular Phanes	62
2.3	Conclusions	62
2.4	Acknowledgement	74
2.5	References	74

3 **Highly Strained Cyclophanes** 81

Takashi Tsuji

3.1	Introduction	81
3.2	$[n]$ Metacyclophanes	81
3.2.1	Synthesis	81
3.2.2	Structures and Physical Properties	83
3.2.3	Reactions of Strained $[n]$ Metacyclophanes	84
3.2.3.1	Thermal and Photochemical Reactions	84
3.2.3.2	Addition Reactions	85
3.2.3.3	Reactions with Electrophiles	86
3.2.3.4	Reactions with Nucleophiles	87
3.3	$[n]$ Paracyclophanes	89
3.3.1	Synthesis	90
3.3.2	Structures and Physical Properties	91
3.3.3	Reactions of Strained $[n]$ Paracyclophanes	92
3.3.3.1	Thermal and Photochemical Reactions	92
3.3.3.2	Reactions with Electrophiles	94
3.3.3.3	Diels-Alder and Other Reactions	94
3.3.3.4	Kinetic Stabilization of $[4]$ Paracyclophane Systems	95
3.4	Aromaticity of Bent Benzene Rings	96
3.5	Cyclophanes containing Polycyclic Aromatic Rings: (2,7)Pyrenophanes	96
3.6	$[1.1]$ Paracyclophanes	98
3.6.1	Synthesis	99
3.6.2	Kinetic Stabilization of $[1.1]$ Paracyclophane Systems	100
3.6.3	Structures and Physical Properties	100
3.7	References and Notes	102

4 **Superphanes** 105

Rolf Gleiter and Rolf Roers

4.1	Introduction	105
4.2	$[n_2]$ Cyclopropenonophanes	106
4.2.1	Synthesis	106
4.3	Superbridged Cyclopropenyliophanes	109
4.4	C_4 -Superphanes	109
4.4.1	Properties of Cyclobutadieno Superphanes	115

4.4.2	Oxidative Demetallations	116
4.5	C ₅ -Superphanes	117
4.6	Superbridged Benzene Rings	120
4.7	Concluding Remarks	126
4.8	Acknowledgement	127
4.9	References	127
5	Carbon-Bridged Ferrocenophanes	131
	<i>Joon-Seo Park and T. Randall Lee</i>	
5.1	Introduction	131
5.2	Nomenclature	132
5.3	Mononuclear Carbon-Bridged Ferrocenophanes	133
5.3.1	[1]Ferrocenophanes	133
5.3.2	[2]Ferrocenophanes	134
5.3.3	[3]Ferrocenophanes	136
5.3.4	[4]Ferrocenophanes	140
5.3.5	[5]Ferrocenophanes	144
5.3.6	[<i>m</i>]Ferrocenophanes (<i>m</i> >5)	146
5.3.7	Multiply-Bridged Mononuclear Ferrocenophanes ([<i>m</i>] _{<i>n</i>} Ferrocenophanes)	147
5.4	Multinuclear Ferrocenophanes	150
5.4.1	[0.0]Ferrocenophanes	150
5.4.2	[1 ^{<i>n</i>}]Ferrocenophanes	151
5.4.3	[<i>m</i> ^{<i>n</i>}] Ferrocenophanes	153
5.5	Summary	154
5.6	References	154
6	Endohedral Metal Complexes of Cyclophanes	159
	<i>Rolf Gleiter, Bernhard J. Rausch, and Rolf J. Schaller</i>	
6.1	Introduction	159
6.2	Singly-Bridged Group IVB Metallocenes	160
6.2.1	Singly-Carbon-Bridged Group IVB Metallocenes	160
6.2.2	Singly-Silicon-Bridged Group IVB Metallocenes	161
6.2.3	Singly-Boron- and Phosphorous-Bridged Group IVB Metallocenes	162
6.3	Doubly-Bridged Group IVB Metallocenes	163
6.3.1	Doubly-Carbon-Bridged Group IVB Metallocenes	163
6.3.2	Doubly-Silicon-Bridged Group IVB Metallocenes	165
6.3.3	Structural Features of Doubly-Bridged Group IVB Metallocenes	166
6.4	Endohedral Group VIB and VIIIB Metal π -Complexes	168
6.4.1	Bridged Bis[benzene]chromium Complexes	168
6.4.2	Bridged Metallocenophanes of Group VIIIB Metals	170
6.5	Cavities as Hosts for Cations of Group IB, IIIA and IVA Metals	176
6.5.1	Endohedral Silver Complexes with Cyclophanes	176
6.5.2	Silver Complexes with π -Prismands	180
6.5.3	Group IIIA and IVA Complexes of Cyclophanes	181

6.5.4	Group IIIA Complexes with π -Prismands	182
6.6	Concluding Remarks	184
6.7	Acknowledgments	184
6.8	References	184
7	Intramolecular Reactions in Cyclophanes	189
	<i>Henning Hopf</i>	
7.1	Introduction	189
7.2	Reactions between the Benzene Rings of Cyclophanes	190
7.3	The <i>Pseudo-gem</i> Effect	198
7.4	Intramolecular Reaction between Functional Groups in Cyclophanes	201
7.5	Conclusions	207
7.5	References	208
8	Reactive Intermediates from Cyclophanes	211
	<i>Wolfram Sander</i>	
8.1	Thermolysis of [2.2]Paracyclophanes	211
8.2	Photolytical Cleavage of [2.2]Paracyclophanes	217
8.3	Cleavage of [2.2]Paracyclophanes via Electron Transfer	220
8.4	Cleavage of Cyclophanes with Unsaturated Bridges	221
8.5	Cleavage of Cyclophanes with Carbonyl Groups in the Bridge	223
8.6	References	226
9	X-ray Crystal Structures of Porphyrinophanes as Model Compounds for Photoinduced Electron Transfer	229
	<i>Hermann Irngartinger and Thomas Oeser</i>	
9.1	Introduction	229
9.2	Porphyrin–Quinone Cyclophanes	230
9.2.1	Double-Bridged Porphyrin–Quinone Cyclophanes	230
9.2.2	Single-Bridged Porphyrin–Quinone Cyclophanes	232
9.2.2.1	Phenyl-Spacered Porphyrin–Quinone Cyclophanes	232
9.2.2.2	Naphthalene-Spacered Porphyrin–Quinone Cyclophanes	235
9.3	Porphyrin–Aromatic–Ring Cyclophanes	238
9.3.1	Single Bridge from Opposite <i>meso</i> -Positions	238
9.3.1.1	Substituted Phenyl Rings as Cyclophane Components	238
9.3.1.2	Polycyclic Aromatic Ring-Systems as Cyclophane Components	243
9.3.1.3	Aromatic Heterocycles as Cyclophane Components	245
9.3.2	Single Bridge from Opposite <i>non-meso</i> -Positions	248
9.3.3	Capped Porphyrins	249
9.4	Porphyrinophanes with Fullerene Hosts	253
9.5	Concluding Remarks	254
9.6	References	255

10	Ultraviolet Photoelectron Spectra of Cyclophanes	
	<i>Heidi M. Muchall</i>	259
10.1	Introduction	259
10.2	[2.2]Paracyclophane	263
10.3	Modified Bridges in [2.2]Paracyclophane	265
10.4	Conjugating Substituents in [2.2]Paracyclophane	266
10.5	Donor–Acceptor Cyclophanes	268
10.6	Heterocyclophanes	270
10.7	Miscellaneous Compounds	270
10.7.1	[6]Phanes with Higher Aromatic Systems	270
10.7.2	Cyclopropenophanes	271
10.7.3	Metacyclophanediynes	272
10.8	References	273
11	UV/Vis Spectra of Cyclophanes	275
	<i>Paul Rademacher</i>	
11.1	Introduction	275
11.2	Characteristic Properties of Cyclophanes with Implications for their Electronic Spectra	275
11.3	[<i>n</i>]Cyclophanes	277
11.4	[<i>m.n</i>]Paracyclophanes	279
11.4.1	[1.1]Paracyclophane	279
11.4.2	[2.2]Paracyclophane	280
11.4.3	[<i>m.n</i>]Paracyclophanes	284
11.5	Multibridged [<i>2_n</i>]Cyclophanes and Related Compounds	285
11.6	[<i>m.n</i>]Arenophanes	288
11.6.1	[<i>m.n</i>]Naphthalenophanes	288
11.6.2	[<i>n.n</i>]Pyrenophanes	289
11.7	Fluorinated Cyclophanes	290
11.8	Heterocyclophanes	290
11.9	Multi-layered Cyclophanes	297
11.10	Donor–Acceptor Cyclophanes	298
11.10.1	Donor–Acceptor Substituted [<i>n.n</i>]Paracyclophanes	298
11.10.2	[<i>n.n</i>]Paracyclophane Quinhydrones	300
11.10.3	[<i>n.n</i>]Metacyclophane Quinhydrones	303
11.10.4	Oligooxa[<i>m.n</i>]paracyclophane Quinhydrones and Related Compounds	304
11.10.5	Donor–Acceptor Substituted [<i>n.n</i>]Cyclophanes with Extensive π -Electron Systems	305
11.11	References	307
12	Electronic Circular Dichroism of Cyclophanes	311
	<i>Stefan Grimme and Arnold Bahlmann</i>	
12.1	Introduction	311
12.2	Theoretical Methods	312

12.3	Excited States of Model Compounds	314
12.3.1	Boat-type Deformation in Benzene	314
12.3.2	Interacting Benzene Fragments	319
12.4	Theoretical and Experimental CD Spectra of Cyclophanes	323
12.4.1	[<i>n</i>]Cyclophanes	323
12.4.1.1	9,12-Dimethyl-4-oxa[7]paracyclophane	323
12.4.1.2	[6]Paracyclophane-8-carboxylic Acid	324
12.4.2	[2.2]Paracyclophanes	325
12.4.2.1	[2.2]Paracyclophane	325
12.4.2.2	4-Fluoro-[2.2]paracyclophane	327
12.4.2.3	4-Methyl[2.2]paracyclophane	329
12.4.3	[2.2]Metacyclophanes	331
12.4.3.1	1-Thia[2.2]metacyclophane	331
12.4.3.2	1-Thia-10-aza[2.2]metacyclophane	332
12.4.4	Cyclophanes with Two Different Aromatic Rings	333
12.4.4.1	14,17-Dimethyl[2](1,3)azuleno[2]paracyclophane	333
12.5	Conclusions	334
12.6	Acknowledgement	335
12.7	References	335
13	Fully Conjugated Beltenes (Belt-Like and Tubular Aromatics)	337
	<i>Rainer Herges</i>	
13.1	Introduction	337
13.2	Approaches Towards Fully-Conjugated Beltenes	341
13.3	Belt-like Benzoannulenes (8)	342
13.4	[0 _{<i>n</i>}]Paracyclophanes 7	344
13.5	Möbius Belts	354
13.6	Conjugated Belts from Fullerenes	355
13.7	References	356
14	Molecular Electrochemistry of Cyclophanes	359
	<i>Bernd Speiser</i>	
14.1	Introduction	359
14.1.1	Electron Transfer and Molecular Electrochemistry	359
14.1.2	Molecular Electrochemistry of Cyclophanes	359
14.1.3	Methods of Molecular Electrochemistry	360
14.2	Molecular Electrochemistry of Hydrocarbon Cyclophanes	361
14.3	Molecular Electrochemistry of Functionalized Cyclophanes	363
14.3.1	Substituted Cyclophanes	363
14.3.2	Cyclophanes with Non-metallic Heteroatoms as Ring Members	364
14.4	Molecular Electrochemistry of Organometallic Cyclophane Derivatives	367
14.4.1	A Possible Classification of Organometallic Cyclophane Derivatives	367
14.4.2	Metallocenophanes and Metallametalocenophanes	367

14.4.3	Cyclophanes as π -Complex Ligands	370
14.4.4	Cyclophanes as Supramolecular Components	374
14.5	Conclusions	376
14.6	References	376
15	NMR Spectra of Cyclophanes	381
	<i>Ludger Ernst and Kerstin Ibrom</i>	
15.1	Introduction and Scope	381
15.2	[<i>n</i>]Phanes	382
15.2.1	[<i>n</i>]Metacyclophanes	382
15.2.2	[<i>n</i>]Paracyclophanes	385
15.2.3	Other [<i>n</i>]Phanes	387
15.3	[2.2]Phanes	389
15.4	[3.3]Phanes	396
15.5	[<i>m.n</i>]Phanes (<i>m</i> > 2, <i>n</i> ≥ 2)	400
15.6	[<i>m_n</i>]Phanes	406
15.7	Other Phanes	409
15.8	Conclusion	412
15.9	References	412
16	Strained Heteroatom-Bridged Metallocenophanes	415
	<i>Ian Manners and Ulf Vogel</i>	
16.1	Introduction	415
16.2	Synthesis and Structure	416
16.2.1	[1]Ferrocenophanes	416
16.2.1.1	Group 13 Bridged [1]Ferrocenophanes	417
16.2.1.2	Group 14 Bridged [1]Ferrocenophanes	417
16.2.1.3	Group 15 Bridged [1]Ferrocenophanes	419
16.2.1.4	Group 16 Bridged [1]Ferrocenophanes	419
16.2.2	Other [1]Metallocenophanes	419
16.2.3	[2]Ferrocenophanes	420
16.2.4	Other [2]Metallocenophanes	422
16.3	Ring-Opening Polymerization of Strained Ferrocenophanes	422
16.3.1	Stoichiometric Insertion and Ring-Opening Reactions	422
16.3.2	Ring-Opening Polymerizations (ROP)	423
16.3.2.1	Thermal ROP	423
16.3.2.2	Living Anionic ROP of Silicon-Bridged [1]Ferrocenophanes	424
16.3.2.3	Transition Metal-Catalyzed ROP of [1]Ferrocenophanes	425
16.3.2.4	Other Initiation Methods for ROP	426
16.4	Properties and Applications of Ring-Opened Polyferrocenes	426
16.4.1	Polyferrocenylsilanes	426
16.4.2	Other Polymetallocenes via ROP	430
16.5	References	431

17	Cyclophanes as Templates in Stereoselective Synthesis 435
	<i>Valeria Rozenberg, Elena Sergeeva, and Henning Hopf</i>
17.1	Introduction 435
17.2	Chiral [2.2]Paracyclophanes: Nomenclature and Stereochemical Assignment 435
17.3	The Resolution of Representative Mono- and Disubstituted [2.2]Paracyclophanes 437
17.4	Monosubstituted [2.2]Paracyclophanes as Chiral Inductors 437
17.5	Disubstituted [2.2]Paracyclophanes as Chiral Inductors 440
17.5.1	<i>Ortho</i> - and <i>syn-latero</i> -Substituted Derivatives 440
17.5.2	<i>Pseudo-ortho</i> -Derivatives 449
17.5.3	<i>Pseudo-gem</i> -Derivatives 452
17.6	Chiral Templates from Substituted [2.2]Paracyclophanes as Building Blocks 454
17.7	Stereoselective Reactions in the Side Chain of the Paracyclophanyl Moiety 457
17.8	Concluding Remarks 458
17.9	References 459
18	Vapor-Based Polymerization of Functionalized [2.2]Paracyclophanes: A Unique Approach towards Surface-Engineered Microenvironments 463
	<i>Doris Klee, Norbert Weiss, and Jörg Lahann</i>
18.1	Introduction 463
18.2	Synthesis of Functionalized [2.2]Paracyclophanes 464
18.3	CVD Polymerization of Functionalized [2.2]Paracyclophanes 466
18.4	Immobilization of Bioactive Substances via Functionalized Poly(<i>p</i> -xylylenes) 472
18.4.1	Introduction 472
18.4.2	Surfaces for Blood Contact 475
18.4.3	Surfaces for Tissue Contact 476
18.4.4	Surface Engineering of Microfluidic Devices 478
18.5	Conclusions 481
18.6	References 482
19	From Cyclophanes to Molecular Machines 485
	<i>Amar H. Flood, Yi Liu, and J. Fraser Stoddart</i>
19.1	Introduction 485
19.1.1	Control over the Location and Motion of Moving Parts in Molecular Machines 485
19.2	The Creation of the Tetracationic Cyclophane 486
19.3	Host–Guest Chemistry with the Tetracationic Cyclophane 488
19.3.1	Location Control: Host–Guest Chemistry 488
19.4	Catenane Chemistry with the Tetracationic Cyclophane 492
19.4.1	Going for Gold – The Story of Olympiadane 495
19.5	Rotaxane Chemistry with the Tetracationic Cyclophane 497

19.5.1	Location Control – Catenanes and Rotaxanes	499
19.6	Switchable Rotaxanes, Catenanes and Pseudorotaxanes	499
19.6.1	Controllable Molecular Shuttles	500
19.6.2	Switchable Catenanes	502
19.6.3	Switchable Rotaxanes	503
19.6.4	Photochemically Switchable Pseudorotaxanes	503
19.6.5	Control of Motion	504
19.7	Electronic Devices Containing Molecular Switches	505
19.7.1	A [2]Catenane-Based Electronic Device	506
19.7.2	Bistable [2]Rotaxane Electronic Devices	508
19.7.3	Memory Devices	510
19.7.4	Miniaturization of the Crossbar	510
19.8	Mechanical Devices with Molecular Machines	513
19.9	Conclusions	515
19.10	References	516
20	Molecular Recognition Studies with Cyclophane Receptors in Aqueous Solutions	519
	<i>François Diederich</i>	
20.1	Introduction	519
20.2	Complexation of Aromatic Solutes	520
20.2.1	Polar Effects in Cyclophane Complexation	520
20.2.2	Complexation of Polycyclic Aromatic Hydrocarbons (PAHs) in Water	521
20.2.3	The Combination of Apolar Binding and Ion Pairing Leads to Very High Substrate Selectivity	523
20.2.4	Solvent Dependency of Cyclophane-Arene Complexation and the Nonclassical Hydrophobic Effect: Enthalpic Driving Forces in Aqueous Solution	525
20.3	Steroid Recognition by Cyclophane Receptors	527
20.3.1	The Search for Cyclophanes with Larger Preorganized Cavity-Binding Sites	527
20.3.2	Steroid Complexation by Cyclophanes 9 and 10	529
20.3.3	Double-decker Cyclophanes for Efficient Steroid Complexation: Dissolution of Cholesterol in Water	533
20.4	Catalytic Cyclophanes	536
20.5	Dendritically Encapsulated Cyclophanes (Dendrophanes)	541
20.6	Conclusions	543
20.7	Acknowledgments	543
20.8	References	544
	Subject Index	547

Preface

[2.2]Paracyclophane, first prepared in 1949 by Brown and Farthing and then systematically investigated by Cram and his co-workers from 1951 onward, is now a molecule with a history spanning half a century. A history, however, as confirmed *inter alia* by this monograph, that is not only active and living, but also increasingly extending into novel fields and applications. This is rare for any organic compound and certainly rare for hydrocarbons (excluding such household chemicals such as natural gas and gasoline).

So, what are the reasons for this unabated interest?

The first reason was already obvious to the pioneer workers in this area: The interaction between layered π -systems combined with an unusual three-dimensional structure. Indeed, most of the early studies in cyclophane chemistry were connected to the problem of the “bent and battered benzene rings” (so the title of a classical review article by Cram) and the consequence this structural feature had on the chemical behavior of the cyclophanes. Obviously, the distance between the aromatic subunits (the “decks”) could be altered by changing the nature of the molecular bridges, and one can say that this classical period of cyclophane chemistry culminated in the preparation of superphane by Boekelheide and certainly with the preparation of [1.1]paracyclophane by Tsuji.

What – secondly – had become more and more obvious during these studies was the realization of what may be called the “phane concept”. Just as benzene is the epitome of a planar (flat) aromatic molecule, [2.2]paracyclophane is the example par excellence of a three-dimensional aromatic molecule. Not only did leaving the π -plane lead to the realization that a new, widely variable class of chiral molecules had become available – with all the effects on, for example, stereoselective synthesis – but the bridged aromatic concept could be connected effortlessly to the simultaneously evolving areas of crown ether, cryptand, cavitand, and supramolecular chemistry. To put it another way: numerous supramolecular structures are cyclophanes (and it may be added in passing that – viewed from this angle – there is also a huge group of naturally occurring cyclophanes).

Thirdly, the zwitter character of the cyclophanes became evident during this period. Whereas, for example, adamantane or cubane are only aliphatic molecules, and benzene or the condensed aromatic systems or even the fullerenes are only (highly) unsaturated compounds, the cyclophanes are both: aromatics and aliphatic.

ics (as already indicated by the euphonious name), making the use of both types of chemistries applicable. As far as the aromatic part of the cyclophanes is concerned, this not only meant the application of the typical reactions of classical aromatic chemistry to these compounds, but also the preparation of reactive intermediates derived from the aromatic rings: radical cations, radical anions, etc. Furthermore there was also the unique possibility of combining cyclophane chemistry with the archetypical layered inorganic chemistry, as exemplified by the classics ferrocene and dibenzene chromium. The cyclophanes, however, offer the additional topologically interesting and important possibility of bonding the metal to the interior or the exterior of the organic ligand. Incidentally, the chemical behavior of the molecular bridges in cyclophanes has not been studied to the same extent as the reactivity of the aromatic subunits.

Finally, the crucial test for the significance of a molecular assembly is its practical application. After all, if the properties of a chemical compound are really “interesting”, there must also be practical, industrial applications. With the increased use of cyclophanes as monomers for the preparation of new polymers by CVD techniques with unusual and characteristic properties, polymers not just of importance as chemically very stable surfaces, e.g. for printed circuits, but also as new materials for medical application and drug delivery, cyclophanes are finding more and more applications in materials science. Together with their use in all kinds of optical devices, and as ligands in stereoselective synthesis, we are observing a very rich field developing here.

It is our aim with this volume to show these different developments in cyclophane chemistry. That we succeeded was, of course, only possible because we found a group of excellent contributors and cyclophane specialists willing to give us their time and share their knowledge. As far as the practical side is concerned, we are very grateful to Drs. Gudrun Walter, Bettina Bems from Wiley/VCH, Mary Korndorffer for copy editing and Petra Krämer and C. Mlynek for drawing the numerous and often very complex structures.

We dedicate this volume to the memory of Donald Cram and Virgil Boekelheide.

Heidelberg/Braunschweig
April 2004

Rolf Gleiter
Henning Hopf

List of Contributors

ARNOLD BAHLMANN
Universität Münster
Organisch-Chemisches Institut
Corrensstraße 40
48149 Münster
Germany

FRANÇOIS DIEDERICH
ETH Zürich
Department of Chemistry and Applied
Biosciences
Wolfgang-Pauli-Straße 10
8093 Zürich
Switzerland

LUDGER ERNST
Technische Universität Braunschweig
NMR-Labor der Chemischen Institute
Hagenring 30
38106 Braunschweig
Germany

AMAR H. FLOOD
University of California, Los Angeles
Department of Chemistry
and Biochemistry
405 Milgard Avenue
CA 90095-1569
Los Angeles
USA

ROLF GLEITER
Universität Heidelberg
Organisch-Chemisches Institut
Im Neuenheimer Feld 270
69120 Heidelberg
Germany

STEFAN GRIMME
Universität Münster
Organisch-Chemisches Institut
Corrensstraße 40
48149 Münster
Germany

UWE HAHN
Rheinische Friedrich-Wilhelms-
Universität Bonn
Kekulé-Institut für Organische Chemie
und Biochemie
Gerhard-Domagk-Str. 1
53121 Bonn
Germany

RAINER HERGES
Universität Kiel
Institut für Organische Chemie
Otto-Hahn-Platz 4
24098 Kiel
Germany

HENNING HOPF

Technische Universität Braunschweig
Institut für Organische Chemie
Hagenring 30
38106 Braunschweig
Germany

KERSTIN IBROM

Technische Universität Braunschweig
Institut für Organische Chemie
Hagenring 30
38106 Braunschweig
Germany

HERMANN IRNGARTINGER

University of Heidelberg
Institute of Organic Chemistry
Im Neuenheimer Feld 270
69120 Heidelberg
Germany

DORIS KLEE

RWTH Aachen
Department of Textile
and Macromolecular Chemistry
52074 Aachen
Germany

JOERG LAHANN

University of Michigan
Departments of Chemical Engineering,
Material Science and Engineering,
and Macromolecular Science
and Engineering
3414 G.G. Brown
2300 Hayward
Ann Arbor MI – 48109-2139
USA

T. RANDALL LEE

University of Houston
Department of Chemistry
4800 Calhoun Road
Houston, TX 77204-5003
USA

YI LIU

University of California, Los Angeles
Department of Chemistry
and Biochemistry
405 Milgard Avenue
CA 90095-1569
Los Angeles
USA

IAN MANNERS

University of Toronto
Department of Chemistry
80 St. George Street
Toronto, Ontario M5S 3H6
Canada

HEIDI M. MUCHALL

Concordia University
Department of Chemistry
and Biochemistry and Centre
for Research in Molecular Modeling
Montreal, Quebec H4B 1R6
Canada

THOMAS OESER

University of Heidelberg
Institute of Organic Chemistry
Im Neuenheimer Feld 270
69120 Heidelberg
Germany

JOON-SEO PARK

University of Houston
Department of Chemistry
4800 Calhoun Road
Houston, TX 77204-5003
USA

GREGOR PAWLITZKI

Rheinische Friedrich-Wilhelms-
Universität Bonn
Kekulé-Institut für Organische Chemie
und Biochemie
Gerhard-Domagk-Str. 1
53121 Bonn
Germany

PAUL RADEMACHER
 Universität Duisburg-Essen
 Institut für Organische Chemie
 Universitätsstr. 5–7
 45117 Essen
 Germany

BERNHARD J. RAUSCH
 Clariant BL Pharma Germany
 Molecules Synthesis Centre
 Industriepark Höchst, D 569
 65926 Frankfurt/Main
 Germany

ROLF ROERS
 Bayer MaterialScience AG
 Building B 103
 51368 Leverkusen
 Germany

VALERIA ROZENBERG
 Russian Academy of Science
 A. N. Nesmeyanov Institute
 of Organoelement Compounds
 Vavilova str., 28
 119991 Moscow
 Russia

WOLFRAM SANDER
 Ruhr University Bochum
 Department of Organic Chemistry II
 44780 Bochum
 Germany

ROLF J. SCHALLER
 EMS-Dottikon AG
 Hembrunnstr. 17
 5605 Dottikon
 Switzerland

ELENA SERGEEVA
 Russian Academy of Science
 A. N. Nesmeyanov Institute
 of Organoelement Compounds
 Vavilova str., 28
 119991 Moscow
 Russia

MOTOHIRO SONODA
 Osaka University
 Graduate School of Engineering
 Science, Division of Frontier Materials
 Science
 1-3 Machikaneyama
 Toyonaka 560-8531
 Japan

BERND SPEISER
 Universität Tübingen
 Institut für Organische Chemie
 Auf der Morgenstelle 18
 72076 Tübingen
 Germany

J. FRASER STODDART
 University of California, Los Angeles
 Department of Chemistry
 and Biochemistry
 405 Milgard Avenue
 CA 90095-1569
 Los Angeles
 USA

TAKASHI TSUJI
 Hokkaido University
 Division of Chemistry, Graduate
 School of Science
 Sapporo 060-0810
 Japan

YOSHITO TOBE
 Osaka University
 Graduate School of Engineering
 Science, Division of Frontier Materials
 Science
 1-3 Machikaneyama
 Toyonaka 560-8531
 Japan

ULF VOGEL
 University of Toronto
 Department of Chemistry
 80 St. George Street
 Toronto, Ontario M5S 3H6
 Canada

FRITZ VÖGTLE
Rheinische Friedrich-Wilhelms-
Universität Bonn
Kekulé-Institut für Organische Chemie
und Biochemie
Gerhard-Domagk-Str. 1
53121 Bonn
Germany

NORBERT WEISS
RWTH Aachen
Department of Textile
and Macromolecular Chemistry
52074 Aachen
Germany

1

Cyclophynes

Yoshito Tobe and Motohiro Sonoda

1.1

Introduction

The chemistry of cyclophynes having carbon–carbon triple bond bridges has been one of the most actively investigated fields in modern cyclophane chemistry, particularly in connection with the evolving fields of carbon-rich materials and shape-persistent macrocyclic compounds. In accordance with the remarkable progress in this field over the last decade, a number of reviews has already been published based on the authors' own perspectives [1–9]. Since, in this review, the authors wish to provide a comprehensive survey of cyclophynes chemistry, some overlaps are inevitable.

The properties of cyclophynes are characterized by the geometric and electronic properties of triple bonds and the substitution pattern of the aromatic rings. With regard to the geometrical properties, the macrocyclic frameworks of cyclophynes can be expanded by incorporation of triple bonds because of their linearity. The substitution pattern of the aromatic rings, on the other hand, fixes the direction of the bridging triple bonds, defining the whole molecular shape. As a result, a variety of two- and three-dimensional architectures can be built by connecting aromatic rings with triple bond linkages. The ortho substitution pattern would lead to conjugated cyclophynes, also regarded as dehydrobenzoannulenes (DBAs), with either planar or nonplanar shape depending principally on their ring size. Although planar DBAs had been studied extensively during the late 1950s through mid-1970s in connection with the aromaticity/antiaromaticity of annulenes, these are currently being investigated with renewed interest regarding their potential as opto-electronic materials. On the other hand, nonplanar macrocycles of this type have been studied with regard to their conformation, chirality, and their potential application to liquid crystalline and sensing materials. Some highly unsaturated members of this type of compound have been shown to serve as precursors of ordered carbon materials. Since the meta substitution pattern would make the size of the cyclophynes larger than the ortho analogs, keeping the planarity of the macrocyclic rings, a number of shape-persistent metacyclophynes have been synthesized in order to investigate their supramolecular properties such as guest-binding and self-association in solution, solid state, and at solid–liq-

uid and air–liquid interfaces. To form cyclic structures of para substitution pattern, macrocycles adopt a cylindrical shape, and with 1,3,5-trisubstitution, a cage-like structure can be constructed. These three-dimensional molecules have been studied in view of the host–guest chemistry of curved π -systems, strain of triple bonds, and their transformation into all-carbon molecules.

Concerning the electronic properties of C–C triple bonds, acetylene linkages are capable of transmitting electronic perturbation efficiently from one end of conjugated π -systems to the other because of the isotropic distribution of the π electrons along the C–C axis. This characteristic is responsible for the unique optoelectronic properties predicted for planar cyclic compounds like DBAs. Because C–C triple bonds have a high energy content, it is well known that structurally unhindered alkynes react violently, frequently with explosion. Similarly, diynes are known to undergo topochemical polymerization to form poly(butadiyne)s, potential materials with nonlinear optical properties. The high potential of the triple bonds toward spontaneous chemical reactions has been exploited to provide the transformation of some cyclophynes into all-carbon and carbon-rich materials. In addition, because triple bonds serve as good ligands for transition metals, a number of metal complexes of cyclophynes with unique structures has been readily prepared.

Besides the interest from materials science, recent development of new synthetic methods, particularly those based on transition metal-catalyzed bond formation between sp – sp^2 and sp – sp carbons, have been playing a crucial role in the rapid growth of this field. This chapter covers the recent development of the two- and three-dimensionally shaped cyclophynes from the above points of view. For convenience, the molecules are classified into three types: (1) orthocyclophynes with either planar or nonplanar shape; (2) metacyclophynes; and (3) paracyclophynes and 1,3,5-bridged cyclophynes, on the basis of the substitution pattern. Those compounds with mixed substitution pattern such as ortho, para and meta, para are included in one of the above three types depending on their most characteristic structural and/or functional properties.

1.2

Orthocyclophynes and Related Systems

Fully conjugated planar orthocyclophynes are regarded as dehydrobenzoannulenes (DBAs) [2, 3, 4, 6]. 12-, 14- and 18-membered macrocycles of this type are investigated intensively in connection with their opto-electronic properties, theoretical and experimental inspection of aromaticity, and as substructures of hitherto unknown two-dimensional carbon networks, graphyne and graphdiyne [9]. On the other hand, since the macrocyclic frameworks of larger orthocyclophynes are nonplanar, these molecules are investigated in view of their structure and the chirality of twisted π -systems [7]. In addition, highly unsaturated compounds of this class are studied as possible precursors to nanocarbon materials such as carbon onions and tubes.




Fig. 1.1

1.2.1

Planar Dehydrobenzoannulenes

It was predicted that the two-dimensional carbon network, graphyne, would exhibit semiconductive and nonlinear optical properties [10]. Recent calculations from first principles confirmed that graphyne should be a semiconductor with a moderate band gap and should be quite stable once synthesized [11]. It was predicted that potassium-intercalated graphyne would become metallic with a remarkably short distance (4.30 Å when each 12-membered ring holds a potassium atom) between the intercalant layers [12]. In connection with the nonlinear optical properties of graphyne, hyperpolarizabilities of ladder oligomers of dehydrobenzo[12]annulene ([12]DBA) have been estimated, based on semiempirical calculations, indicating that these can be promising third-order nonlinear optical materials [13]. Force field calculations for networks related to graphdiyne revealed that they would exhibit negative Poisson's ratios, an unusual mechanical property caused by the rotation of triangles [14].

Most cyclophynes were synthesized by using the Stephens-Castro and Sonogashira reactions for the $C(sp^2)-C(sp)$ bond formation or copper(II)-mediated oxidative coupling for $C(sp)-C(sp)$ bond formation. Remarkable improvements in the synthesis of cyclophynes have been achieved as exemplified by the synthesis of the known [12]DBA **1a**. These include (1) the Stephens-Castro reaction of the pre-formed copper(I) acetylide [15]; (2) one-step synthesis by the Sonogashira coupling using acetylene [16]; (3) selective construction of the 12-membered ring based on a masking/protection protocol for both aryl iodide using dialkyltriazene group and alkyne terminal using trialkylsilyl groups [17]; and (4) the alkyne metathesis method [18], which has been developed recently (Scheme 1.1). The first two methods were also utilized for the synthesis of thiophene analogs **2a** [16] and **3** [19]. By stepwise construction of the requisite precursor followed by intramolecular Sonogashira coupling, Haley synthesized bisDBAs **4** and **5a** [17]. Another derivative of the latter compound **5b** was also prepared by an *in situ* deprotection–cyclization method to prove the theoretical prediction for hyperpolarizability of this sys-






Fig. 1.2



Scheme 1.1



Scheme 1.2

tem (Scheme 1.2) [20]. Not only the parent hydrocarbon **1a** but also a number of its derivatives **1b–d** have been prepared by the metathesis method, except for those with substituents adjacent to the alkyne bridges. Similarly, the metathesis method was applied to the synthesis of bisDBA **4** and orthometacyclophylene **6** [18].

Since the late 1980s, Youngs and Tessier have been developing new frontiers in the chemistry of the [12]DBA system [6]. A number of transition metal complexes of **1a** were prepared such as Co(0), Cu(I), Ag(I), and Ni(0) complexes [21]. Significantly among these metal complexes, the Ni(0) complex becomes semiconductive [22]. For example, the Ni(0) complex doped with potassium [2.2.2]cryptate (1:2 ratio) exhibits conductivity of a moderate semiconductor regime ($2 \times 10^{-3} \text{ S cm}^{-1}$). The hexamethoxy derivative **1g**, hexa-alkoxy derivatives **1e–f** having long alkyl chains, and their Ni(0) complexes were also prepared, as well as the quinone **7** and thiophene derivatives **2b–c** [23]. In spite of the expectation based on their structural similarities to hexasubstituted triphenylene derivatives, DBAs **1e–f** and **2b–c** with long alkyl chains did not exhibit liquid crystalline properties.


A dihydro derivative of **1a**, cyclophanediyne **8**, was prepared by Iyoda *et al.* using modified McMurry coupling conditions, followed by dehydroxylation [24]. Although the X-ray structure analysis of **8** revealed its short transannular distance (the shortest C(sp)–C(sp) distance: 2.77 Å), it was inert to photoirradiation. On the other hand, heating **8** at 250 °C in the presence of cyclohexa-1,4-diene resulted in the formation of dibenzonaphthocyclooctatetraene, a product of the Bergman cyclization. The corresponding transannular distances of the oxygen- and sulfur-bridged diynes **9a** and **9b**, respectively, are longer than that of **8** (3.44 and 3.50 Å, respectively), and it is reported that these compounds are inert to thermal and photochemical cyclization [25].

Hexa-ethynyl derivatives **1i–j** were prepared as the key intermediates for the synthesis of antikekulene **12**, an antiaromatic counterpart of kekulene [26]. Thus

the first intramolecular CpCo-catalyzed cyclotrimerization of **1i** and **1j** yielded the triphenylene-based cyclophynes **10a** and **10b**, respectively, whose second cycloisomerization furnished the corresponding pentaphenylene-based cyclophynes **11a** and **11b** (Scheme 1.3). The final cyclization to the target **12** is yet to be realized. Besides this spectacular synthetic endeavor, the parent hydrocarbon **1h** was found to form crystals with highly organized channels including tetrahydrofuran induced by multiple alkyne C–H...O hydrogen bonds [27].

Perethynyl-substituted dibenzodehydro[12]annulene **13a** was prepared by Rubin *et al.* by oxidative coupling of a differentially substituted hexa-ethynylbenzene derivative [28]. Hydrocarbon **13a** can be regarded as a constituent unit of a two-dimensional network with different topology from that of graphdiyne and as a monomer of ladder polymers constructed from the dibenzodehydro[12]annulene ring, called acynes. In this connection, Anthony prepared tetra-ethynyl-substituted [14]DBA **14** having acenaphthene units as a possible monomer unit of the related ladder polymers [29]. In connection with the topochemical polymerization of butadiynes, Komatsu *et al.* prepared octafluoro[12]DBA **13c**, whose reduction potential was more negative than that of the parent hydrocarbon **13b** [30]. In the two crystal forms of **13c**, the molecules are stacked in a slanted manner, while in the co-crystal of **13b** and **13c** (1:2), the molecules are stacked in a columnar arrangement forming a sandwich-like complex. In spite of the favorable packing geometries for topochemical polymerization, none of the crystals underwent photochemical polymerization, probably because of the high potential energy of the polymerization product.

As a quantitative measure of aromaticity of DBAs, the nuclear independent chemical shift (NICS) values at the center of the macrocyclic ring of **1a**, as well as






Fig. 1.3

those for [14]annulene **15**, [16]annulene **16**, and [18]annulene **17a**, were estimated by the DFT method, indicating that **1a** and **16** are moderately antiaromatic while **15** and **17a** are weakly aromatic [31]. Similarly we can predict that highly strained [10]annulene **18** is fully aromatic [31c]. In order to investigate the effect of strain on the tropicity of DBA rings, tetrakisdehydro[12]annulene **19** was generated by photochemical [2+2] fragmentation of the propellane precursor **20** in an argon matrix at low temperature and was characterized by comparison of UV and IR spectra with those calculated by the DFT method (Scheme 1.4) [32]. Unfortunately, since the [12]annulene system **19** was too reactive to observe its tropicity in solution, the next higher homolog **21** was produced by the same method from the homologous precursor **22**. The calculated ^1H NMR chemical shift by the DFT method and experimental observations revealed that strained pentakisdehydro[14]annulene **21** was as diatropic as the corresponding tetrakisdehydro derivative **22**.

Interestingly, the 14-membered DBA ([14]DBA) **15** underwent topochemical polymerization induced by photoirradiation or by applying pressure (20000 psi) [33]. From the packing diagram of the crystal, *trans-trans* geometry of the polymer chain was suggested. This represents the first example of the topochemical polymerization of a conjugated cyclic diyne derivative, although the reactions of acyclic butadiyne derivatives have been well known to form poly(butadiyne)s, which are known to exhibit interesting optical properties. In connection with the theoretical prediction regarding the tropicity of dehydrobenzo[14]annulenes, Haley prepared a number of monobenzo, dibenzo, and monothiopheno derivatives such as **23–27** and studied their diatropcities based on the ^1H NMR chemical shifts [34]. As a result, fusion of a benzene ring to the dehydro[14]annulene ring induces stepwise loss of aromaticity with increasing number of the fused rings. The relative aromaticity (RA) was defined by the chemical shifts relative to those of a non-conjugated reference compound. The experimental RA_{expt} values are in accord with the theoretical values (RA_{NICS}) derived from the calculated NICS values. To further confirm the aromaticity of the [14]annulene system, hybrid systems **28** and **29** formed by fusion of the [14]DBA ring to dimethyldihydropyrene (DHP) unit were prepared [35]. The chemical shift of the probe methyl group in the DHP system of mono-DBA-fused compound **28** was compared with appropriate references. However, the effect of benzoannulation on aromaticity of the [14]DBA ring is not conclusive probably because of the small chemical shift change. Bis-DBA fused system **29** cannot be used for the examination of aromaticity because




Fig. 1.4

the effect of one DBA unit on the DHP probe should counteract that of the other DBA. Since Mitchell *et al.* developed reversible photochemical isomerization of fused DHPs with the corresponding [2.2]metacyclophane systems, the isomerization of **29** was examined, because its absorption band extended to 600–800 nm region, which may be used as “read out” without scrambling of information. However, photochemical isomerization was not successful because the cyclophane form reverts very easily to the starting material **29**. A hybrid system **30** between [14]DBA **15** and [2.2]paracyclophane was synthesized in order to investigate the global transannular delocalization through the overall π system [36]. By comparison of the UV absorption with those of appropriate acyclic reference compounds, it was concluded that **30** sustained π -electron delocalization through the circuit composed of the DBA and [2.2]paracyclophane systems.