



Marcello Lappa

# Rotating Thermal Flows

in Natural  
and Industrial  
Processes



 WILEY



# Table of Contents

[Title Page](#)

[Copyright](#)

[Dedication](#)

[Preface](#)

[A Survey of the Contents](#)

[Acknowledgements](#)

[Chapter 1: Equations, General Concepts and Nondimensional Numbers](#)

[1.1 The Navier-Stokes and Energy Equations](#)

[1.2 Some Considerations about the Dynamics of Vorticity](#)

[1.3 Incompressible Formulation](#)

[1.4 Buoyancy Convection](#)

[1.5 Surface-Tension-Driven Flows](#)

[1.6 Rotating Systems: The Coriolis and Centrifugal Forces](#)

[1.7 Some Elementary Effects due to Rotation](#)

[Chapter 2: Rayleigh-Bénard Convection with Rotation](#)

[2.1 Rayleigh-Bénard Convection with Rotation in Infinite Layers](#)

[2.2 The Küppers-Lortz Instability and Domain Chaos](#)

[2.3 Patterns with Squares](#)

[2.4 Typical Phenomena for  \$Pr \cong 1\$  and Small Values of the Coriolis Number](#)

[2.5 The Low-Pr Hopf Bifurcation and Mixed States](#)

[2.6 Laterally Confined Convection](#)

[2.7 Centrifugal Effects](#)

[2.8 Turbulent Rotating RB Convection](#)

## [Chapter 3: Spherical Shells, Rossby Waves and Centrifugally Driven Thermal Convection](#)

[3.1 The Coriolis Effect in Atmosphere Dynamics](#)

[3.2 Self-Gravitating Rotating Spherical Shells](#)

[3.3 Centrifugally Driven Thermal Convection](#)

## [Chapter 4: The Baroclinic Problem](#)

[4.1 Energetics of Convection and Heuristic Arguments](#)

[4.2 Linear Stability Analysis: The Classical Eady's Model](#)

[4.3 Extensions of the Eady's model](#)

[4.4 Fully Developed Nonlinear Waveforms](#)

[4.5 The Influence of the Prandtl Number](#)

[4.6 The Route to Chaos](#)

[4.7 Hybrid Baroclinic Flows](#)

[4.8 Elementary Application to Atmospheric Dynamics](#)

## Chapter 5: The Quasi-Geostrophic Theory

5.1 The Potential Vorticity Perspective

5.2 The Perturbation Energy Equation

5.3 Derivation of Necessary Conditions for Instability

5.4 A Generalization of the Potential Vorticity Concept

5.5 The Concept of Interlevel Interaction

5.6 The Counter-Propagating Rossby-Wave Perspective on Baroclinic Instability

5.7 Barotropic Instability

5.8 Extensions of the CRW Perspective

5.9 The Over-reflection Theory and Its Connections to Other Theoretical Models

5.10 Nonmodal Growth, Optimal Perturbations and Resonance

5.11 Limits of the CRW Theory

## Chapter 6: Planetary Patterns

6.1 Jet Sets

6.2 A Rigorous Categorization of Hypotheses and Models

6.3 The Weather-Layer Approach

6.4 The Physical Mechanism of Vortex Merging

6.5 Freely Decaying Turbulence

6.6 Geostrophic Turbulence

6.7 The Reorientation of the Inverse Cascade into Zonal Modes

[6.8 Baroclinic Effects, Stochasting Forcing and Barotropization](#)

[6.9 Hierarchy of Models and Scales](#)

[6.10 One-Layer Model](#)

[6.11 Barotropicality, Baroclinicity and Multilayer Models](#)

[6.12 The Ocean-Jupiter Connection](#)

[6.13 Wave-Mean-Flow Dynamics](#)

[6.14 Solitary Vortex Dynamics](#)

[6.15 Penetrative Convection Model](#)

[6.16 Extension and Unification of Existing Theories and Approaches](#)

## [Chapter 7: Surface-Tension-Driven Flows in Rotating Fluids](#)

[7.1 Marangoni-Bénard Convection](#)

[7.2 The Return Flow](#)

[7.3 The Hydrothermal Instability](#)

[7.4 The Annular Pool](#)

## [Chapter 8: Crystal Growth from the Melt and Rotating Machinery](#)

[8.1 The Bridgman Method](#)

[8.2 The Floating Zone](#)

[8.3 The Czochralski Method](#)

[8.4 Rotating Machinery](#)

## [Chapter 9: Rotating Magnetic Fields](#)

[9.1 Physical Principles and Characteristic Numbers](#)

[9.2 Stabilization of Thermo-gravitational Flows](#)

[9.3 Stabilization of Surface-Tension-Driven Flows](#)

[9.4 Combining Rotation and RMF](#)

## [Chapter 10: Angular Vibrations and Rocking Motions](#)

[10.1 Equations and Relevant Parameters](#)

[10.2 The Infinite Layer](#)

[10.3 The Vertical Coaxial Gap](#)

[10.4 Application to Vertical Bridgman Crystal Growth](#)

[References](#)

[Index](#)

# **Rotating Thermal Flows in Natural and Industrial Processes**

MARCELLO LAPPA  
Naples, Italy



A John Wiley & Sons, Ltd., Publication

This edition first published 2012

2012 John Wiley & Sons, Ltd

*Registered office*

John Wiley & Sons Ltd, The Atrium, Southern Gate,  
Chichester, West Sussex, PO19 8SQ, United Kingdom

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at [www.wiley.com](http://www.wiley.com).

The right of the author to be identified as the author of this work has been asserted in accordance with the Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. This publication is designed to provide accurate and authoritative information in regard to the subject matter covered. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.



The publisher and the author make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of fitness for a particular purpose. This work is sold with the understanding that the publisher is not engaged in rendering professional services. The advice and strategies contained herein may not be suitable for every situation. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of experimental reagents, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each chemical, piece of equipment, reagent, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. The fact that an organization or Website is referred to in this work as a citation and/or a potential source of further information does not mean that the author or the publisher endorses the information the organization or Website may provide or recommendations it may make. Further, readers should be aware that Internet Websites listed in this work may have changed or disappeared between when this work was written and when it is read. No warranty may be created or extended by any promotional statements for this work. Neither the publisher nor the author shall be liable for any damages arising herefrom.

*Library of Congress Cataloging-in-Publication Data*

Lappa, Marcello.

Rotating thermal flows in natural and industrial processes /  
Marcello Lappa.

pages cm

Includes bibliographical references and index.

ISBN 978-1-119-96079-9 (hardback)

1. Heat—Transmission. 2. Rotating masses of fluid. I. Title.

QC320.2.L37 2012

536'.2—dc23

2012008650

A catalogue record for this book is available from the British  
Library.

Print ISBN: 978-111-996-0799

*To a red rose*

# Preface

The relevance of self-organization, pattern formation, nonlinear phenomena and non-equilibrium behaviour in a wide range of fluid-dynamics problems in *rotating systems*, somehow related to the science of materials, crystal growth, thermal engineering, meteorology, oceanography, geophysics and astrophysics, calls for a concerted approach using the tools of thermodynamics, fluid-dynamics, statistical physics, nonlinear dynamics, mathematical modelling and numerical simulation, in synergy with experimentally oriented work.

The reason behind such a need, of which the present book may be regarded as a natural consequence, is that in many instances pertaining to such fields one witnesses remarkable affinities between *large-scale-level processes* and the same entities on the *smaller* (laboratory) *scale*; despite the common origin (they are related to 'rotational effects'), such similarities (and the important related implications) are often ignored in typical analyses related to one or the other category of studies.

With the specific intent to extend the treatment given in an earlier Wiley text (*Thermal Convection: Patterns, Evolution and Stability*, Chichester, 2010, which was conceived in a similar spirit), the present book is entirely focused on *hybrid* regimes of convection in which one of the involved forces is represented by standard gravity or surface tension gradients (under various heating conditions: from below, from the side, etc.), while the other arises by virtue of *rotation*.

The analogies and kinships between the two fundamental classes of models mentioned above, one dealing with issues of complex behaviour at the laboratory (technological application) level and the second referring to the strong

nonlinear nature of large-scale (terrestrial atmosphere, oceans and more) evolution, are defined and discussed in detail.

The starting point for such a development is the recognition that such phenomena share an important fundamental feature, a group of equations, strictly related, from a mathematical point of view, to model mass, momentum and energy transfer, and the mathematical expressions used therein for the 'driving forces'.

Although other excellent monographs that have appeared in the literature (e.g. to cite the most recent ones: Marshall and Plumb, 2007, *Atmosphere, Ocean, and Climate Dynamics*, Academic Press; Vallis, 2006, *Atmospheric and Oceanic Fluid Dynamics*, Cambridge University Press) have some aspects in common with the present book, they were expressly conceived for an audience consisting of meteorologists.

Here the use of jargon is avoided, this being done under the declared intent to increase the book's readability and, in particular, make it understandable also for those individuals who are not 'pure' meteorologists (or 'pure' professionals/researchers working in the field of materials science), thereby promoting the exchange of ideas and knowledge integration.

In this context, it is expressly shown how the aforementioned isomorphism between small and large scale phenomena becomes beneficial to the definition and ensuing development of an integrated comprehensive framework, allowing the reader to understand and assimilate the underlying quintessential mechanisms without requiring familiarity with specific literature on the subject.

## **A Survey of the Contents**

In Chapter 1 the main book topics are placed in a precise theoretical context by introducing some necessary notions and definitions, such a melange of equations and nondimensional numbers being propaedeutical to the subsequent elaboration of more complex concepts and theories.

Chapter 2 deals with *Rayleigh-Bénard convection* in simplified (infinite and finite) geometrical models, which is generally regarded as the simplest possible laboratory system incorporating the essential forces that occur in natural phenomena (such as circulations in the atmosphere and ocean currents) and many technological applications (too numerous to list).

The astonishing richness of possible convective modes for this case is presented with an increasing level of complexity as the discussion progresses, starting from the ideal case of a system of infinite (in the horizontal direction) extent in which the role of centrifugal force is neglected (with related phenomena including the Küppers-Lortz instability, domain chaos, the puzzling appearance of patterns with square symmetry, spiral defect chaos and associated dynamics of chiral symmetry breaking), passing through the consideration of finite-sized geometries and the reintroduction of the centrifugal force, up to a presentation of the myriad of possible solutions and bifurcations in cylindrical containers under the combined effects of vertical (gravity), radial (centrifugal) and azimuthal (Coriolis) forces.

Similar concepts apply to the case of convection driven by internal heating in rotating self-gravitating spherical shells (Chapter 3), whose typical manifestation under the effect of radial buoyancy is represented by an unsteady *columnar mode* able to generate differential rotation under given circumstances. Exotic modes of convection (such as hexagons, oblique rolls, hexarolls, knot convection and so

on) are also reviewed and linked to specific regions of the parameter space.

Then, attention is switched from rotating systems with bottom (or internal) heating to laterally heated configurations (temperature gradient directed horizontally, gravity directed vertically), which leads in a more or less natural way to the treatment of so-called *sloping convection* (Chapter 4), known to be the dominant mechanism producing large-scale spiralling eddy structures in Earth's atmosphere, but also eddy structures and wavy patterns in typical problems of crystal growth from the melt.

Apart from providing a general overview of so-called *quasigeostrophic theory*, Chapter 5 also gives some insights into the fundamental difference between the two main categories of fluid-dynamic instabilities in rotating fluids: one associated with problems for which the unstable modes essentially involve *mass and temperature redistribution* (e.g. Rayleigh-Bénard or Marangoni-Bénard convection considered in Chapters 2 and 7, respectively); and the other including problems such as stably stratified and unstratified shear instabilities, *barotropic and baroclinic instabilities*, which appear to be connected to the *self-excitation of waves* rather than to the direct redistribution of mass and temperature.

A number of works are reviewed, which focus on the mechanism by which mechanical and wave signals interplay to control how individual convective structures decide whether to grow, differentiate, move or die, and thereby promote pattern formation during the related process. Moreover, starting from the cardinal concept of the *Rossby wave*, some modern approaches, such as the so-called CRW (counter-propagating-Rossby-wave) perspective, an ingenious application of what has become known as 'potential vorticity thinking', are also invoked and used to

elaborate a specific mathematical formalism and some associated important microphysical reasoning.

As a natural continuation of preceding chapters, Chapter 6 develops the important topic of *geostrophic turbulence*.

The basic ideas of inertial range theory are illustrated and extended phenomenologically by incorporating ideas of vortex-vortex and vortex-strain interactions that are normally present in physical and not spectral space. Then, a critical analysis of the distinctive marks of geostrophic turbulence (and its relationship with other classical models of turbulence) is developed. The main theories for *jet formation* and stability are discussed, starting from the fundamental concept of turbulent 'decascade' of energy. Subsequent arguments deal with the role played in maintaining turbulence by baroclinic effects and/or other types of 3D instabilities and on the so-called *baroclinic life cycle*. An overview of the main characteristic wavenumbers and scales relating to distinct effects is also elaborated.

Similarities between Earth's phenomena and typical features of *outer planet* (Jupiter and Saturn) dynamics are discussed as well. After the exposition of the general theory for vortex-vortex coalescence, a similar treatment is also given for phenomena of wave-wave and wave-mean-flow interference.

The remaining chapters are entirely devoted to phenomena occurring on the lab scale, thereby allowing most of the arguments introduced in earlier chapters *to spread from their traditional heartlands of meteorology and geophysics to the industrial field* (and related applications).

Along these lines, Chapter 7 is concerned with the interplay between rotation and flows induced by surface tension gradients (more specifically, *Marangoni-Bénard convection* and so-called *hydrothermal waves*, considered as typical manifestations of surface-tension-driven flows in configurations of technological interest subjected to



temperature gradient *perpendicular or parallel*, respectively, to the liquid/gas interface).

The modification of the classical hydrothermal mechanism due to rotation, in particular, is discussed on the basis of concepts of system invariance breaking (due to rotation) and of the fundamental processes allowing waves to derive energy from the basic flow (an interpretation is given as well for still unexplained observations appeared in the literature).

Chapter 8 provides specific information on cases with important background applications in the realm of crystal growth from the melt, for example the Bridgman, floating zone and Czochralski (CZ) techniques, considering, among other things, the interesting subject of *interacting baroclinic and hydrothermal waves*, together with an exposition of the most recent theories about the origin of the so-called *spoke patterns*.

The CZ configuration is used as a classical example of situations in which fluid motion is brought about by different coexisting mechanisms: Marangoni convection, generated by the interfacial stresses due to horizontal temperature gradients along the free surface and gravitational convection driven by the volumetric buoyancy forces caused by thermally and/or solutally generated density variations in the bulk of the fluid, without forgetting the presence of phenomena of a rotational nature (baroclinic instability) and those deriving from temperature contrasts induced in the vertical direction by radiative or other (localized) effects.

The exposition of turbulence given in Chapter 6 about typical planetary dynamics is *extended* in this chapter to topics of crystal growth showing commonalities and differences due to 'contamination' exerted on the geostrophic flow by effects of surface-tension or gravitational nature (thermal plumes and jets).

Then a survey is given of very classical problems in rotating fluids which come under the general heading of *differential-rotation-driven flows*. This subject includes a variety of prototypical laboratory-scale models of industrial devices (among them: centrifugal pumps, rotating compressors, turbine disks, computer storage drives, turbo-machinery, cyclone separators, rotational viscometers, pumping of liquid metals at high melting point, cooling of rotating electrical motors, rotating heat exchangers, etc.).

Rotating magnetic fields are also considered (Chapter 9) as a potential technological means for counteracting undesired flow instabilities. Some attention is also devoted to so-called *swirling flow* and related higher modes of convection (Taylor-vortex flow, Görtler vortices, instabilities of the Bodewadt layer, etc.).

Last, but not least, a synthetic account is elaborated for flows produced by *angular vibrations* (i.e. situations in which the constant rotation rate considered in earlier chapters is replaced by an angular displacement varying sinusoidally with time with respect to an initial reference position) and rocking motions (Chapter 10), which complements, from a theoretical point of view, the analogous treatment given in Wiley's earlier book on *Thermal Convection* (2010) of purely translational vibrations, and may be of interest for researchers and scientists who are now coordinating their efforts to conceive new strategies for flow control.

# Acknowledgements

The present book should be regarded as a natural and due extension of my earlier monograph *Thermal Convection: Patterns, Evolution and Stability* (published by Wiley at the beginning of 2010) in which I presented a critical, focused and ‘comparative’ study of different types of thermal convection typically encountered in natural or technological contexts (thermogravitational, thermocapillary and thermovibrational), including the effect of magnetic fields and other means of flow control. That book attracted much attention and comments, as witnessed by the many reviews that have appeared in distinct important scientific journals (R.D. Simitev (2011) *Geophys. Astrophys. Fluid Dyn.*, **105** (1), 109–111; A. Nepomnyashchy (2011) *Eur. J. Mech.—B/Fluids*, **30** (1), 135; A. Gelfgat (2011) *Cryst. Res. Technol.*, **46** (8), 891–892; J. A. Reizes (2011) *Comput. Therm. Sci.*, **3** (4), 343–344).

The success of the 2010 book and the express requests of many referees to ‘complete’ the treatment of thermal convection, including the influence of Coriolis and centrifugal forces, as well as the development of turbulence, led me to undertake the present new work, for which I gratefully acknowledge also the many unknown reviewers selected by John Wiley & Sons, who initially examined the new book project, for their critical reading and valuable comments.

Following the same spirit of the earlier 2010 monograph, I envisaged to consider both natural and industrial processes, and develop a common framework so to promote the exchange of ideas between researchers and professionals working in distinct fields (in particular between the materials science and geophysical communities).

Along these lines, deep gratitude goes to many colleagues around the world pertaining to both such categories for generously sharing with me their precious recent experimental and numerical data (in alphabetical order): Prof. R. Bessaih, Prof. F.H. Busse, Prof. R.E. Ecke, Prof. A.Yu. Gelfgat, Prof. N. Imaishi, Prof. A. Ivanova, Prof. V. Kozlov, Dr. R.P.J. Kunnen, Prof. I. Mutabazi, Prof. P.B. Rhines, Prof. P. Read, Prof. V. Shevtsova, Prof. I. Ueno.

In particular, I wish to express my special thanks to Prof. P. Read (Atmospheric, Oceanic and Planetary Physics, Clarendon Laboratory, University of Oxford, United Kingdom) and to Prof. N. Imaishi (University of Kyushu, Institute for Materials, Chemistry and Engineering, Division of Advanced Device Materials, formerly Department of Advanced Material Study, Fukuoka, Japan), for several relevant suggestions which significantly improved both the clarity and the accuracy of some arguments in Chapters 3–4 and 7–8, respectively. In addition, there were several people (too numerous to be listed here) who were instrumental in keeping me updated on the latest advancements in several fields relevant to this book, especially the many authors who over recent years published their work in the Journal of Fluid Dynamics and Materials Processing (FDMP), for which I serve as the Editor-in-Chief. Last but not least, I am also indebted to my wife Paola, to whom this book is dedicated. Writing is a solitary activity; nevertheless, her good humour, sensitivity and vitality have made this project a pleasure, especially in the evening and during weekends. This new treatise is largely due to her optimism, encouragement, and patience.

Marcello Lappa

Author contact information:

Marcello Lappa, Via Salvator Rosa 53, San Giorgio a Cremano (Na), 80046—Italy

Email: [marlappa@unina.it](mailto:marlappa@unina.it), [lappa@thermalconvection.net](mailto:lappa@thermalconvection.net),  
[fdmp@techscience.com](mailto:fdmp@techscience.com)

Websites: [www.thermalconvection.net](http://www.thermalconvection.net),  
[www.fluidsandmaterials.com](http://www.fluidsandmaterials.com), [www.techscience.com/FDMP](http://www.techscience.com/FDMP)

# Chapter 1

## Equations, General Concepts and Nondimensional Numbers

Prior to expanding on the subject of convection in rotating fluids and related myriad manifestations, some propaedeutical concepts and accompanying fundamental mathematics must be provided to help the reader in the understanding of the descriptions and elaborations given later.

Along these lines, the goal of this introductory chapter is to stake out some common ground by providing a survey of *overarching principles, characteristic nondimensional parameters and governing equations*.

Such a theoretical framework, in its broadest sense, attempts to classify and characterize all forces potentially involved in the class of phenomena considered in the present book.

As the chapter progresses, in particular, balance equations are first introduced assuming an inertial frame of reference, hence providing the reader with fundamental information about the nature and properties of forces of nonrotational origin (Sections 1.1 and 1.5); then such equations are reformulated in a rotating coordinate system (Section 1.6) in which the so-called centrifugal and Coriolis forces emerge naturally as 'noninertial' effects.

While such a practical approach justifies the use of continuum mechanics and of macrophysical differential equations for the modelling of the underlying processes, it is insufficient, however, for the understanding/introduction of a microscopic phenomenological theory. Such development requires some microphysical reasoning. The cross-link between macro- and micro-scales is, in general, a challenging problem.

Due to page limits, here we limit ourselves to presenting the Navier–Stokes and energy equations directly in their macroscopic (continuum) form, the reader being referred to other texts (e.g. Lappa, 2010) for a complete elaboration of the approach leading from a microscopic phenomenological model to the continuum formalism.

## 1.1 The Navier-Stokes and Energy Equations

The Navier–Stokes equations, named after Claude-Louis Navier and George Gabriel Stokes (Navier, 1822; Stokes, 1845), describe the motion of a variety of fluid substances, including gases, liquids and even solids of geological sizes and time-scales. Thereby, they can be used to model flows of technological interest (too many to mention; e.g. fluid motion inside a crucible used for crystal growth or for the processing of metal alloys), but also weather, ocean currents and even motions of cosmological interest.

In their macroscopic (continuum) form these equations establish that the overall mass must be conserved and that changes in momentum can be simply expressed as the sum of dissipative viscous forces, changes in pressure, gravity, surface tension (in the presence of a free surface) and other forces (electric, magnetic, etc.) acting on the fluid.

### 1.1.1 The Continuity Equation

The mass balance equation (generally referred to in the literature as the continuity equation) reads:

$$\mathbf{1.1a} \quad \frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \underline{V}) = 0$$

that, in terms of the substantial derivative  $D/Dt = \partial/\partial t + \underline{V} \cdot \nabla$  (also known as ‘material’ or ‘total’ derivative), can be rewritten as

$$\mathbf{1.1b} \quad \frac{D\rho}{Dt} + \rho \nabla \cdot \underline{V} = 0$$

where  $\rho$  and  $\underline{V}$  are, respectively, the fluid density and velocity.

## 1.1.2 The Momentum Equation

The momentum equation reads:

$$\mathbf{1.2} \quad \frac{\partial}{\partial t} (\rho \underline{V}) + \nabla \cdot \underline{\Phi}_{mt} = \underline{F}_b$$

where  $\underline{F}_b$  is the generic body force acting on the fluid and  $\underline{\Phi}_{mt}$  the flux of momentum, which can be written as

$$\mathbf{1.3} \quad \underline{\Phi}_{mt} = \rho \underline{V} \underline{V} - \underline{\tau}$$

where  $\underline{\tau}$  is known as the *stress tensor*. Such a tensor can be regarded as a stochastic measure of the exchange of microscopic momentum induced at molecular level by particle random motion (it provides clear evidence of the fact that viscous forces originate in molecular interactions; we shall come back to this concept later).

Substituting [Equation 1.3](#) into [Equation 1.2](#), it follows that:

$$\mathbf{1.4a} \quad \frac{\partial}{\partial t} (\rho \underline{V}) + \nabla \cdot (\rho \underline{V} \underline{V} - \underline{\tau}) = \underline{F}_b$$

$$\mathbf{1.4b} \quad \rho \frac{D\underline{V}}{Dt} - \nabla \cdot \underline{\tau} = \underline{F}_b$$

## 1.1.3 The Total Energy Equation

Introducing the total energy as:

$$\mathbf{1.5} \quad E = \frac{1}{2} \rho V^2 + \rho u_{\text{int}}$$

the total energy balance equation can be cast in condensed form as:

$$\mathbf{1.6a} \quad \frac{\partial}{\partial t} \left[ \rho \left( \frac{1}{2} V^2 + u_{\text{int}} \right) \right] + \nabla \cdot \left[ \rho \left( \frac{1}{2} V^2 + u_{\text{int}} \right) \underline{V} + \underline{J}_u - \underline{V} \cdot \underline{\tau} \right] = \underline{F}_b \cdot \underline{V}$$

or in terms of the substantial derivative:

$$\mathbf{1.6b} \quad \rho \frac{D}{Dt} \left( \frac{1}{2} V^2 + u_{\text{int}} \right) + \nabla \cdot [\underline{J}_u - \underline{V} \cdot \underline{\tau}] = \underline{F}_b \cdot \underline{V}$$



where

$$\mathbf{1.7} \quad \Phi_e = \left( \frac{1}{2} \rho V^2 + \rho u_{\text{int}} \right) \underline{V} + \underline{J}_u - \underline{V} \cdot \underline{\underline{\tau}}$$

$\underline{J}_u$  being the diffusive flux of internal energy (it measures transport at the microscopic level of molecular kinetic energy due to molecular random motion).

## 1.1.4 The Budget of Internal Energy

A specific balance equation for the single internal energy can be obtained from subtracting the kinetic energy balance equation from the total energy balance equation ([Equation 1.6](#)).

Obviously, a balance equation for the pure kinetic energy can be introduced by taking the product of the momentum balance equation with  $\underline{V}$ :

$$\mathbf{1.8a} \quad \rho \frac{D}{Dt} \left( \frac{V^2}{2} \right) - (\underline{\nabla} \cdot \underline{\underline{\tau}}) \cdot \underline{V} = \underline{E}_b \cdot \underline{V}$$

this equation, using the well-known vector identity  $\underline{\nabla} \cdot (\underline{V} \cdot \underline{\underline{\tau}}) = (\underline{\nabla} \cdot \underline{\underline{\tau}}) \cdot \underline{V} + \underline{\underline{\tau}} : \underline{\nabla V}$  can be rewritten as

$$\mathbf{1.8b} \quad \rho \frac{D}{Dt} \left( \frac{V^2}{2} \right) - \underline{\nabla} \cdot (\underline{V} \cdot \underline{\underline{\tau}}) = \underline{E}_b \cdot \underline{V} - \underline{\underline{\tau}} : \underline{\nabla V}$$

from which, among other things, it is evident that the diffusive flux of kinetic energy can be simply expressed as the scalar product between  $\underline{V}$  and the stress tensor. Subtracting, as explained before, [Equation 1.8b](#) from [Equation 1.6b](#), one obtains:

$$\mathbf{1.9a} \quad \rho \frac{Du_{\text{int}}}{Dt} + \underline{\nabla} \cdot \underline{J}_u = \underline{\underline{\tau}} : \underline{\nabla V}$$

or

$$\mathbf{1.9b} \quad \frac{\partial \rho u_{\text{int}}}{\partial t} + \underline{\nabla} \cdot [\rho u_{\text{int}} \underline{V} + \underline{J}_u] = \underline{\underline{\tau}} : \underline{\nabla V}$$

that is the aforementioned balance equation for the internal energy.

## 1.1.5 Closure Models

In general, the ‘closure’ of the thermofluid–dynamic balance equations given in the preceding sections, i.e. the determination of a precise mathematical formalism relating the diffusive fluxes (stress tensor and the diffusive flux of internal energy) to the macroscopic variables involved in the process, is not as straightforward as many would assume.

For a particular but fundamental category of fluids, known as ‘newtonian fluids’ the treatment of this problem, however, becomes relatively simple.

For the case considered in the present book (nonpolar fluids and absence of torque forces), the stress tensor can be taken symmetric, i.e.  $\tau_{ij} = \tau_{ji}$  (conversely, a typical example of fluids for which the stress tensor is *not* symmetric is given by ‘micropolar fluids’, which represent fluids consisting of rigid, randomly oriented particles suspended in a viscous medium).

If the considered fluid is in quiescent conditions (i.e. there is no macroscopic motion) the stress tensor is diagonal and simply reads

$$\mathbf{1.10a} \quad \underline{\underline{\tau}} = -p\underline{\underline{I}}$$

where  $\underline{\underline{I}}$  is the unity tensor and  $p$  is the pressure.

In the presence of bulk convection the above expression must be enriched with the contributions induced by macroscopic fluid motion.

In the most general case such a contribution should be related to the gradient of velocity  $\underline{\underline{\nabla V}}$  via a tensor having tensorial order 4 (from a mathematical point of view a relationship between two tensors having order two has to be established using a four-dimension tensor). According to some simple considerations based on the isotropy of fluids (i.e. their property of not being dependent upon a specific direction) and other arguments provided over the years by various authors (Isaac Newton's landmark observations in liquids; later, the so-called Chapman–Enskog expansion elaborated for gases by Grad (1963) and Rosenau (1989)), the four-dimensional tensor

relating the stress tensor to the gradients of mass velocity simply reduces to a proportionality (scalar) constant that does not depend on such gradients:

$$\underline{\underline{\tau}} = -p\underline{\underline{I}} + 2\mu(\underline{\underline{\nabla V}})_o^s$$

where the constant of proportionality  $\mu$  is known as the dynamic viscosity (it may be regarded as a macroscopic measure of the intermolecular attraction forces) and the tensor  $(\underline{\underline{\nabla V}})_o^s$  (known as the viscous stress tensor or the dissipative part of the stress tensor) comes from the following decomposition of  $\underline{\underline{\nabla V}}$ :

$$\underline{\underline{\nabla V}} = \frac{1}{3}(\underline{\underline{\nabla}} \cdot \underline{\underline{V}})\underline{\underline{I}} + (\underline{\underline{\nabla V}})_o^s + (\underline{\underline{\nabla V}})^a$$

where

$$\underline{\underline{(\nabla V)}}_o^s = (\underline{\underline{\nabla V}})^s - \frac{1}{3}(\underline{\underline{\nabla}} \cdot \underline{\underline{V}})\underline{\underline{I}}, \quad (\underline{\underline{\nabla V}})^s = \frac{\underline{\underline{\nabla V}} + \underline{\underline{\nabla V}}^T}{2}$$

$$\underline{\underline{(\nabla V)}}^a = \frac{\underline{\underline{\nabla V}} - \underline{\underline{\nabla V}}^T}{2}$$

The three contributions in [Equation 1.11](#) are known to be responsible for volume changes, deformation and rotation, respectively, of a generic (infinitesimal) parcel of fluid (moving under the effect of the velocity field  $\underline{\underline{V}}$ ; the reader being referred to Section 1.2 for additional details about the meaning of  $(\underline{\underline{\nabla V}})^a$  and its kinship with the concept of vorticity).

Moreover, in general, the diffusive flux of internal energy can be written as (Fourier law):

$$\underline{\underline{J}}_u = -\lambda\underline{\underline{\nabla T}}$$

where  $\lambda$  is the thermal conductivity and  $T$  the fluid temperature.

Using such closure relationships, and taking into account the following vector and tensor identities:

$$\underline{\underline{\nabla}} \cdot (p\underline{\underline{I}}) = \underline{\underline{\nabla}} p$$

$$\underline{\underline{\nabla}} \cdot (p\underline{\underline{V}}) = p\underline{\underline{\nabla}} \cdot \underline{\underline{V}} + \underline{\underline{V}} \cdot \underline{\underline{\nabla}} p$$

$$p\underline{\underline{I}} : \underline{\underline{\nabla V}} = p\underline{\underline{\nabla}} \cdot \underline{\underline{V}}$$

$$\mathbf{1.17} \quad (\nabla V)_o^s : \nabla V = (\nabla V)_o^s : \left[ \frac{1}{3} (\nabla \cdot \underline{V}) \underline{I} + (\nabla V)_o^s + (\nabla V)^a \right] = (\nabla V)_o^s : (\nabla V)_o^s$$

the balance equations can be therefore rewritten in compact form as follows:

Momentum:

$$\mathbf{1.18a} \quad \frac{\partial}{\partial t} (\rho \underline{V}) + \nabla \cdot (\rho \underline{V} \underline{V}) + \nabla p = \nabla \cdot [2\mu (\nabla V)_o^s] + \underline{F}_b$$

$$\mathbf{1.18b} \quad \rho \frac{D\underline{V}}{Dt} + \nabla p = \nabla \cdot [2\mu (\nabla V)_o^s] + \underline{F}_b$$

Kinetic energy:

**1.19a**

$$\frac{\partial}{\partial t} \left( \rho \frac{V^2}{2} \right) + \nabla \cdot \left( \rho \frac{V^2}{2} \underline{V} \right) + \underline{V} \cdot \nabla p = \nabla \cdot [2\mu \underline{V} \cdot (\nabla V)_o^s] - 2\mu (\nabla V)_o^s : (\nabla V)_o^s + \underline{F}_b \cdot \underline{V}$$

$$\mathbf{1.19b} \quad \rho \frac{D}{Dt} \left( \frac{V^2}{2} \right) + \underline{V} \cdot \nabla p = \nabla \cdot [2\mu \underline{V} \cdot (\nabla V)_o^s] - 2\mu (\nabla V)_o^s : (\nabla V)_o^s + \underline{F}_b \cdot \underline{V}$$

Internal energy:

$$\mathbf{1.20a} \quad \frac{\partial \rho u_{\text{int}}}{\partial t} + \nabla \cdot [\rho u_{\text{int}} \underline{V}] = \nabla \cdot (\lambda \nabla T) - p \nabla \cdot \underline{V} + 2\mu (\nabla V)_o^s : (\nabla V)_o^s$$

$$\mathbf{1.20b} \quad \rho \frac{D u_{\text{int}}}{Dt} = \nabla \cdot (\lambda \nabla T) - p \nabla \cdot \underline{V} + 2\mu (\nabla V)_o^s : (\nabla V)_o^s$$

Total (Internal+Kinetic) energy

**1.21a**

$$\frac{\partial}{\partial t} \left[ \rho \left( \frac{1}{2} V^2 + u_{\text{int}} \right) \right] + \nabla \cdot \left[ \rho \left( \frac{1}{2} V^2 + u_{\text{int}} \right) \underline{V} \right] = \nabla \cdot (\lambda \nabla T - p \underline{V} + 2\mu \underline{V} \cdot (\nabla V)_o^s) + \underline{F}_b \cdot \underline{V}$$

$$\mathbf{1.21b} \quad \rho \frac{D}{Dt} \left( \frac{1}{2} V^2 + u_{\text{int}} \right) = \nabla \cdot (\lambda \nabla T - p \underline{V} + 2\mu \underline{V} \cdot (\nabla V)_o^s) + \underline{F}_b \cdot \underline{V}$$

## 1.2 Some Considerations about the Dynamics of Vorticity

## 1.2.1 Vorticity and Circulation

Apart from the classical fluid-dynamic variables such as mass, momentum, (kinetic, internal or total) energy, whose balance equations have been shortly presented in the preceding section, 'vorticity' should be regarded as an additional useful mathematical concept for a better characterization of certain types of flow. Generally used in synergetic combination with the other classical concepts, this quantity has been found to play a fundamental role in the physics of vortex-dominated flows, *its dynamics being the primary tool to understand the time evolution of dissipative vortical structures.*

In the following we provide some related fundamental notions, together with a short illustration of the related interdependencies with other variables, as well as a derivation of the related balance equation.

Along these lines, it is worth starting the discussion with the observation that, in general, vorticity can be related to the amount of 'circulation' or 'rotation' (or more strictly, the local angular rate of rotation) in a fluid (it is intimately linked to the moment of momentum of a generic small fluid particle about its own centre of mass). The average vorticity in a small region of fluid flow, in fact, can be defined as the circulation  $\Gamma$  around the boundary of the small region, divided by the area  $A$  of the small region.


$$\mathbf{1.22a} \quad \bar{\zeta} = \frac{\Gamma}{A}$$

where the fluid circulation  $\Gamma$  is defined as the line integral of the velocity  $\underline{V}$  around the closed curve  $\ell$  in [Figure 1.1](#).

$$\mathbf{1.22b} \quad \Gamma = \oint_{\ell} \underline{V} \cdot \hat{t} d\ell$$

$\hat{t}$  being the unit vector tangent to  $\ell$ .

**Figure 1.1** Vorticity as a measure of the rate of rotational spin in a fluid.



In practice, the vorticity at a point in a fluid can be regarded as the limit of [Equation 1.22a](#) as the area of the small region of fluid approaches zero at the point:

$$\mathbf{1.22c} \quad \zeta = \frac{d\Gamma}{dA}$$

In addition to the previous modelling, using the Stokes theorem (purely geometrical in nature), which equates the circulation  $\Gamma$  around  $\ell$  to the flux of the curl of  $\underline{V}$  through any surface area bounded by  $\ell$ :

$$\mathbf{1.23} \quad \Gamma = \oint_{\ell} \underline{V} \cdot \hat{t} dl = \int_A (\nabla \wedge \underline{V}) \cdot \hat{n} dS$$

where  $\hat{n}$  is the unit vector perpendicular to the surface  $A$  bounded by the closed curve  $\ell$  (it is implicitly assumed that  $\ell$  is smooth enough, i.e. it is locally lipschitzian; this implies that the existence of the unit vector perpendicular to the surface is guaranteed), it becomes evident from a mathematical point of view that the vorticity at a point can be defined as the curl of the velocity:

$$\mathbf{1.24} \quad \underline{\zeta} = \nabla \wedge \underline{V}$$

Therefore, it is a vector quantity, whose direction is along the axis of rotation of the fluid.

Notably,  $\underline{\zeta}$  has the same components as the anti-symmetric part of the tensor  $\nabla \underline{V}$ , that is in line with the explanation given in Section 1.1.5 about the physical meaning of  $(\nabla \underline{V})^a$ .

Related concepts are the vortex line, which is a line that is at any point tangential to the local vorticity; and a vortex tube which is the surface in the fluid formed by all vortex lines passing through a given (reducible) closed curve in the fluid.