

STUDENT SOLUTIONS  
MANUAL TO ACCOMPANY  
**LOSS MODELS**

FROM DATA TO DECISIONS

————— Fourth Edition —————

STUART A. KLUGMAN • HARRY H. PANJER • GORDON E. WILLMOT



*Student Solutions Manual to Accompany*  
**LOSS MODELS**

WILEY SERIES IN PROBABILITY AND STATISTICS

Established by WALTER A. SHEWHART and SAMUEL S. WILKS

Editors: *David J. Balding, Noel A. C. Cressie, Garrett M. Fitzmaurice,  
Harvey Goldstein, Iain M. Johnstone, Geert Molenberghs, David W. Scott,  
Adrian F. M. Smith, Ruey S. Tsay, Sanford Weisberg*

Editors Emeriti: *Vic Barnett, J. Stuart Hunter, Joseph B. Kadane, Jozef L. Teugels*

A complete list of the titles in this series appears at the end of this volume.

---

*Student Solutions Manual to Accompany*

***LOSS MODELS***

*From Data to Decisions*

*Fourth Edition*

---

**Stuart A. Klugman**

*Society of Actuaries*

**Harry H. Panjer**

*University of Waterloo*

**Gordon E. Willmot**

*University of Waterloo*

SOCIETY OF ACTUARIES

 **WILEY**

A JOHN WILEY & SONS, INC., PUBLICATION

Copyright © 2012 by John Wiley & Sons, Inc. All rights reserved

Published by John Wiley & Sons, Inc., Hoboken, New Jersey  
Published simultaneously in Canada

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4470, or on the web at [www.copyright.com](http://www.copyright.com). Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008, or online at <http://www.wiley.com/go/permission>.

**Limit of Liability/Disclaimer of Warranty:** While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services or for technical support, please contact our Customer Care Department within the United States at (800) 762-2974, outside the United States at (317) 572-3993 or fax (317) 572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic formats. For more information about Wiley products, visit our web site at [www.wiley.com](http://www.wiley.com).

***Library of Congress Cataloging-in-Publication Data is available.***

ISBN 978-1-118-31531-6

10 9 8 7 6 5 4 3 2 1

# CONTENTS

---

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Chapter 2 solutions</b>	<b>3</b>
2.1	Section 2.2	3
<b>3</b>	<b>Chapter 3 solutions</b>	<b>9</b>
3.1	Section 3.1	9
3.2	Section 3.2	14
3.3	Section 3.3	15
3.4	Section 3.4	16
3.5	Section 3.5	20

<b>4</b>	<b>Chapter 4 solutions</b>	<b>23</b>
4.1	Section 4.2	23
<b>5</b>	<b>Chapter 5 solutions</b>	<b>29</b>
5.1	Section 5.2	29
5.2	Section 5.3	40
5.3	Section 5.4	41
<b>6</b>	<b>Chapter 6 solutions</b>	<b>43</b>
6.1	Section 6.1	43
6.2	Section 6.5	43
6.3	Section 6.6	44
<b>7</b>	<b>Chapter 7 solutions</b>	<b>47</b>
7.1	Section 7.1	47
7.2	Section 7.2	48
7.3	Section 7.3	50
7.4	Section 7.5	52
<b>8</b>	<b>Chapter 8 solutions</b>	<b>55</b>
8.1	Section 8.2	55
8.2	Section 8.3	57
8.3	Section 8.4	59
8.4	Section 8.5	60
8.5	Section 8.6	64
<b>9</b>	<b>Chapter 9 solutions</b>	<b>67</b>
9.1	Section 9.1	67
9.2	Section 9.2	68
9.3	Section 9.3	68
9.4	Section 9.4	78
9.5	Section 9.6	79
9.6	Section 9.7	85
9.7	Section 9.8	87



<b>10</b>	<b>Chapter 10 solutions</b>	<b>93</b>
	10.1 Section 10.2	93
	10.2 Section 10.3	96
	10.3 Section 10.4	97
<b>11</b>	<b>Chapter 11 solutions</b>	<b>99</b>
	11.1 Section 11.2	99
	11.2 Section 11.3	100
<b>12</b>	<b>Chapter 12 solutions</b>	<b>105</b>
	12.1 Section 12.1	105
	12.2 Section 12.2	111
	12.3 Section 12.3	114
	12.4 Section 12.4	116
<b>13</b>	<b>Chapter 13 solutions</b>	<b>119</b>
	13.1 Section 13.1	119
	13.2 Section 13.2	124
	13.3 Section 13.3	137
	13.4 Section 13.4	145
	13.5 Section 13.5	146
<b>14</b>	<b>Chapter 14</b>	<b>147</b>
	14.1 Section 14.7	147
<b>15</b>	<b>Chapter 15</b>	<b>151</b>
	15.1 Section 15.2	151
	15.2 Section 15.3	161
<b>16</b>	<b>Chapter 16 solutions</b>	<b>165</b>
	16.1 Section 16.3	165
	16.2 Section 16.4	166
	16.3 Section 16.5	175
<b>17</b>	<b>Chapter 17 solutions</b>	<b>183</b>
	17.1 Section 17.7	183

<b>18</b>	<b>Chapter 18</b>	<b>187</b>
18.1	Section 18.9	187
<b>19</b>	<b>Chapter 19</b>	<b>219</b>
19.1	Section 19.5	219
<b>20</b>	<b>Chapter 20 solutions</b>	<b>227</b>
20.1	Section 20.1	227
20.2	Section 20.2	228
20.3	Section 20.3	229
20.4	Section 20.4	230

# CHAPTER 1

---

## INTRODUCTION

---

The solutions presented in this manual reflect the authors' best attempt to provide insights and answers. While we have done our best to be complete and accurate, errors may occur and there may be more elegant solutions. Errata will be posted at the ftp site dedicated to the text and solutions manual:

[ftp://ftp.wiley.com/public/sci\\_tech\\_med/loss\\_models/](ftp://ftp.wiley.com/public/sci_tech_med/loss_models/)

Should you find errors or would like to provide improved solutions, please send your comments to Stuart Klugman at [sklugman@soa.org](mailto:sklugman@soa.org).



## CHAPTER 2

---

## CHAPTER 2 SOLUTIONS

---


### 2.1 SECTION 2.2

$$2.1 \quad F_5(x) = 1 - S_5(x) = \begin{cases} 0.01x, & 0 \leq x < 50, \\ 0.02x - 0.5, & 50 \leq x < 75. \end{cases}$$


$$f_5(x) = F_5'(x) = \begin{cases} 0.01, & 0 < x < 50, \\ 0.02, & 50 \leq x < 75. \end{cases}$$

$$h_5(x) = \frac{f_5(x)}{S_5(x)} = \begin{cases} \frac{1}{100-x}, & 0 < x < 50, \\ \frac{1}{75-x}, & 50 \leq x < 75. \end{cases}$$


**2.2** The requested plots follow. The triangular spike at zero in the density function for Model 4 indicates the 0.7 of discrete probability at zero.




Distribution function for Model 3.




Distribution function for Model 4.




Distribution function for Model 5.




Probability function for Model 3.




Density function for Model 4.



Density function for Model 5.



Hazard rate for Model 4.



Hazard rate for Model 5.

**2.3**  $f'(x) = 4(1+x^2)^{-3} - 24x^2(1+x^2)^{-4}$ . Setting the derivative equal to zero and multiplying by  $(1+x^2)^4$  give the equation  $4(1+x^2) - 24x^2 = 0$ . This is equivalent to  $x^2 = 1/5$ . The only positive solution is the mode of  $1/\sqrt{5}$ .

**2.4** The survival function can be recovered as

$$\begin{aligned} 0.5 &= S(0.4) = e^{-\int_0^{0.4} A + e^{2x} dx} \\ &= e^{-Ax - 0.5e^{2x}} \Big|_0^{0.4} \\ &= e^{-0.4A - 0.5e^{0.8} + 0.5} \end{aligned}$$

Taking logarithms gives

$$-0.693147 = -0.4A - 1.112770 + 0.5,$$

and thus  $A = 0.2009$ .



**2.5** The ratio is

$$\begin{aligned}r &= \frac{\left(\frac{10,000}{10,000+d}\right)^2}{\left(\frac{20,000}{20,000+d^2}\right)^2} \\ &= \left(\frac{20,000+d^2}{20,000+2d}\right)^2 \\ &= \frac{20,000^2 + 40,000d^2 + d^4}{20,000^2 + 80,000d + 4d^2}.\end{aligned}$$

From observation or two applications of L'Hôpital's rule, we see that the limit is infinity.



# CHAPTER 3

---

## CHAPTER 3 SOLUTIONS

---

### 3.1 SECTION 3.1

$$\begin{aligned} 3.1 \quad \mu_3 &= \int_{-\infty}^{\infty} (x - \mu)^3 f(x) dx = \int_{-\infty}^{\infty} (x^3 - 3x^2\mu + 3x\mu^2 - \mu^3) f(x) dx \\ &= \mu'_3 - 3\mu'_2\mu + 2\mu^3, \end{aligned}$$

$$\begin{aligned} \mu_4 &= \int_{-\infty}^{\infty} (x - \mu)^4 f(x) dx \\ &= \int_{-\infty}^{\infty} (x^4 - 4x^3\mu + 6x^2\mu^2 - 4x\mu^3 + \mu^4) f(x) dx \\ &= \mu'_4 - 4\mu'_3\mu + 6\mu'_2\mu^2 - 3\mu^4. \end{aligned}$$

**3.2** For Model 1,  $\sigma^2 = 3,333.33 - 50^2 = 833.33$ ,  $\sigma = 28.8675$ .

$$\mu'_3 = \int_0^{100} x^3(0.01) dx = 250,000, \quad \mu_3 = 0, \quad \gamma_1 = 0.$$

$$\mu'_4 = \int_0^{100} x^4(0.01) dx = 20,000,000, \quad \mu_4 = 1,250,000, \quad \gamma_2 = 1.8.$$

For Model 2,  $\sigma^2 = 4,000,000 - 1,000^2 = 3,000,000$ ,  $\sigma = 1,732.05$ .  $\mu'_3$  and  $\mu'_4$  are both infinite so the skewness and kurtosis are not defined.

For Model 3,  $\sigma^2 = 2.25 - .93^2 = 1.3851$ ,  $\sigma = 1.1769$ .  
 $\mu'_3 = 0(0.5) + 1(0.25) + 8(0.12) + 27(0.08) + 64(0.05) = 6.57$ ,  $\mu_3 = 1.9012$ ,  
 $\gamma_1 = 1.1663$ ,  $\mu'_4 = 0(0.5) + 1(0.25) + 16(0.12) + 81(0.08) + 256(0.05) = 21.45$ ,  
 $\mu_4 = 6.4416$ ,  $\gamma_2 = 3.3576$ .

For Model 4,  $\sigma^2 = 6,000,000,000 - 30,000^2 = 5,100,000,000$ ,  $\sigma = 71,414$ .  
 $\mu'_3 = 0^3(0.7) + \int_0^\infty x^3(0.000003)e^{-.00001x}dx = 1.8 \times 10^{15}$ ,  
 $\mu_3 = 1.314 \times 10^{15}$ ,  $\gamma_1 = 3.6078$ .  
 $\mu'_4 = \int_0^\infty x^4(0.000003)e^{-.00001x}dx = 7.2 \times 10^{20}$ ,  $\mu_4 = 5.3397 \times 10^{20}$ ,  
 $\gamma_2 = 20.5294$ .

For Model 5,  $\sigma^2 = 2,395.83 - 43.75^2 = 481.77$ ,  $\sigma = 21.95$ .  
 $\mu'_3 = \int_0^{50} x^3(0.01)dx + \int_{50}^{75} x^3(0.02)dx = 142,578.125$ ,  $\mu_3 = -4,394.53$ ,  
 $\gamma_1 = -0.4156$ .  
 $\mu'_4 = \int_0^{50} x^4(0.01)dx + \int_{50}^{75} x^4(0.02)dx = 8,867,187.5$ ,  $\mu_4 = 439,758.30$ ,  
 $\gamma_2 = 1.8947$ .

**3.3** The standard deviation is the mean times the coefficient of variation, or 4, and so the variance is 16. From (3.3) the second raw moment is  $16 + 2^2 = 20$ . The third central moment is (using Exercise 3.1)  $136 - 3(20)(2) + 2(2)^3 = 32$ . The skewness is the third central moment divided by the cube of the standard deviation, or  $32/4^3 = 1/2$ .

**3.4** For a gamma distribution the mean is  $\alpha\theta$ . The second raw moment is  $\alpha(\alpha + 1)\theta^2$ , and so the variance is  $\alpha\theta^2$ . The coefficient of variation is  $\sqrt{\alpha\theta^2}/\alpha\theta = \alpha^{-1/2} = 1$ . Therefore  $\alpha = 1$ . The third raw moment is  $\alpha(\alpha + 1)(\alpha + 2)\theta^3 = 6\theta^3$ . From Exercise 3.1, the third central moment is  $6\theta^3 - 3(2\theta^2)\theta + 2\theta^3 = 2\theta^3$  and the skewness is  $2\theta^3/(\theta^2)^{3/2} = 2$ .

**3.5** For Model 1,

$$e(d) = \frac{\int_d^{100} (1 - 0.01x)dx}{1 - 0.01d} = \frac{100 - d}{2}.$$

For Model 2,

$$e(d) = \frac{\int_d^\infty \left(\frac{2,000}{x + 2,000}\right)^3 dx}{\left(\frac{2,000}{d + 2,000}\right)^3} = \frac{2,000 + d}{2}.$$

For Model 3,

$$e(d) = \begin{cases} \frac{0.25(1-d) + 0.12(2-d) + 0.08(3-d) + 0.05(4-d)}{0.5} = 1.86 - d, & 0 \leq d < 1, \\ \frac{0.12(2-d) + 0.08(3-d) + 0.05(4-d)}{0.25} = 2.72 - d, & 1 \leq d < 2, \\ \frac{0.08(3-d) + 0.05(4-d)}{0.13} = 3.3846 - d, & 2 \leq d < 3, \\ \frac{0.05(4-d)}{0.05} = 4 - d, & 3 \leq d < 4. \end{cases}$$

For Model 4,

$$e(d) = \frac{\int_d^\infty 0.3e^{-0.00001x} dx}{0.3e^{-0.00001d}} = 100,000.$$

The functions are straight lines for Models 1, 2, and 4. Model 1 has negative slope, Model 2 has positive slope, and Model 4 is horizontal.

**3.6** For a uniform distribution on the interval from 0 to  $w$ , the density function is  $f(x) = 1/w$ . The mean residual life is

$$\begin{aligned} e(d) &= \frac{\int_d^w (x-d)w^{-1} dx}{\int_d^w w^{-1} dx} \\ &= \frac{\left. \frac{(x-d)^2}{2w} \right|_d^w}{\frac{w-d}{w}} \\ &= \frac{(w-d)^2}{2(w-d)} \\ &= \frac{w-d}{2}. \end{aligned}$$

The equation becomes

$$\frac{w-30}{2} = \frac{100-30}{2} + 4,$$

with a solution of  $w = 108$ .

**3.7** From the definition,

$$e(\lambda) = \frac{\int_\lambda^\infty (x-\lambda)\lambda^{-1}e^{-x/\lambda} dx}{\int_\lambda^\infty \lambda^{-1}e^{-x/\lambda} dx} = \lambda.$$

$$\begin{aligned}
 \mathbf{3.8} \quad E(X) &= \int_0^\infty xf(x)dx = \int_0^d xf(x)dx + \int_d^\infty df(x)dx + \int_d^\infty (x-d)f(x)dx \\
 &= \int_0^d xf(x)dx + d[1-F(d)] + e(d)S(d) = E[X \wedge d] + e(d)S(d).
 \end{aligned}$$

**3.9** For Model 1, from (3.8),

$$E[X \wedge u] = \int_0^u x(0.01)dx + u(1 - 0.01u) = u(1 - 0.005u)$$

and from (3.10),

$$E[X \wedge u] = 50 - \frac{100-u}{2}(1 - 0.01u) = u(1 - 0.005u).$$

From (3.9),

$$E[X \wedge u] = - \int_{-\infty}^0 0 dx + \int_0^u 1 - 0.01x dx = u - \frac{0.01u^2}{2} = u(1 - 0.005u).$$

For Model 2, from (3.8),

$$E[X \wedge u] = \int_0^u x \frac{3(2,000)^3}{(x+2,000)^4} dx + u \frac{2,000^3}{(2,000+u)^3} = 1000 \left[ 1 - \frac{4,000,000}{(2,000+u)^2} \right],$$

and from (3.10),

$$E[X \wedge u] = 1,000 - \frac{2,000+u}{2} \left( \frac{2,000}{2,000+u} \right)^3 = 1,000 \left[ 1 - \frac{4,000,000}{(2,000+u)^2} \right].$$

From (3.9),

$$\begin{aligned}
 E[X \wedge u] &= \int_0^u \left( \frac{2,000}{2,000+x} \right)^3 dx = \frac{-2,000^3}{2(2,000+x)^2} \Big|_0^u \\
 &= 1,000 \left[ 1 - \frac{4,000,000}{(2,000+u)^2} \right].
 \end{aligned}$$

For Model 3, from (3.8),

$$E[X \wedge u] = \begin{cases} 0(0.5) + u(0.5) = 0.5u, & 0 \leq u < 1, \\ 0(0.5) + 1(0.25) + u(0.25) = 0.25 + 0.25u, & 1 \leq u < 2, \\ 0(0.5) + 1(0.25) + 2(0.12) + u(0.13) \\ = 0.49 + 0.13u, & 2 \leq u < 3, \\ 0(0.5) + 1(0.25) + 2(0.12) + 3(0.08) + u(0.05) \\ = 0.73 + 0.05u, & 3 \leq u < 4, \end{cases}$$

and from (3.10),

$$E[X \wedge u] = \begin{cases} 0.93 - (1.86 - u)(0.5) = 0.5u, & 0 \leq u < 1, \\ 0.93 - (2.72 - u)(0.25) = 0.25 + 0.25u, & 1 \leq u < 2, \\ 0.93 - (3.3846 - u)(0.13) = 0.49 + 0.13u, & 2 \leq u < 3, \\ 0.93 - (4 - u)0(.05) = 0.73 + 0.05u, & 3 \leq u < 4. \end{cases}$$

For Model 4, from (3.8),

$$\begin{aligned} E[X \wedge u] &= \int_0^u x(0.000003)e^{-0.00001x} dx + u(0.3)e^{-0.00001u} \\ &= 30,000[1 - e^{-0.00001u}], \end{aligned}$$

and from (3.10),

$$E[X \wedge u] = 30,000 - 100,000(0.3e^{-0.00001u}) = 30,000[1 - e^{-0.00001u}].$$

**3.10** For a discrete distribution (which all empirical distributions are), the mean residual life function is

$$e(d) = \frac{\sum_{x_j > d} (x_j - d)p(x_j)}{\sum_{x_j > d} p(x_j)}.$$

When  $d$  is equal to a possible value of  $X$ , the function cannot be continuous because there is jump in the denominator but not in the numerator. For an exponential distribution, argue as in Exercise 3.7 to see that it is constant. For the Pareto distribution,

$$\begin{aligned} e(d) &= \frac{E(X) - E(X \wedge d)}{S(d)} \\ &= \frac{\frac{\theta}{\alpha-1} - \frac{\theta}{\alpha-1} \left[ 1 - \left( \frac{\theta}{\theta+d} \right)^{\alpha-1} \right]}{\left( \frac{\theta}{\theta+d} \right)^{\alpha}} \\ &= \frac{\theta}{\alpha-1} \frac{\theta+d}{\theta} = \frac{\theta+d}{\alpha-1}, \end{aligned}$$

which is increasing in  $d$ . Only the second statement is true.

**3.11** Applying the formula from the solution to Exercise 3.10 gives

$$\frac{10,000 + 10,000}{0.5 - 1} = -40,000,$$

which cannot be correct. Recall that the numerator of the mean residual life is  $E(X) - E(X \wedge d)$ . However, when  $\alpha \leq 1$ , the expected value is infinite and so is the mean residual life.

**3.12** The right truncated variable is defined as  $Y = X$  given that  $X \leq u$ . When  $X > u$ , this variable is not defined. The  $k$ th moment is

$$E(Y^k) = \frac{\int_0^u x^k f(x) dx}{F(u)} = \frac{\sum_{x_i \leq u} x_i^k p(x_i)}{F(u)}.$$

**3.13** This is a single parameter Pareto distribution with parameters  $\alpha = 2.5$  and  $\theta = 1$ . The moments are  $\mu_1 = 2.5/1.5 = 5/3$  and  $\mu_2 = 2.5/.5 - (5/3)^2 = 20/9$ . The coefficient of variation is  $\sqrt{20/9}/(5/3) = 0.89443$ .

**3.14**  $\mu = 0.05(100) + 0.2(200) + 0.5(300) + 0.2(400) + 0.05(500) = 300$ .  
 $\sigma^2 = 0.05(-200)^2 + 0.2(-100)^2 + 0.5(0)^2 + 0.2(100)^2 + 0.05(200)^2 = 8,000$ .  
 $\mu_3 = 0.05(-200)^3 + 0.2(-100)^3 + 0.5(0)^3 + 0.2(100)^3 + 0.05(200)^3 = 0$ .  
 $\mu_4 = 0.05(-200)^4 + 0.2(-100)^4 + 0.5(0)^4 + 0.2(100)^4 + 0.05(200)^4 = 200,000,000$ .  
 Skewness is  $\gamma_1 = \mu_3/\sigma^3 = 0$ . Kurtosis is  $\gamma_2 = \mu_4/\sigma^4 = 200,000,000/8,000^2 = 3.125$ .

**3.15** The Pareto mean residual life function is

$$e_X(d) = \frac{\int_d^\infty \theta^\alpha (x + \theta)^{-\alpha} dx}{\theta^\alpha (x + d)^{-\alpha}} = (d + \theta)/(\alpha - 1),$$

and so  $e_X(2\theta)/e_X(\theta) = (2\theta + \theta)/(\theta + \theta) = 1.5$ .

**3.16** Sample mean:  $0.2(400) + 0.7(800) + 0.1(1,600) = 800$ . Sample variance:  $0.2(-400)^2 + 0.7(0)^2 + 0.1(800)^2 = 96,000$ . Sample third central moment:  $0.2(-400)^3 + 0.7(0)^3 + 0.1(800)^3 = 38,400,000$ . Skewness coefficient:  $38,400,000/96,000^{1.5} = 1.29$ .

## 3.2 SECTION 3.2

**3.17** The pdf is  $f(x) = 2x^{-3}$ ,  $x \geq 1$ . The mean is  $\int_1^\infty 2x^{-2} dx = 2$ . The median is the solution to  $.5 = F(x) = 1 - x^{-2}$ , which is 1.4142. The mode is the value where the pdf is highest. Because the pdf is strictly decreasing, the mode is at its smallest value, 1.

**3.18** For Model 2, solve  $p = 1 - \left(\frac{2,000}{2,000 + \pi_p}\right)^3$  and so  $\pi_p = 2,000[(1-p)^{-1/3} - 1]$  and the requested percentiles are 519.84 and 1419.95.



For Model 4, the distribution function jumps from 0 to 0.7 at zero and so  $\pi_{0.5} = 0$ . For percentile above 70, solve  $p = 1 - 0.3e^{-0.00001\pi_p}$ , and so  $\pi_p = -100,000 \ln[(1 - p)/0.3]$  and  $\pi_{0.8} = 40,546.51$ .

For Model 5, the distribution function has two specifications. From  $x = 0$  to  $x = 50$  it rises from 0.0 to 0.5, and so for percentiles at 50 or below, the equation to solve is  $p = 0.01\pi_p$  for  $\pi_p = 100p$ . For  $50 < x \leq 75$ , the distribution function rises from 0.5 to 1.0, and so for percentiles from 50 to 100 the equation to solve is  $p = 0.02\pi_p - 0.5$  for  $\pi_p = 50p + 25$ . The requested percentiles are 50 and 65.

**3.19** The two percentiles imply

$$\begin{aligned} 0.1 &= 1 - \left( \frac{\theta}{\theta + \theta - k} \right)^\alpha, \\ 0.9 &= 1 - \left( \frac{\theta}{\theta + 5\theta - 3k} \right)^\alpha. \end{aligned}$$

Rearranging the equations and taking their ratio yield

$$\frac{0.9}{0.1} = \left( \frac{6\theta - 3k}{2\theta - k} \right)^\alpha = 3^\alpha.$$

Taking logarithms of both sides gives  $\ln 9 = \alpha \ln 3$  for  $\alpha = \ln 9 / \ln 3 = 2$ .

**3.20** The two percentiles imply

$$\begin{aligned} 0.25 &= 1 - e^{-(1,000/\theta)^\tau}, \\ 0.75 &= 1 - e^{-(100,000/\theta)^\tau}. \end{aligned}$$

Subtracting and then taking logarithms of both sides give

$$\begin{aligned} \ln 0.75 &= -(1,000/\theta)^\tau, \\ \ln 0.25 &= -(100,000/\theta)^\tau. \end{aligned}$$

Dividing the second equation by the first gives

$$\frac{\ln 0.25}{\ln 0.75} = 100^\tau.$$

Finally, taking logarithms of both sides gives  $\tau \ln 100 = \ln[\ln 0.25 / \ln 0.75]$  for  $\tau = 0.3415$ .

### 3.3 SECTION 3.3

**3.21** The sum has a gamma distribution with parameters  $\alpha = 16$  and  $\theta = 250$ . Then,  $\Pr(S_{16} > 6,000) = 1 - \Gamma(16; 6,000/250) = 1 - \Gamma(16; 24)$ . From the

Central Limit Theorem, the sum has an approximate normal distribution with mean  $\alpha\theta = 4,000$  and variance  $\alpha\theta^2 = 1,000,000$  for a standard deviation of 1000. The probability of exceeding 6,000 is  $1 - \Phi[(6,000 - 4,000)/1,000] = 1 - \Phi(2) = 0.0228$ .

**3.22** A single claim has mean  $8,000/(5/3) = 4,800$  and variance

$$2(8,000)^2/[(5/3)(2/3)] - 4,800^2 = 92,160,000.$$

The sum of 100 claims has mean 480,000 and variance 9,216,000,000, which is a standard deviation of 96,000. The probability of exceeding 600,000 is approximately

$$1 - \Phi[(600,000 - 480,000)/96,000] = 1 - \Phi(1.25) = 0.106.$$

**3.23** The mean of the gamma distribution is  $5(1,000) = 5,000$  and the variance is  $5(1,000)^2 = 5,000,000$ . For 100 independent claims, the mean is 500,000 and the variance is 500,000,000 for a standard deviation of 22,360.68. The probability of total claims exceeding 525,000 is

$$1 - \Phi[(525,000 - 500,000)/22,360.68] = 1 - \Phi(1.118) = 0.13178.$$

**3.24** The sum of 2,500 contracts has an approximate normal distribution with mean  $2,500(1,300) = 3,250,000$  and standard deviation  $\sqrt{2,500}(400) = 20,000$ . The answer is  $\Pr(X > 3,282,500) \doteq \Pr[Z > (3,282,500 - 3,250,000)/20,000] = \Pr(Z > 1.625) = 0.052$ .

### 3.4 SECTION 3.4

**3.25** While the Weibull distribution has all positive moments, for the inverse Weibull moments exist only for  $k < \tau$ . Thus by this criterion, the inverse Weibull distribution has a heavier tail. With regard to the ratio of density functions, it is (with the inverse Weibull in the numerator and marking its parameters with asterisks)

$$\frac{\tau^* \theta^{*\tau^*} x^{-\tau^*-1} e^{-(\theta^*/x)^{\tau^*}}}{\tau \theta^{-\tau} x^{\tau-1} e^{-(x/\theta)^\tau}} \propto x^{-\tau-\tau^*} e^{-(\theta^*/x)^{\tau^*} + (x/\theta)^\tau}.$$

The logarithm is

$$(x/\theta)^\tau - (\theta^*/x)^{\tau^*} - (\tau + \tau^*) \ln x.$$

The middle term goes to zero, so the issue is the limit of  $(x/\theta)^\tau - (\tau + \tau^*) \ln x$ , which is clearly infinite. With regard to the hazard rate, for the Weibull distribution we have

$$h(x) = \frac{\tau x^{\tau-1} \theta^{-\tau} e^{-(x/\theta)^\tau}}{e^{-(x/\theta)^\tau}} = \tau x^{\tau-1} \theta^{-\tau},$$