

The Stress Response of Critical Illness: Metabolic and Hormonal Aspects

Jean-Charles Preiser
Editor

 Springer

The Stress Response of Critical Illness: Metabolic and Hormonal Aspects

Jean-Charles Preiser
Editor

The Stress Response of Critical Illness: Metabolic and Hormonal Aspects

 Springer

Editor
Jean-Charles Preiser
Brussels
Belgium

ISBN 978-3-319-27685-4 ISBN 978-3-319-27687-8 (eBook)
DOI 10.1007/978-3-319-27687-8

Library of Congress Control Number: 2016935706

Springer Cham Heidelberg New York Dordrecht London
© Springer International Publishing Switzerland 2016

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

Springer International Publishing AG Switzerland is part of Springer Science+Business Media
(www.springer.com)

Foreword

Life is like riding a bicycle. To keep your balance, you must keep moving.

Albert Einstein

All living organisms share common biological characteristics, developed over hundreds of millions of years, which have allowed them to adapt and survive since the beginning of life on earth. This is particularly true for human beings, so weak and frail before the occurrence of organized civilizations. The importance of effective adaptation abilities was recognized very long ago by the Chinese and Greek physicians, although it was not before the nineteenth and twentieth centuries that the real importance of the stress response with its complex multisystem mechanisms was discovered. Claude Bernard recognized the importance of a constant *milieu intérieur*, insuring the function of body cells in a changing environment via adaptive mechanisms in the vital organs. Later, Walther Cannon developed further the concept of *homeostasis*, leading in case of failure of the homeostatic mechanisms to disequilibrium and illness.

In 1936, Hans Selye published the historical Letter to the Editor of Nature, “A syndrome produced by diverse nocuous agents,” describing stress as the consequence of an inadequate response to harmful physical and psychological agents. The stress response was originally believed to be mainly related to the neuroendocrine system activation, but Selye later realized that nearly all systems were involved and the concept of a multisystem general response was developed.

Although the clinical relevance of such adaptive mechanisms in trauma and acute conditions was long recognized, their long-term effects on the mood and behavior were not acknowledged before the second part of the twentieth century: the post-traumatic stress disorder was only included in 1980 in the third edition of the DSM of the American Psychiatric Association. This underlines the central role of the brain in the response to stress, as a regulation but also as a target organ, when its vulnerability to emotional and psychological challenges, exceeds its resilience capacity and induces undesirable emotional and behavioral symptoms.

In the field of stress response, the critical care environment is unique, since it gathers individuals with different kinds of stress: patients with life-threatening conditions, health-care professionals, mainly physicians and nurses, as well as family members. In the early phase of critical illness, the patients are submitted to acute challenges such as hemorrhage, ischemia, hypoxia, sepsis and pain, as well as psychological or emotional threats. The initial adaptation to the critical illness is mainly related to the multiple autonomic, endocrine, tissue, and immune mediators; it promotes survival and recovery. However, when the critical illness is prolonged, inadequate regulation of this response may occur, inducing damaging effects, such as depressed immunity, metabolic dysfunctions, and malnutrition. At longer term, patients with prolonged or complicated stay are at risk to develop post-traumatic stress disorder, mainly characterized by psychological, emotional, and behavioral symptoms.

The family is submitted to intense psycho-emotional stress, leading to adverse psychological responses, the so-called post-intensive care syndrome-family. The latter is mainly characterized by insomnia, anxiety, depressive symptoms, inability to perform the grief, and difficulty or inability to work. Surprisingly, the clinical importance of post-intensive care syndrome-family was only discovered about 20 year ago, despite quite a high prevalence: about a third of family members are affected by PTSD symptoms in both pediatric and adult ICUs. The process of decision making involving families and proxies plays a critical role, particularly the methods of communication and inclusion. Various strategies of communication with family members have shown to be associated with decreased anxiety, improved resilience, and coping.

In addition to emotional, psychological stress, the health-care givers are submitted to work-related stress, promoting burnout symptoms, as emotional and physical exhaustion, inability to work, depersonalization, and depression. The prevalence of PTSD in ICU nurses is particularly high, due to the daily contact with suffering, uncertainty of therapy, and death. This is also the case of health-care professionals working in emergency and mental health care, which in addition are often submitted to violence and physical assaults. The occurrence and severity of work-related stress is affected by several factors, the type, nature, and severity of the stressor, the presence of a team support, and the quality of professional training, as well as by individual factors, such as the individual personality, mental health, and social-family support.

The publication of *The Stress Response of Critical Illness: Metabolic and Hormonal Aspects* by Jean-Charles Preiser and more than 20 top-level scientists must be highlighted, since it constitutes a remarkable high level and original contribution. This book comes at a right stage, as a large body of recent information has brought new insights on the metabolic and endocrine aspects of the stress response during the last decade, such as the corticoadrenal response in sepsis, the regulation of blood glucose, and substrate metabolism in the settings of critical care.

Contents

1	Introduction	1
	Jean-Charles Preiser	
Part I Metabolic Changes		
2	Successive Phases of the Metabolic Response to Stress	5
	Jean-Charles Preiser, Carole Ichai, and A.B. Johan Groeneveld	
3	Bioenergetics of the Stress Response	19
	Christophe Faisy	
4	Mitochondrial Adaptation and Hibernation	27
	Jonathan Grip, Nicolas Tardif, and Olav Rooyackers	
5	Anabolic Resistance	45
	Jean-Paul Thissen	
6	Use of Lipids as Energy Substrates	61
	Philip C. Calder and Pierre Singer	
7	The Stress Response of Critical Illness: Metabolic and Hormonal Aspects	75
	Luc Tappy	
8	Stress Hyperglycemia	89
	Jean-Charles Preiser, Aurélie Thooft, and Rafael Machado Tironi	
9	Protein Metabolism	95
	Åke Norberg, Felix Liebau, and Jan Wernerman	
10	Micronutrients	107
	Mette M. Berger	

Part II Hormonal Regulation

- 11 Thyroidal Changes During Critical Illness** 125
Lies Langouche and Greet Van den Berghe
- 12 Stress Response: Adrenal Function** 137
Paul E. Marik
- 13 Enterohormones and the Response to Critical Illness** 153
Mark P. Plummer, Annika Reintam Blaser,
and Adam M. Deane
- 14 Adipokines in Critical Illness** 169
Katherine Robinson, John Prins, and Bala Venkatesh

Part III Particular Clinical Situations

- 15 Severe Undernutrition** 187
Pierre Singer and Jonathan Cohen
- 16 The Stress Response after Traumatic Brain Injury:
Metabolic and Hormonal Aspects** 197
Hervé Quintard, Carole Ichai, and Jean-Francois Payen
- 17 Sepsis and Multiple Organ Failure** 207
Jean-Charles Preiser and Vincent Fraipont
- 18 The Stress Response of Critical Illness: Metabolic
and Hormonal Aspects, Hormonal Regulation,
Particular Clinical Situations “Morbidity Obesity”** 217
Moise Coeffier and Fabienne Tamion
- 19 Hypermetabolic Response to Burn Injury** 227
Abdikarim Abdullahi, David Patsouris, Sheila R. Costford,
and Marc G. Jeschke

Chapter 1

Introduction

Jean-Charles Preiser

The important thing in science is not so much to obtain new facts as to discover new ways of thinking about them. William Henry Bragg

One of the most challenging tasks on earth is to bring people together and to build bridges to promote the cross-fertilization of knowledge!

In the field of care of the critically ill, this statement can be translated into the integration of new discoveries or major advances in the understanding of physiology into unbiased evaluations of new therapeutic strategies adapted from novel experimental findings. Conversely, basic science should be open to clinical data and able to understand and to integrate clinical concerns into research questions. This book aims to bridge the new knowledge gathered in experimental research with new clinical results.

Historically, the “*stress response*” was quoted by the Canadian physiologist Hans Selye, who discovered the mechanisms of the “fly or fight response” designed to restore the homeostasis needed for an independent life, as described by Claude Bernard. Some of the basic and adaptive mechanisms were preserved over the evolution, in keeping with the Darwinian theory of evolution.

Now, the changes and improvements in the practice of medicine allow patients to survive critical illness, allow surgeons to perform risky interventions, allow anesthesiologists to sedate very weak patients, etc. The support of the vital functions prolongs the lifespan of a critically ill thanks to improvements in pharmacological agents, in the technology of ventilators, renal replacement therapies, and extracorporeal membrane oxygenators. However, the metabolic and functional consequences of the critical illness can last weeks or months, representing a major burden for the society and cannot be supported by any dedicated device. Only a few drugs

J.-C. Preiser, MD, PhD

Department of Intensive Care, Erasme University Hospital, Université Libre de Bruxelles,
808 route de Lennik, Brussels 1070, Belgium

e-mail: Jean-Charles.Preiser@erasme.ulb.ac.be

© Springer International Publishing Switzerland 2016

J.-C. Preiser (ed.), *The Stress Response of Critical Illness:*

Metabolic and Hormonal Aspects, DOI 10.1007/978-3-319-27687-8_1

and nutritional formulas can slightly influence metabolism. Therefore, a detailed knowledge of the intimate mechanisms of the stress response is warranted to help clinicians to support the adaptive and desirable metabolic responses while trying to minimize the maladaptive ones.

In this book, several world-leading experts accepted to share their knowledge and to summarize the current understanding of the metabolic response to stress, including the cellular and subcellular mechanisms, the use of macro- and micronutrient as energy substrates during catabolism or anabolic resistance typically associated with critical illness. The roles and patterns of endocrine mediators will be discussed in detail. The final section will address difficult clinical situations, as examples of how the new findings can be translated into daily practice.

Specifically, in the first section, the bioenergetics of the stress response has been revisited in detail by C Faisy, using the concept of stress as a challenge to the equilibrium at each level of the body. The successive phases of the metabolic response to stress were being reviewed in a temporal and clinically relevant sequence. Novel insights as hibernation and mitochondrial mechanisms of adaptation have been updated by J Grip, N Tardif, and O Rooyackers, while the development of anabolic resistance has been reviewed by JP Thissen. The ensuing alterations in the use of lipids, carbohydrates, and protein metabolism were updated by Ph Calder, P Singer, L Tappy, A Norberg, F Liebau, and J Wernerman, whereas the current roles of micronutrients have been reviewed by MM Berger. A Thoof, R Machado, and myself summarized the related issue of stress hyperglycemia.

In Part 2, the current understanding of the functional changes of hormonal systems has been described by L Langouche and G Van den Berghe for the thyroid axis and P Marik for the adrenal system. The roles and relevance of new important endocrine mediators, the enterohormones released from the gastrointestinal tract, and the adipokines released from fat tissue have been reviewed by M Plummer, A Reintam, A Deane, K Robinson, J Prins, and B Venkatesh.

In Part 3, the clinical views and attitudes in situations accompanied by challenging metabolic alterations have been addressed. The issue of severe undernutrition was revisited by P Singer and J Cohen; the specificities of traumatic brain injury by H Quintard, C Ichai, and JF Payen; the particular aspects of sepsis and organ failures by V Fraipont and myself; morbid obesity by M Coeffier and F Tamion; and the issues related to burn injury by A Abdullahi, D Patsouris, SR. Costford, and MG Jeschke.

I would like to thank wholeheartedly each one of the authors who brought his own contribution and his personal stone to the huge enterprise of understanding the metabolic response to stress, an indispensable step to improve the quality of care and the quality of the lives of the survivors of critical illness.

Part I
Metabolic Changes

Chapter 2

Successive Phases of the Metabolic Response to Stress

Jean-Charles Preiser, Carole Ichai, and A.B. Johan Groeneveld

Abstract The metabolic response to stress have been selected as an adaptive response to survive critical illness. Several mechanisms well preserved over the evolution, including the stimulation of the sympathetic nervous system, the release of pituitary hormones, a peripheral resistance to the effects of these and other anabolic factors are triggered to increase the provision of energy substrates to the vital tissues. After an acute insult, alternative substrates are used as a result of the loss of control of energy substrate utilization. The clinical consequences of the metabolic response to stress include sequential changes in energy expenditure, stress hyperglycemia, changes in body composition, psychological and behavioral problems. The loss of muscle proteins and function is a major long-term consequence of stress metabolism. Specific therapeutic interventions, including hormone supplementation, enhanced protein intake and early mobilization are investigated.

2.1 Introduction

The understanding and knowledge of metabolic response to critical illness has dramatically changed during the last decade, following several important discoveries in line with the findings of pioneering scientists of the nineteenth and twentieth

J.-C. Preiser, MD, PhD (✉)

Department of Intensive Care, Erasme University Hospital, Free University of Brussels,
808 route de Lennik, Brussels 1070, Belgium
e-mail: Jean-Charles.Preiser@erasme.ulb.ac.be

C. Ichai, MD, PhD

Department of Anesthesiology and Intensive Care, Hôpital Saint-Roch, University of Nice,
Nice, France

A.B.J. Groeneveld, MD, PhD

Department of Intensive Care, Erasmus Medical Center, Rotterdam, The Netherlands

century. In his theory of the evolution, Charles Darwin reported that “It is not the strongest or the most intelligent that survives. It is the most adaptable to change.” This statement is particularly relevant after any life-threatening injury triggering a “critical illness,” when survival in a hostile environment strongly relies on the ability to mount an appropriate adaptive response. In terms of the metabolic response to stress, the principle of homeostasis of Claude Bernard (“The constancy of the internal environment is the condition for a free and independent life”) is highly relevant to the critically ill whose homeostasis must be restored as rapidly as possible to survive the injury. The mechanisms allowing the maintenance of homeostasis, vital functions, and ultimately survival in a hostile environment have been unraveled by Hans Selye, who described the “fight or flight” response, “a nonspecific response to a wide variety of stimuli.” Sir David Cuthbertson described several phases of the metabolic response over time, including the ebb phase and the flow phase. A third sequence, the chronic phase, preceding recovery, was more recently suggested and is probably relevant to the post-injury phase frequently encountered in modern intensive care [1, 2]. The mechanisms of these successive adaptive changes mounted to survive a stress are increasingly understood and are now gathered into a general theory.

2.2 Pathophysiological Mechanisms

The metabolic response to stress involves a neuroendocrine and an inflammatory/immune component. Recent data suggest that hormones released from the adipose tissue and from the gastrointestinal tract can play an important role as well (Fig. 2.1).

The neuroendocrine component is triggered in a region located near the hypothalamus, paraventricular nucleus/locus coeruleus. When a stressor is detected and signaled to the central nervous system, a prototypical response will be triggered, resulting in the activation of the sympathetic nervous system (SNS), the hypothalamic-pituitary axis, and later by behavioral changes. Many different stressors can be sensed and transmitted; for instance, a peripheral tissular injury induced by a trauma will activate afferent nerves, hypoxemia or hypercapnia will trigger chemoreceptors, hypovolemia will activate baroreceptors, and inflammatory mediators will change the phenotype of microglial cells.

The SNS is involved in the fast control of most of the body’s internal organs, via the activation of adrenergic receptors. After any stress, an immediate release of norepinephrine occurs from the postganglionic neuron in response to the stimulation of its nicotinic receptors by acetylcholine released from the preganglionic neurons [3]. The adrenal medulla is a functional sympathetic ganglion, where chromaffin cells release norepinephrine and epinephrine into the bloodstream upon stimulation by the preganglionic neuron.

The activation of the hypothalamus-pituitary axis results in the release of adrenocorticotrophic hormone, thyroid-stimulating hormone, growth hormone, and follicle-stimulating and follicle-luteinizing hormones by the anterior pituitary gland. The circulating levels of hormones released from peripheral glands in response to these




Fig. 2.1 Different levels of the metabolic response to stress. Once a stressor has been sensed, systems/organs are activated (first level). Mediators are released upon activation (second level). Physiological and phenotypical changes are triggered (third level).

pituitary factors are decreased, with the notable exception of cortisol. Peripheral inactivation of the active hormones is the likely mechanism [2], while recently reported alterations in the cortisol breakdown [4] could account for its increased concentration. During the chronic phase, the plasma levels of both pituitary factors and peripheral hormones are lowered, while a peripheral resistance to the effects of growth hormone, insulin, thyroid hormone, and cortisol persists. These hormonal alterations profoundly and sequentially affect the energy, protein, and fat metabolism. The metabolic response to stress thus depends on the time lag after the initial insult.

In addition to these well-characterized pathways, adipokines released from the different cell types of the fat tissue, including leptin, resistin, and adiponectin, are currently being investigated as potential contributors to the metabolic changes related to sepsis [5–8]. The role played by hormones released from the gut is also under scrutiny. Recent data reviewed by Deane et al. [9] indicate that the circulating levels of ghrelin are mostly decreased, while the levels of cholecystokinin and peptide YY are increased [10, 11]. These changes have been related to anorexia, a common feature of the behavioral adaptation to stress. Of note, the metabolic changes associated with adipokines and with the gastrointestinal hormones vary according to the clinical circumstances. The elucidation of the metabolic roles of these hormones requires more clinical research.

The inflammatory component is partially regulated at the level of the central nervous system, via cytokines and inflammatory mediators. The immune response of the host to an infection comprises an innate and a specific immune response. This latter response is subdivided into cell-mediated and humoral components, including antibodies and cytokines. These cytokines can impair some of the body's physiological functions. For example, tumor necrosis factor (TNF), interleukin (IL)-1, and IL-6 play pivotal roles in the metabolic changes associated with sepsis. In addition to typical clinical signs of sepsis (fever, lethargy), these cytokines also induce weight loss and increase proteolysis and lipolysis. In addition, these cytokines trigger anorexia at the hypothalamic level. Several other metabolic effects are indirectly exerted by cytokines via the activation of other cells [12, 13].

The final common pathway of the metabolic response to stress implies the development of a resistance to anabolic signals, including insulin, in order to reset the hierarchy of the delivery of energy substrates to prioritize vital tissues over the insulin-dependent organs, mainly fat and muscle [14, 15]. Therefore, insulin resistance is considered as an adaptive mechanism designed to provide enough glucose to the vital organs, unable to use other energy substrates in stress conditions [16, 17], which results in the inability to suppress central hepatic glucose production [14, 18] and to a decrease of insulin-mediated glucose uptake in the periphery. Insulin resistance is mediated through the reduction of post-receptor insulin signaling defects and downregulation of glucose transporter (GLUT)-4, especially in skeletal muscle. Moreover, impaired nonoxidative glucose disposal results from a reduction in skeletal muscle glycogen synthesis. Despite decreased insulin-mediated glucose uptake, there is an early increase in whole-body glucose uptake, primarily a result of cytokine-mediated upregulation of GLUT-1 [18].

The complexity of the metabolic response is further enhanced by the currently increasing prevalence of obesity and the (type of) metabolic and nutritional support

that is given and may either attenuate or aggravate some of the metabolic responses to stress. The latter depends, among others, on the level of feeding – under- and overnutrition – as well as, indirectly, the level of inflammation that is either evoked or attenuated by nutrition. Also preoperative fasting is a metabolic stress, and losses of energy and proteins following bleeding, hemofiltration, gastrointestinal dysfunction, and others may further compound the metabolic response to stress [19]. Some of the hormones released early from endocrine glands such as (nor)epinephrine, cortisol, thyroid hormone, and glucagon are clearly associated with hypermetabolism aimed at survival, whereas the later changes, with impaired production and/or increased resistance, are more likely adaptive and aimed at a long-term protection of the organism. The latter may, theoretically, be associated with mitochondrial changes, some type of hibernation, and a shutdown of excessive organ function and may thereby, together with an inflammatory response, herald development of multiple organ dysfunction syndrome [19]. Some of these chronic hormonal changes may, however, be regarded as maladaptive when contributing to ultimate mortality by increasing organ dysfunction, immunodepression, and wasting [20–25].

2.3 Clinical Consequences

The clinical consequences of the metabolic response to stress include several different aspects, from changes in resting energy expenditure, use of macronutrients as sources of energy, stress hyperglycemia, and changes in body composition to behavioral changes (Table 2.1 and Fig. 2.2).

2.3.1 Energy Expenditure (EE)

Traditionally, the EE is thought to be lower during the first ebb phase described by Cuthbertson. During the later flow phase, EE is considered to be higher than the EE predicted for a matched healthy subject [26–28]. During the third chronic phase of critical illness, EE decreases slightly. Kreymann et al. serially measured EE in patients with sepsis and septic shock and found lower values during severe sepsis

Table 2.1 Typical patterns of metabolic changes

	Usual patterns of change
Energy expenditure	Decrease (ebb phase or early phase) followed by increase (flow phase or late and recovery phases)
Use of energy substrates	Increased oxidation of carbohydrates, more than lipids/proteins Use of alternative substrates (lactate)
Stress hyperglycemia	Systematic
Changes in body composition	Decreased active cell mass Decreased fat-free mass, increased or unchanged fat mass
Behavior	Lethargy, anorexia




Fig. 2.2 Schematic representation of the three successive phases of the metabolic response to stress, depicting the changes in energy expenditure, and use of energy substrates occurring during the early, late, and recovery phases

[29]. Due to these temporal changes, the actual EE is extremely difficult to predict during critical illness [30]. Indeed, EE is influenced of several physiological derangements, such as fever or hypothermia, changes in heart rate, shivering, agitation, as well as by therapeutic interventions such as sedative agents, nonselective beta-blockers, and active cooling. The use of indirect calorimetry is the best way to assess EE, even though its use to guide the caloric prescription is debatable [31–34].

2.3.2 Use of Energy Substrates

The metabolism of macronutrients is altered at several levels, including the digestive absorption, the intracellular intermediate metabolism, and the oxidation of substrates.

Facing the increased requirements, the oxidation of macronutrients is largely increased during critical illness, and the relative contribution and metabolism of each type of macronutrient is regulated by the circulating hormones (Table 2.2). Overall, the oxidation of carbohydrates is globally more increased than the oxidation of lipids and proteins [35]. Later on, decreased glucose utilization, increased fat turnover, and loss of muscle and visceral (organ) protein mass with wasting occur. A negative nitrogen balance – pointing to increased protein breakdown over protein synthesis – is the ultimate result, even when reprioritization leads to an increased overall hepatic protein synthesis. Indeed, muscle may lose amino acids at the benefit of the liver. These changes are hardly amenable to any fruitful intervention to improve protein synthesis, attenuate lipogenesis, and thereby conserve lean body mass needed for rehabilitation.

Carbohydrates Glucose is the preferential energy substrate during critical illness and will be able to yield 2 ATP after anaerobic glycolysis and 36 additional molecules of ATP by the Krebs cycle when the mitochondrion is functional. At the whole-body level, changes in the metabolism of carbohydrates include the rapid

Table 2.2 Use of substrates during the successive phases

Macronutrient	Anabolic	Catabolic
Carbohydrates	Insulin	Cortisol Glucagon Growth hormone Catecholamines
Lipids	Insulin	Catecholamines
Proteins	Insulin Growth hormone – IGF-1 Testosterone Catecholamines	Cortisol Glucagon Catecholamines

utilization of the glycogen stores, followed by a high level of endogenous glucose production from lactate, glycerol, and alanine in the liver, the kidney, and the intestine [36, 37]. As the turnover of glucose is increased, plasma concentrations of glucose will rise, resulting in the typical stress hyperglycemia [18]. While nonoxidative metabolism (e.g., glycogen synthesis) is impaired, oxidative glucose metabolism is upregulated early [38]. Alterations in the digestion of dietary carbohydrates occur as well: once ingested, the long molecules of polysaccharides are cleaved into oligosaccharides (3–10 sugars) by the amylase enzymes. The resulting oligosaccharides will be cleaved by enzymes of the intestinal brush border. The activity of one of these enzymes, lactase, can be inhibited in the critically ill, thereby reducing the absorption of enteral carbohydrates [39].

Use of lactate as an alternative substrate Alteration of lactate metabolism is one of the prominent component of the metabolic stress response. Lactate is a physiological substrate (carbohydrate) issued from pyruvate reduction during glycolysis. In stable conditions, lactate production and elimination are equivalent, i.e., 1200–1500 mmol per day, leading to a stable blood lactate concentration of 0.8–1.2 mmol/L [40]. Most organs, except those without mitochondria, simultaneously release and take up lactate. As a result, the net flux of lactate depends on the difference between release and uptake and varies upon organs and their energetic condition [41]. In stable conditions, the brain, muscles, and digestive tract are producing lactate organs, whereas the liver is responsible for more than 70 % of lactate clearance. Lactatemia and lactate metabolism (turnover) are often confused. Lactatemia indicates an instantaneous equilibrium between total body lactate production and clearance. Accordingly, lactatemia can be in a normal value, while lactate turnover can be normal, high, or low, just indicating that there is an equilibrium between production and elimination.

Lactate, is a physiological intermediate energetic substrate. The Cori cycle (conversion of lactate into glucose) confirms the ability of lactate to serve as a very efficient interorgan shuttle, allowing to provide fuel useable by organs in various stress conditions [42]. For instance, red blood cells not equipped with mitochondria produce ATP only *via* an anaerobic glycolysis leading to lactate production, the latter being further metabolized in glucose in the liver in the presence of oxygen. Growing data support that these exchanges are favored during stress condition and that lactate “per se” is at least a useful if not an obligatory substrate used by organs

and tissues during energetic crisis conditions and has been particularly demonstrated to fuel the heart and brain.

At rest, the heart consumes energy issued for 60–90 % from fatty acids β -oxidation. But in case of hypoxia such as during myocardial ischemia, increased O_2 consumption, or decreased O_2 delivery, metabolic pathways shift toward a preferential use of carbohydrate oxidation for ATP production [43]. The role of lactate as a myocardial fuel has been confirmed experimentally during septic and hemorrhagic shocks [44, 45].

Stress hyperglycemia The etiology of hyperglycemia in type 2 diabetes is a combination of insulin resistance and beta cell secretory defects [14, 18]. The development of stress hyperglycemia involves a much more dramatic, complex interplay of counterregulatory hormones such as catecholamines, growth hormone, and cortisol, and cytokine resulting in excessive hepatic glucose production (from gluconeogenesis and glycogenolysis) and insulin resistance. Increased hepatic output of glucose, particularly through gluconeogenesis, appears to be the most important contributor to stress hyperglycemia (see Chap. 8 for further discussion).

Numerous association studies [46, 47] confirm the presence of a U-shaped relationship between admission BG value and outcome, i.e., low and high BG are associated with poor outcome. An admission BG value of 5.5–6.1 mmol/L is associated with the lowest mortality rate. Similarly, high GV and low BG complexity are also associated with a worsened outcome.

Lipids The use of lipids as energy substrate is relatively less increased than carbohydrates, during critical illness [35]. Indeed, the conversion of lipids into ATP requires large amounts oxygen and functional mitochondria. During critical illness, endogenous triglycerides stored in the adipose tissue and exogenous triglycerides released from chylomicrons and other lipoproteins are avidly hydrolysed to release FFAs and glycerol into the bloodstream. In contrast to physiological conditions, this increased lipolysis cannot be efficiently inhibited by infusion of carbohydrates. The oxidation of FFAs is increased in peripheral tissues, while in the liver they are converted to ketone bodies or re-esterified to triglycerides and released into the bloodstream as very-low-density lipoprotein (VLDL), which is subject to impaired clearance. However, the production of FFAs from exogenous and endogenous triglycerides still exceeds the utilization of FFAs, and plasma FFA levels are typically increased in critically ill patients. Overall, the metabolism of lipids is increased, although complete oxidation can only be achieved in tissues where mitochondria are functional.

Proteins Under normal conditions, proteins are constantly broken down and replaced in a highly selective and closely balanced process. The majority of intracellular proteins are degraded via activation of the ubiquitin-proteasome pathway. In a series of enzymatic reactions, ubiquitin forms a chain on a protein to be degraded. Once tagged, the protein is recognized by a proteasome. The protein unravels and is injected into the central core of the proteasome where it is broken down into peptides.

Stress metabolism is characterized by over-activation of the ubiquitin-proteasome pathway that causes excessive protein degradation and muscle wasting.

Overall, the large increases in protein breakdown are partially balanced by increased protein synthesis (of inflammatory mediators). The amino acids released during the degradation of proteins will be either reused (cf. alanine, glutamine) or oxidized and will provide waste products: urea and ammonium. The nitrogen balance will be negative, with a rate of breakdown largely exceeding the rate of synthesis. Consequently, the stores of proteins, i.e., the skeletal muscles, will be rapidly depleted. These losses are related to the large wastage of muscles, which is involved in ICU-acquired weakness [48–52]. This is one of the most devastating consequences of the metabolic response to stress. A major complaint of patients who had a prolonged stay in an ICU is weakness, even a considerable time after discharge. In a study by Herridge et al., survivors of an acute respiratory distress syndrome had persistent muscle wasting and weakness 5 years after discharge from the ICU [48]. Thus, it is essential to take muscle function into consideration when assessing and monitoring the nutritional status of ICU patients.

2.3.3 Changes in Body Composition

The changes in body composition systematically found during critical illness include a loss of lean body mass and a relative preservation of the fat tissue [6]. As a result, body cell mass is typically decreased, while extracellular fluid is increased. Recently, functional and morphological changes of the fat tissue have been identified. These changes can be summarized as a preservation of the fat mass, with increased number of small adipocytes and increased infiltration of the fat tissue by macrophages [53]. Functionally, these changes result in increased lipid storage.

2.3.4 Psychosocial and Behavioral Problems

Long-term psychosocial and behavioral issues have been consistently reported in different cohorts of critically ill patients [54]. Some of these changes, such as prolonged catabolism, are clearly related to the metabolic response to critical illness. Behavioral changes, including anorexia, might be related to changes in the release of gastrointestinal hormones [10].

2.4 Therapeutic Implications

Generally speaking, hormone repletion in the chronic phase by exogenous administration, even though attempted in the past on numerous occasions, has not been successful in attenuation morbidity and mortality of the critically ill, even though successful from a metabolic point of view [55]. For instance, growth or

thyroid hormone supplementation may have anabolic effects by increasing protein synthesis and ameliorating protein breakdown in the critically ill but may even increase morbidity and mortality because of other unwanted effects [56–59]. Although insulin may have some albeit controversial anabolic effects and may help, by overcoming resistance and glucose control, the patient-centered outcome effects are highly controversial. Recently, expert guidelines recommend to avoid severe hyperglycemia, although an universally acceptable high limit to titrate insulin therapy cannot be defined [60–62]. Sex steroid hormones are still explored to increase anabolism [63]. Intervening with intermediary metabolism by administering (pharmacologic quantities of) substrates, often together with other nutritional supplements seemed promising in the last decades, but recent evidence suggests that this may be less helpful. For instance, glutamine supplementation of nutrition improved immunologic, gut function and protein metabolism, and even patient-centered outcomes in prior studies, but recent, large studies demonstrate that this may be associated with worse rather than better vital outcomes [64]. This is not to say that adequate nutrition lacks sufficient evidence in improving patient outcomes. Whatever route, quantities, or composition is chosen – these issues remain highly controversial – there is no doubt that prolonged starvation and resultant malnutrition in the critically ill substantially contribute to morbidity and mortality. The question, however, remains whether altered composition – branched chain amino acids, immunonutrition with L-arginine and glutamine, antioxidants, and others – is meaningfully contributing to altered utilization and metabolic processes, particularly in sepsis and trauma patients where metabolism is driven by underlying inflammatory and host defense mechanisms rather than by exogenous supply. Then, metabolic or nutritional support may have some cosmetic effects, including normalization of altered plasma and tissue amino acid and protein levels, but without large effects on gluconeogenesis from protein breakdown, protein synthesis and lean body mass, and even in the presence of hyperinsulinemia. Early mobilization and the avoidance of prolonged sedation are other daily therapeutic measures that are likely to attenuate catabolism.

Hence, other interventions include raising ambient temperature (to decrease energy-consuming heat production) and administering beta-blockers to attenuate sympathetic overstimulation, inflammation, and protein breakdown and to improve organ and muscle function, particularly in burns and sepsis [65, 66]. The latter is still under investigation and certainly not uniformly and routinely accepted. Animal studies suggest that gut-derived ghrelin has anabolic properties, and studying its administration in critically ill patients has been proposed, since circulating levels have been found to be increased or lowered depending on the phase of disease.

2.5 Conclusion

The metabolic response to stress is a complex combination of neurological, endocrine, immune, and inflammatory mechanisms which lead to multiple functional changes in each tissue of the body. A better understanding of the physiology of this

response is needed when the progresses of intensive care medicine allow the survival of patients whose adaptive metabolic mechanisms are developing. Therapeutic interventions need to account for the complexity and sequential pattern of the metabolic response to critical illness.

References

1. Preiser JC, Ichai C, Orban JC, Groeneveld ABJ (2014) Metabolic response to the stress of critical illness. *Br J Anaesth* 113:945–954
2. Van den Berghe G, de Zegher F, Bouillon R (1998) Clinical review 95: acute and prolonged critical illness as different neuroendocrine paradigms. *J Clin Endocrinol Metab* 83:1827–1834
3. Hamill RW, Woolf PD, McDonald JV, Lee LA, Kelly M (1987) Catecholamines predict outcome in traumatic brain injury. *Ann Neurol* 21:438–443
4. Boonen E, Vervenne H, Meersseman P, Andrew R, Mortier L, Declercq PE, Vanwijngaerden YM, Spriet I, Wouters PJ, Vander Perre S, Langouche L, Vanhorebeek I, Walker BR, Van den Berghe G (2013) Reduced cortisol metabolism during critical illness. *N Engl J Med* 368:1477–1488
5. Koch A, Gressner OA, Sanson E, Tacke F, Trautwein C (2009) Serum resistin levels in critically ill patients are associated with inflammation, organ dysfunction and metabolism and may predict survival of non-septic patients. *Crit Care* 13:R95
6. Marques MB, Langouche L (2013) Endocrine, metabolic, and morphologic alterations of adipose tissue during critical illness. *Crit Care Med* 41:317–325
7. Hillenbrand A, Weiss M, Knippschild U, Wolf AM, Huber-Lang M (2012) Sepsis-induced adipokine change with regard to insulin resistance. *Int J Inflam* 2012:972368
8. Fantuzzi G (2009) Adiponectin and inflammation. *Am J Physiol Endocrinol Metab* 296(2), E397
9. Deane A, Chapman MJ, Fraser RJL, Horowitz M (2010) Bench-to-bedside review: the gut as an endocrine organ in the critically ill. *Crit Care* 14:228
10. Nematy M, O'Flynn JE, Wandrag L, Brynes AE, Brett SJ, Patterson M, Ghatei MA, Bloom SR, Frost GS (2006) Changes in appetite related gut hormones in intensive care unit patients: a pilot cohort study. *Crit Care* 10:R10
11. Nematy M, Brynes AE, Hornick PI, Patterson M, Ghatei MA, Bloom SR, Brett SJ, Frost GS (2007) Postprandial ghrelin suppression is exaggerated following major surgery; implications for nutritional recovery. *Nutr Metab (Lond)* 4:20
12. Losser MR, Damoisel C, Payen D (2010) Bench-to-bedside review: glucose and stress conditions in the intensive care unit. *Crit Care* 14:231
13. Plank LD, Hill GL (2000) Sequential metabolic changes following induction of systemic inflammatory response in patients with severe sepsis or major blunt trauma. *World J Surg* 24:630–638
14. Lena D, Kalfon P, Preiser JC, Ichai C (2011) Glycemic control in the intensive care unit and during the postoperative period. *Anesthesiology* 114:438–444
15. Biolo G, Grimble G, Preiser JC, Leverve X, Jolliet P, Planas M, Roth E, Wernerman J, Pichard C, European Society of Intensive Care Medicine Working Group on Nutrition and Metabolism (2002) Position paper of the ESICM Working Group on Nutrition and Metabolism. Metabolic basis of nutrition in intensive care unit patients: ten critical questions. *Intensive Care Med* 28:1512–1520
16. Soeters MR, Soeters PB (2012) The evolutionary benefit of insulin resistance. *Clin Nutr* 31:1002–1007
17. Marik PE, Bellomo R (2013) Stress hyperglycemia: an essential survival response! *Crit Care* 17:305

18. Dungan KM, Braithwaite SS, Preiser JC (2009) Stress hyperglycaemia. *Lancet* 373:1798–1807
19. Singer M, De Santis V, Vitale D, Jeffcoate W (2004) Multiorgan failure is an adaptive, endocrine-mediated metabolic response to overwhelming systemic inflammation. *Lancet* 364:545–548
20. Desborough JP (2000) The stress response to trauma and surgery. *Br J Anaesth* 85: 109–117
21. Siegel JH, Cerra FB, Coleman B, Giovannini I, Shetye M, Border JR, McMenamy RH (1979) Physiological and metabolic correlations in human sepsis. *Surgery* 86:163–193
22. Wilmore DW (2000) Metabolic response to severe surgical stress: overview. *World J Surg* 24:705–711
23. Kyle UG, Jolliet P, Genton L, Meier CA, Mensi N, Graf JD, Chevrolet JC, Pichard C (2005) Clinical evaluation of hormonal stress state in medical ICU patients: a prospective blinded observational study. *Intensive Care Med* 31:1669–1675
24. Donatelli F, Corbella D, Di Nicola M, Carli F, Lorini L, Fumagalli R, Biolo G (2011) Preoperative insulin resistance and the impact of feeding on postoperative protein balance: a stable isotope study. *J Clin Endocrinol Metab* 896:E1789–E1797
25. Hoffer LJ, Bistrian BR (2013) Why critically ill patients are protein depleted. *J Parenter Enteral Nutr* 37(3):300–309
26. Magnuson B, Peppard A, Auer Flomenhoft D (2011) Hypocaloric considerations in patients with potentially hypometabolic disease States. *Nutr Clin Pract* 26:253–260
27. McClave SA, Martindale RG, Kiraly L (2013) The use of indirect calorimetry in the intensive care unit. *Curr Opin Clin Nutr Metab Care* 16:202–208
28. Siirala W, Olkkola KT, Noponen T, Vuori A, Aantaa R (2010) Predictive equations overestimate the resting energy expenditure in amyotrophic lateral sclerosis patients who are dependent on invasive ventilation support. *Nutr Metab (Lond)* 7:70
29. Kreyman G, Grosser S, Buggisch P, Gottschall C, Matthaei S, Greten H (1993) Oxygen consumption and resting metabolic rate in sepsis, sepsis syndrome, and septic shock. *Crit Care Med* 21:1012–1019
30. Uehara M, Plank LD, Hill GL (1999) Components of energy expenditure in patients with severe sepsis and major trauma: a basis for clinical care. *Crit Care Med* 27:1295–1302
31. Vincent JL, Preiser JC (2013) When should we add parenteral to enteral nutrition? *Lancet* 381:354–355
32. Heidegger CP, Berger MM, Graf S, Zingg W, Darmon P, Costanza MC, Thibault R, Pichard C (2013) Optimisation of energy provision with supplemental parenteral nutrition in critically ill patients: a randomised controlled clinical trial. *Lancet* 381:385–393
33. Casaer MP, Mesotten D, Hermans G, Wouters PJ, Schetz M, Meyfroidt G, Van Cromphaut S, Ingels C, Meersseman P, Muller J, Vlasselaers D, Debaveye Y, Desmet L, Dubois J, Van Assche A, Vanderheyden S, Wilmer A, Van den Berghe G (2011) Early versus late parenteral nutrition in critically ill adults. *N Engl J Med* 365:506–517
34. Schetz M, Casaer MP, Van den Berghe G (2013) Does artificial nutrition improve outcome of critical illness? *Crit Care* 17:302
35. Tappy L, Schwarz JM, Schneiter P, Cayeux C, Revely JP, Fagerquist CK, Jéquier E, Chioléro R (1998) Effects of isoenergetic glucose-based or lipid-based parenteral nutrition on glucose metabolism, de novo lipogenesis, and respiratory gas exchanges in critically ill patients. *Crit Care Med* 26:860–867
36. Watford M (2005) Is the small intestine a gluconeogenic organ. *Nutr Rev* 63:356–360
37. Battezzati A, Caumo A, Martino F et al (2004) Nonhepatic glucose production in humans. *Am J Physiol Endocrinol Metab* 286:E129–E135
38. Shangraw RE, Jahoor F, Wolfe RR, Lang CH (1996) Pyruvate dehydrogenase inactivity is not responsible for sepsis-induced insulin resistance. *Crit Care Med* 24:566–574
39. Burgstad CM, Besanko LK, Deane AM, Nguyen NQ, Saadat-Gilani K, Davidson G, Burt E, Thomas A, Holloway RH, Chapman MJ, Fraser RJ (2013) Sucrose malabsorption and impaired mucosal integrity in enterally fed critically ill patients: a prospective cohort observational study. *Crit Care Med* 41:1221–1228

40. Orban JC, Leverve X, Ichai C (2011) Lactate: métabolisme et physiopathologie. In: Ichai C, Quintard H, Orban JC (eds) *Désordres métaboliques et réanimation : de la physiopathologie au traitement*. Springer, Paris, pp 181–198
41. Van Hall G, Stromstadt M, Rasmussen P et al (2004) Blood lactate is an important source of energy for the human brain. *J Cereb Blood Flow Metab* 29:1121–1129
42. Leverve XM (1999) Energy metabolism in critically ill patients: lactate is a major oxidizable substrate. *Curr Opin Clin Nutr Metab Care* 2:165–169
43. Ichai C, Armando G, Orban JC, Berthier F, Rami L, Samat-Long C, Grimaud D, Leverve X (2009) Sodium lactate versus mannitol in the treatment of intracranial hypertensive episodes in severe traumatic brain-injured patients. *Intensive Care Med* 35(3):471–479
44. Stanley WC, Recchia FA, Lopasschuk GD (2005) Myocardial substrate metabolism in the normal and failing heart. *Physiol Rev* 85:1093–1129
45. Levy B, Gibot S, Franck P, Cravoisy A, Bollaert PE (2005) Relation between muscle Na⁺ + K⁺ ATPase activity and raised lactate concentrations in septic shock: a prospective study. *Lancet* 365:871–875
46. Krinsley JS, Egi M, Kiss A, Devendra AN, Schuetz P, Maurer PM, Schultz MJ, van Hooijdonk RT, Kiyoshi M, Mackenzie IM, Annane D, Stow P, Nasraway SA, Holewinski S, Holzinger U, Preiser JC, Vincent JL, Bellomo R (2013) Diabetic status and the relation of the three domains of glycemic control to mortality in critically ill patients: an international multicenter cohort study. *Crit Care* 17:R37
47. Falciglia M, Freyberg RW, Almenoff PL, D'Alessio DA, Render ML (2009) Hyperglycemia-related mortality in critically ill patients varies with admission diagnosis. *Crit Care Med* 37:3001–3009
48. Herridge MS, Tansey CM, Matte A et al (2011) Functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med* 364:1293–1304
49. Hill GL (1992) Jonathan E. Rhoads Lecture. Body composition research: implications for the practice of clinical nutrition. *JPEN J Parenter Enteral Nutr* 16:197–218
50. Lecker SH (2006) Protein degradation by the ubiquitin-proteasome pathway in normal and disease states. *J Am Soc Nephrol* 17:1807–1819
51. Mitch WE, Goldberg AL (1996) Mechanisms of muscle wasting. The role of the ubiquitin-proteasome pathway. *N Engl J Med* 335:1897–1905
52. Hill NE, Murphy KG, Singer M (2012) Ghrelin, appetite and critical illness. *Curr Opin Crit Care* 18:199–205
53. Langouche L, Perre SV, Thiessen S, Gunst J, Hermans G, D'Hoore A, Kola B, Korbonits M, Van den Berghe G (2010) Alterations in adipose tissue during critical illness: an adaptive and protective response? *Am J Respir Crit Care Med* 2010(182):507–516
54. Broomhead LR, Brett SJ (2002) Clinical review: intensive care follow-up – what has it told us? *Crit Care* 6:411–417
55. Ligtenberg JJ, Girbes AR, Beentjes JA, Tulleken JE, Van der Werf TS, Zijlstra JG (2001) Hormones in the critically ill patients: to intervene or not to intervene? *Intensive Care Med* 27:1567–1577
56. Takala J, Ruokonen E, Webster NR et al (1999) Increased mortality associated with growth hormone treatment in critically ill adults. *N Engl J Med* 341:785–792
57. Ruokonen E, Takala J (2002) Dangers of growth hormone therapy in critically ill patients. *Curr Opin Clin Nutr Metab* 5:199–209
58. Voerman HJ, Strack van Schijndel RJM, Groeneveld ABJ, de Boer H, Nauta JJP, van der Veen EA, Thijs LG (1992) Effects of recombinant human growth hormone in patients with severe sepsis. *Ann Surg* 216:648–655
59. Schulman RC, Mechanick JI (2012) Metabolic and nutrition support in the chronic critical illness syndrome. *Respir Care* 57:958–978
60. Ichai C, Preiser JC, Société Française d'Anesthésie-Réanimation; Société de Réanimation de langue Française; Experts group (2010) International recommendations for glucose control in adult non diabetic critically ill patients. *Crit Care* 14:R166
61. Groeneveld ABJ, Beishuizen A, Visser FC (2002) Insulin: a wonder drug in the critically ill? *Crit Care* 6:102–105