

BIOMASS AND BIOFUELS SERIES

Bioenergy Feedstocks

Breeding and Genetics

EDITED BY

Malay C. Saha
Hem S. Bhandari
Joseph H. Bouton



WILEY-BLACKWELL

Bioenergy Feedstocks

Bioenergy Feedstocks

Breeding and Genetics

Edited by

Malay C. Saha

Hem S. Bhandari

Joseph H. Bouton

 **WILEY-BLACKWELL**

A John Wiley & Sons, Inc., Publication

This edition first published 2013 © 2013 by John Wiley & Sons, Inc.

Wiley-Blackwell is an imprint of John Wiley & Sons, formed by the merger of Wiley's global Scientific, Technical and Medical business with Blackwell Publishing.

Editorial offices: 2121 State Avenue, Ames, Iowa 50014-8300, USA
The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK
9600 Garsington Road, Oxford, OX4 2DQ, UK

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell.

Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by Blackwell Publishing, provided that the base fee is paid directly to the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923. For those organizations that have been granted a photocopy license by CCC, a separate system of payments has been arranged. The fee codes for users of the Transactional Reporting Service are ISBN-13: 978-0-4709-6033-2/2013.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. This publication is designed to provide accurate and authoritative information in regard to the subject matter covered. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

Limit of Liability/Disclaimer of Warranty: While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with the respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. It is sold on the understanding that the publisher is not engaged in rendering professional services and neither the publisher nor the author shall be liable for damages arising herefrom. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

Library of Congress Cataloging-in-Publication Data is available upon request.

A catalogue record for this book is available from the British Library.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Cover design by Matt Kuhns

Set in 10/11.5pt Times New Roman by Aptara® Inc., New Delhi, India

Contents

<i>The Editors</i>	xi
<i>List of Contributors</i>	xiii
<i>Preface</i>	xix
1 Introduction	1
1.1 Historical Development	2
1.2 Cultivar Development	2
1.3 Breeding Approach	3
1.4 Molecular Tools	3
1.5 Future Outlook	4
References	4
2 Switchgrass Genetics and Breeding Challenges	7
2.1 Introduction	7
2.2 Origin and Distribution	9
2.3 Growth and Development, Genome Structure and Cytogenetics	9
2.3.1 Growth and Development	10
2.3.2 Genome Structure and Cytogenetics	12
2.4 Genetic Diversity	12
2.5 Phenotypic Variability and Inheritance	13
2.6 Conventional Breeding Approaches	14
2.6.1 Early Work	15
2.6.2 Systematic Recurrent Selection	15
2.6.3 Heterosis	17
2.7 Molecular Breeding	18
2.7.1 Molecular Markers Used for Switchgrass and Other Polyploids	18
2.7.2 Molecular Mapping	20
2.7.3 Association Mapping	22
2.7.4 Transgenic Approaches	23
2.8 Conclusions and Future Directions	23
References	24
3 Switchgrass Genomics	33
3.1 Introduction	33
3.2 Genome Sequencing	34
3.2.1 Other Available Sequence Resources	35
3.3 Analysis of Expressed Sequences in Switchgrass	36

3.4	Linkage Mapping	40
3.5	Cytoplasmic Genome	42
3.6	Genome-enabled Improvement of Switchgrass	42
3.7	Conclusions	45
	References	45
4	Germplasm Resources of <i>Miscanthus</i> and Their Application in Breeding	49
4.1	Introduction	49
4.2	Species Belonging to <i>Miscanthus</i> Genus, Their Characteristics, and Phylogenetic Relationships	50
4.2.1	Section: Eumiscanthus	50
4.2.2	Section: Triarrhena	53
4.2.3	Section: Kariyasu	54
4.3	Natural Hybrids between <i>Miscanthus</i> Species	55
4.4	Karyotype Analysis	55
4.5	Phylogenetic Relationships between <i>Miscanthus</i> Species	56
4.6	Genetic Improvement of <i>Miscanthus</i>	57
4.6.1	Germplasm Collection and Management	57
4.6.2	Artificial Hybridization	57
4.6.3	Polyploidization	58
4.7	Variations in Several Agronomical Traits Related to Yield and Plant Performance	58
4.7.1	Variation in Flowering Time	58
4.7.2	Variation in Cold Tolerance	58
4.7.3	Variation in Lignin, Cellulose, and Mineral Content	59
4.8	Molecular Resources	60
4.8.1	Development of Linkage Map for <i>Miscanthus</i>	60
4.8.2	QTL Analysis of Traits Related to Yield and Mineral Content	60
4.8.3	Molecular Markers for Hybrids Identification	61
4.9	Transgenic <i>Miscanthus</i>	61
4.10	Future Studies	62
	References	62
5	Breeding <i>Miscanthus</i> for Bioenergy	67
5.1	Introduction	67
5.2	<i>Miscanthus</i> as a Biomass Crop	67
5.3	Breeding Strategy	68
5.3.1	Collection and Characterization	68
5.3.2	Hybridization	68
5.3.3	Ex Situ Phenotypic Characterization	69
5.3.4	Large-scale Demonstration Trials	69
5.4	Genetic Diversity	69
5.5	Breeding Targets	70
5.5.1	Biomass Yield	70
5.5.2	Morphological Traits Contributing to High Yield Potential	75
5.5.3	Seed Propagation: Crop Diversification and Reducing the Cost of Establishment	77

5.6	Incorporating Bioinformatics, Molecular Marker-Assisted Selection (MAS), and Genome-Wide Association Selection (GWAS)	77
5.7	Summary	78
	Acknowledgments	79
	References	79
6	Breeding Sorghum as a Bioenergy Crop	83
6.1	Introduction	83
6.2	Botanical Description and Evolution	84
6.2.1	Basic Characteristics	84
6.2.2	Evolution and Distribution	85
6.3	Traditional Breeding and Development	86
6.3.1	Initial Sorghum Improvement	86
6.3.2	Development of Hybrid Sorghum and Heterosis	86
6.3.3	Current Sorghum Breeding Approaches	88
6.3.4	Germplasm Resources	88
6.4	Approaches to Breeding Sorghum as a Bioenergy Crop	90
6.4.1	Grain Sorghum	90
6.4.2	Sweet Sorghum	90
6.4.3	Biomass Sorghum	93
6.5	Composition in Energy Sorghum Breeding	93
6.6	Genetic Variation and Inheritance	95
6.6.1	Grain Sorghum	95
6.6.2	Grain Quality/Starch Composition	96
6.6.3	Dual Purpose—Grain and Stalk	97
6.6.4	Soluble Carbohydrates	97
6.6.5	Breeding for Stress Tolerance	99
6.7	Wide Hybridization	106
6.7.1	Interspecific Hybridization	106
6.7.2	Intergeneric Hybridization	107
6.8	Conclusions	107
	References	107
7	Energy Cane	117
7.1	Introduction	117
7.2	Sugar and Energy Production Systems	118
7.2.1	Current Global Sugarcane Production	118
7.2.2	Bioenergy Production from Sugarcane in Brazil	120
7.2.3	Overview of Main Components in Existing Sugarcane Production Systems	120
7.2.4	Overview and Potential Trends	123
7.3	Sugarcane Improvement	124
7.3.1	Taxonomy and Crop Physiology	124
7.3.2	History of Sugarcane Breeding	127
7.3.3	Basic Features of Sugarcane Breeding Programs	128
7.3.4	Composition of Cane for Sugar or Energy Production	130
7.3.5	Application of Molecular Genetics in Developing Energy Cane	131

7.4	Selection of Sugarcane Genotypes for Energy Production	134
7.4.1	Overall Directions	134
7.4.2	Example of Economic Weightings for Selecting Sugarcane for Energy Products	136
7.4.3	Progress in Breeding for Energy Production	138
7.5	Conclusion	141
	Acknowledgments	141
	References	141
8	Breeding Maize for Lignocellulosic Biofuel Production	151
8.1	Introduction	151
8.2	General Attributes of Maize as a Biofuel Crop	151
8.3	Potential Uses of Maize Stover for Bioenergy	153
8.4	Breeding Maize for Biofuels	154
8.4.1	Selection Criteria	154
8.4.2	Stover Yield	157
8.4.3	Maximum Biomass Yield and the Effects of Time and Latitude	159
8.4.4	Stover Quality	161
8.4.5	Sustainability Parameters	163
8.4.6	Breeding Methods	164
8.5	Single Genes and Transgenes	165
8.6	Future Outlook	167
	References	167
9	Underutilized Grasses	173
9.1	Introduction	173
9.2	Prairie Cordgrass	174
9.2.1	Importance	174
9.2.2	Genetic Variation and Breeding Methods	176
9.2.3	Future Goals	180
9.3	Bluestems	181
9.3.1	Importance	181
9.3.2	Genetic Variation and Breeding Methods	184
9.3.3	Future Goals	190
9.4	Eastern Gamagrass	191
9.4.1	Importance	191
9.4.2	Genetic Variation and Breeding Methods	192
9.4.3	Future Goals	196
	References	197
10	Alfalfa as a Bioenergy Crop	207
10.1	Introduction	207
10.2	Biomass for Biofuels	208
10.2.1	Lignocellulose-based Biofuels	208
10.2.2	Plant Cell Wall Components	209
10.3	Why Alfalfa?	211
10.3.1	Background	211
10.3.2	Prospect as a Biofuel Feedstock	212

10.4	Breeding Strategies	213
10.4.1	Germplasm Resources	213
10.4.2	Cultivar Development	214
10.4.3	Synthetic Cultivars and Heterosis	214
10.4.4	Molecular Breeding	215
10.4.5	Trait Integration Through Biotechnology	216
10.5	Breeding Targets	217
10.5.1	Biomass Yield	217
10.5.2	Forage Quality and Composition	218
10.5.3	Stress Tolerance	219
10.5.4	Winter Hardiness	220
10.6	Management and Production Inputs	221
10.7	Processing for Biofuels	222
10.8	Additional Value from Alfalfa Production	223
10.8.1	Environmental Benefits	223
10.8.2	Alfalfa Co-products	223
10.9	Summary	223
	Acknowledgments	224
	References	224
11	Transgenics for Biomass	233
11.1	Introduction	233
11.1.1	Biomass for Biofuels	233
11.1.2	Biofuels	234
11.1.3	Lignocellulosic Biomass	234
11.2	Transgenic Approaches	235
11.2.1	Biolistics Transformation	235
11.2.2	Agrobacterium-mediated Transformation	236
11.3	Transgenic Approaches for Biomass Improvement	237
11.3.1	Improving Biomass Yield	237
11.3.2	Modifying Biomass Composition	240
11.3.3	Regulatory Issues of Transgenic Bioenergy Crops	242
11.4	Summary	242
	Acknowledgments	242
	References	243
12	Endophytes in Low-input Agriculture and Plant Biomass Production	249
12.1	Introduction	249
12.2	What are Endophytes?	249
12.3	Endophytes of Cool Season Grasses	251
12.4	Endophytes of Warm Season Grasses	251
12.5	Endophytes of Woody Angiosperms	253
12.6	Other Fungal Endophytes	253
12.7	Endophytes in Biomass Crop Production	254
12.8	The Use of Fungal Endophytes in Bioenergy Crop Production Systems	256
12.9	Endophyte Consortia	256

12.10	Source of Novel Compounds	257
12.11	Endophyte in Genetic Engineering of Host Plants	258
12.12	Conclusions	258
	Acknowledgments	259
	References	259

<i>Index</i>		267
--------------	--	-----

Color plate is located between pages 172 and 173.

The Editors

Malay C. Saha Forage Improvement Division (FID)
The Samuel Roberts Noble Foundation, Inc.
Ardmore, OK
mcsaha@noble.org

Hem S. Bhandari Department of Plant Sciences
University of Tennessee
Knoxville, TN
hsbhandari@utk.edu

Joseph H. Bouton Forage Improvement Division (FID)
The Samuel Roberts Noble Foundation, Inc.
Ardmore, OK
jhbouton@noble.org

List of Contributors

Laura Bartley	Department of Botany and Microbiology University of Oklahoma Norman, OK
Kishor Bhattarai	Forage Improvement Division The Samuel Roberts Noble Foundation Ardmore, OK
Arvid Boe	South Dakota State University Brookings, SD
John Clifton Brown	Institute of Biological, Environmental and Rural Sciences (IBERS) Aberystwyth University Gogerddan, Aberystwyth, Ceredigion, Wales, UK
E. Charles Brummer	Forage Improvement Division The Samuel Roberts Noble Foundation Ardmore, OK
Kelly D. Craven	Plant Biology Division The Samuel Roberts Noble Foundation Ardmore, OK
Chris Davey	Institute of Biological, Environmental and Rural Sciences (IBERS) Aberystwyth University Gogerddan, Aberystwyth, Ceredigion, Wales, UK
Natalia de Leon	Department of Agronomy University of Wisconsin Madison, WI
Iain Donnison	Institute of Biological, Environmental and Rural Sciences (IBERS) Aberystwyth University Gogerddan, Aberystwyth, Ceredigion, Wales, UK

- Maria Stefanie Dwiyanti** Energy Biosciences Institute
University of Illinois
Urbana, IL
- Kerrie Farrar** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- Sita R. Ghimire** Center for Agricultural and Environmental Biotechnology
RTI International
Research Triangle Park, NC
- J. Gonzalez-Hernandez** South Dakota State University
Brookings, SD
- C. Frank Hardin** Forage Improvement Division
The Samuel Roberts Noble Foundation
Ardmore, OK
- Charlotte Hayes** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- Maurice Hinton-Jones** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- Lin Huang** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- Elaine Jensen** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- Phillips Jackson** CSIRO Plant Industry
Australian Tropical Science Innovation Precinct
Townsville, Australia
- Laurence Jones** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK

-
- Shawn M. Kaeppler** Department of Agronomy
University of Wisconsin
Madison, WI
- Joe G. Lauer** Department of Agronomy
University of Wisconsin
Madison, WI
- D. K. Lee** University of Illinois
Urbana, IL
- Anne Maddison** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- Heike Meyer** Julius Kühn-Institute (JKI)
Federal Research Centre for Cultivated Plants
Bundesallee, Germany
- Maria J. Monteros** Forage Improvement Division
The Samuel Roberts Noble Foundation
Ardmore, OK
- John Norris** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- Sarah Purdy** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- A. Lane Rayburn** University of Illinois
Urbana, IL
- Paul Robson** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- W. L. Rooney** Department of Soil and Crop Sciences
Texas A&M University
College Station, Texas
- Aaron Saathoff** Grain, Forage, and Bioenergy Research Unit
USDA-ARS and Department of Agronomy and Horticulture
University of Nebraska
Lincoln, NE

- Cosentino Salvatore** Department of Agriculture and Food Science
University of Catania
Via Valdisavoia, Catania, Italy
- Gautam Sarath** Grain, Forage, and Bioenergy Research Unit
USDA-ARS and Department of Agronomy and Horticulture
University of Nebraska
Lincoln, NE
- Kai-Uwe Schwarz** Julius Kühn-Institute (JKI)
Federal Research Centre for Cultivated Plants
Bundesallee, Braunschweig, Germany
- Gancho Slavov** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- Tim Springer** Southern Plains Range Research Station
Woodward, OK
- T. R. Stefaniak** Department of Soil and Crop Sciences
Texas A&M University
College Station, TX
- J. Ryan Stewart** Brigham Young University
Provo, UT
- Charlie Rodgers** Ceres, Inc.
Somerville, TX
- Christian Tobias** USDA-ARS
Western Regional Research Center
Albany, CA
- John Valentine** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK
- Zeng-Yu Wang** Forage Improvement Division
The Samuel Roberts Noble Foundation
Ardmore, OK
- Richard Webster** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK

- Yanqi Wu** Department of Plant and Soil Science
Oklahoma State University
Stillwater, OK
- Toshihiko Yamada** Field Science Center for Northern Biosphere
Hokkaido University
Kita-ku, Sapporo, Hokkaido, Japan
- Sue Youell** Institute of Biological, Environmental and Rural Sciences
(IBERS)
Aberystwyth University
Gogerddan, Aberystwyth, Ceredigion, Wales, UK

Preface

The world energy use grew by 39% from 1990 to 2008. It is estimated that the global demand for energy will increase by at least 50% over the next 20 years. Energy consumption growth of several developing nations remains vigorous. Hydrocarbons, petroleum, coal, and natural gas are now the chief sources of energy. All are finite resources and their natural reserves are depleting every day. In addition, during their conversion and use several greenhouse gases are emitted with a potential for climatic warming.

Bioenergy, both biofuels and biopower, produced from renewable sources are sustainable alternatives to hydrocarbons. Bioenergy use has the potential to lower greenhouse gas emissions, boost rural economy, and ensure energy security. Interest in bioenergy began in early 20th century but it was reinforced in the recent decades. Biopower includes co-firing bioenergy feedstocks with coal to reduce problem emissions. Due to government incentives during 1995–2005, commercial scale biofuels, mainly ethanol, became available in the European Union, UK, USA, Brazil, and many other countries around the world. Most of the biofuels are derived from corn grain, sugarcane, and vegetable oil feedstocks thus creating a food versus fuel controversy. Second-generation biofuels are now being made from nonfood, lignocellulosic materials such as municipal waste and wood chips, along with dedicated crops such as switchgrass and *Miscanthus*.

Plant breeding is critical for crop improvement. Due to intensive breeding efforts in both public and private sectors average maize grain yield has increased by 745% since 1930. Several of the dedicated feedstock crops, for example, switchgrass and *Miscanthus*, are only recently removed from the wild and need serious breeding efforts for improvement. Increased biomass yield and improved quality through breeding efforts can make feedstock more economical and attractive. This book on *Bioenergy Feedstocks: Breeding and Genetics* should greatly contribute to these breeding efforts.

We are grateful to John Wiley & Sons, Inc. for their prudence and supporting us in publishing this book. Contribution from many prominent scientists on bioenergy research has greatly enhanced this publication. We extend our sincere appreciation to all the chapter contributors for their invaluable contribution. We also appreciate the efforts of all who directly or indirectly supported our endeavor. We sincerely believe that this book will be a useful reference for cultivar improvement of lignocellulosic biomass feedstock crops.

*Malay C. Saha
Hem S. Bhandari
Joseph H. Bouton*

Chapter 1

Introduction

Joseph H. Bouton¹, Hem S. Bhandari², and Malay C. Saha³

¹*Former Director and Senior Vice President, Forage Improvement Division, The Samuel Roberts Noble Foundation, Ardmore, OK 73401 and Emeritus Professor, Crop and Soil Sciences, University of Georgia, Athens, GA 30602, USA*

²*Department of Plant Sciences, University of Tennessee, Knoxville, TN37996, USA*

³*Forage Improvement Division, The Samuel Roberts Noble Foundation, Ardmore, OK 73401, USA*

By most estimates, world population growth has more than tripled during the past 100 years, going from approximately 2–7 billion persons (Anonymous, 2012). To sustain the economies needed to support this type of unprecedented population growth, readily available, cheap, scalable, and efficient energy sources were required. These sources turned out to be hydrocarbons, both oil and coal, and after the Second World War, nuclear power. Heavy hydrocarbon use resulted in their depletion and increased cost and a concurrent increase in environmental problems due to gas emissions. Although “clean” as far as gas emissions, nuclear power has its own problems associated with safety and disposal of its highly toxic waste products. Therefore, alternative energy sources such as wind, solar, and bioenergy that are capable of offsetting some of the hydrocarbons and nuclear use and mitigating their environmental problems are now being investigated and, in some cases, implemented on a commercial scale.

Lignocellulosic feedstocks derived from plant biomass emerged as a sustainable and renewable energy source that underpins the bioenergy industry (McLaughlin, 1992; Sanderson et al., 2006). Bioenergy, both biopower and biofuels, could contribute significantly to meet growing energy demand while mitigating the environmental problems. The Energy Independence and Security Act RFS2 in the United States mandates that annual biofuels’ use increase to 36 billion gallons per year by 2022, of which 21 billion gallons should come from advanced biofuels (EISA, 2007). Waste products, both agricultural and forest residues, are obvious choices as base feedstocks; however, it is the use of “dedicated” energy crops where the ability to achieve the billion tons of biomass USA goals will be realized (Perlack et al., 2005). Several plant species such as switchgrass, Miscanthus, corn fodder, sorghum, energy canes, and other grass and legume species have demonstrated tremendous potential for use as dedicated bioenergy feedstocks especially for the production of advanced biofuels. Their adaptation patterns along most agro-ecological gradients also offer options for optimizing a crop species mix for any bioenergy feedstock production system.

1.1 Historical Development

The concept of bioenergy is not new. Early human civilization witnessed energy potential of plant biomass and used it in cooking and as a source of light. By 1912, Rudolf Diesel demonstrated that diesel obtained from plant biomass can be used in automobile operation (Korbitz, 1999). The shortage of crude oil during the 1970s reinforced the world's motivation toward plant biomass as alternative energy source. In Brazil, use of ethanol to power automobile dates back to the late 1920s. Brazil's National Alcohol Program under government funding was launched in 1975 to promote ethanol production from sugarcane. In 2007, Brazil produced more than 16 billion liters of ethanol (Goldemberg, 2007).

In the United States, during the past decade, billions of dollars were invested annually by the federal and state governments, venture capitalists, and major private companies for the development of new technology to convert feedstock species into renewable biofuels. Major breakthroughs have happened during the past few years and the biofuel production increased significantly. Significant improvements have also noticed on conversion technologies thus moving the biofuel from pilot scale to near-commercial scale.

At present, biofuels are produced from corn grain, sugar cane, and vegetable oil. In the United States, corn is the main feedstock used to produce ethanol. In 2010, corn-based ethanol production was about 50 billion liters (USDOE, 2011). With the increasing world's food demand there is serious economic (animal feed costs are rising) and even ethical concern with using corn grain in ethanol production. In the mid-1985s, U.S. Department of Energy (DOE) Herbaceous Energy Crops Program (HECP), coordinated by Oakridge National Laboratory (ORNL), funded research to identify potential herbaceous species as potential bioenergy feedstock. Over 30 plant herbaceous crop species including grasses and legumes were studied, and consequently switchgrass was chosen as the "model bioenergy species" (McLaughlin and Kszos, 2005). Under optimum conditions, switchgrass demonstrated annual biomass yield as high as 24 Mg ha⁻¹, and each ton of biomass can produce about 380 L of ethanol (Schmer et al., 2008). Carbon sequestration by 5-year-old switchgrass stand can add 2.4 Mg C ha⁻¹ year⁻¹ for 10,000 Mg ha⁻¹ of soil mass (Schmer et al., 2011). Other plant species with high bioenergy potential include *Miscanthus*, corn fodder, sorghum, sugarcane, prairie cordgrass, bluestems, eastern gamagrass, and alfalfa. *Miscanthus* hybrids have the potential to produce high biomass and can make a significant contribution to biofuel production and to the mitigation of climate change. Plant breeding will play an important role in improving the genetic potential of these species, as well as other potential species, and make them suitable as bioenergy feedstock.

1.2 Cultivar Development

Genetic improvement of plant species targeting biomass feedstock production, particularly the dedicated energy crops such as switchgrass and *Miscanthus*, is in a very early stage, posing both challenges and opportunities for genetic improvement. The current emphasis of most biomass feedstock cultivar development research is based on biomass yield. Due to extensive breeding efforts, maize grain yield has increased 745% from 1930 to the present (USDA-NASS, 2011). Biomass yield per unit of land is a function of many traits; thus plant breeders also have to address problems related to establishment, seed shattering, and resistance to abiotic/biotic stresses. Equally important is improvement in feedstock quality for sustainable bioeconomy.

Research is still evolving on processes to convert biomass to bioenergy/biofuel that will dictate the quality targets of dedicated bioenergy crops. One likely scenario is that both enzymatic and thermochemical conversion technologies will be required depending on the biomass feedstock availability and the targeted bioenergy end product.

1.3 Breeding Approach

The fundamentals of feedstock cultivar development will be the same as ones that have been successfully employed in several agricultural crops for thousands of years. Most of the potential bioenergy crops are outcrossing polyploids and great genetic diversity exists both within and among populations. This reinforces the potential for genetic improvement of these crops. Most of the named switchgrass cultivars were developed only by seed increases of desirable plants identified from the wild or selected through two or three generations under cultivation (Casler et al., 2007). The improvement of quantitative traits will require several cycles of selection (Bouton, 2008). The traits that are qualitatively inherited can be improved rapidly. Exploitation of heterosis would require identification of genes involved in heterosis and development of heterotic pools, similar to the one that was followed in hybrid breeding in maize. Different crop species would need different plant breeding methodologies depending on their mode of reproduction, ploidy systems, and germplasm availability. For example, corn has a well-developed hybrid production system using inbred lines, which may not be directly applicable to crops like switchgrass that has nearly 100% self-incompatibility. Some species of *Miscanthus* and sugarcane that do not produce seeds require a different approach. The hundreds of years of experience gained in the development of modern cultivars of food and other agricultural crops can directly benefit the cultivar development research of bioenergy crops.

1.4 Molecular Tools

Rapid development in high-throughput genotyping, genotyping based on sequencing, and computational biology continues to shape modern plant breeding into a new approach called “molecular breeding.” Rapid discoveries of DNA-based markers at significantly reduced cost have impacted cultivar development methodologies in the recent years. Advances in molecular biological research have uncovered several plant biological functions and enhanced the understanding of gene function at the molecular level (Bouton, 2008). Rapidly growing genome, transcriptome, proteome, and metabolome resources of several important biofuel crops can speed the process of feedstock development which can lead to improved economics of renewable bioenergy production. Lignin polymer is found to be interfering with enzymatic digestion of lignocellulosic biomass necessitating the pretreatment of biomass feedstock, making biofuel production an economic challenge (Dien et al., 2011). However, plant biologists have been able to characterize and modify lignin pathway and produce low lignin plants by silencing genes involved in lignin pathway (Dien et al., 2011; Fu et al., 2011). Transgenic technologies have also enabled plant breeders to look beyond target species for genes conferring desirable traits, but current regulatory aspects could curtail gains from transgenics, especially for bioenergy crops, without deregulation reforms that better balance both risk and benefit (Strauss et al., 2010).

1.5 Future Outlook

Changing climates as seen by frequent unprecedented drought cycles have become a serious challenge in the recent decades. This will require an “adjustment philosophy” in that breeding strategies will need to continually adjust trait targets for greater stress extremes with programs concentrating on stress tolerances growing in importance (Bouton, 2010). As biomass feedstock production scales up to a commercial level, there will also be a significant shift in agricultural landscapes, leading to occurrence of new pest and diseases specific to the feedstock species. Exploration and exploitation of microbial endophytes implicated in protection of plants from a broad range of biotic and abiotic stresses are important areas for future research (Ghimire et al., 2011). Bioenergy crop breeders should therefore take proactive action to integrate all conventional and modern tools into their cultivar development research.

There are government policy issues that may assist the growth of bioenergy industry. However, these are political issues not within the scope of this book and will need to be hashed out at that level. But one thing is certain, bioenergy cultivar development research will benefit by always striving for a cost-effective product that competes in the free market with hydrocarbons and nuclear power. This should become more possible by leveraging facilities/resources established for traditional agricultural crops and implementation of regional/national/international collaborations between institutions involved in bioenergy feedstock research. Finally, sharing germplasms between participating institutes would help maintain genetic diversity of the breeding pools needed for long-term use.

References

- Anonymous. Environment, 2012. <http://one-simple-idea.com/Environment1.htm>. Accessed on 3 September 2012.
- Bouton J. Improvement of switchgrass as a bioenergy crop. In: Vermerris W (ed.) *Genetic Improvement of Bioenergy Crops*, 2008, pp. 295–308. Springer Science, New York.
- Bouton JH. Future developments and uses. In: Boller B, Posselt UK, Veronesi F (eds) *Handbook of Plant Breeding*, Vol. 5. Fodder Crops and Amenity Grasses, 2010, pp. 201–209. Springer, New York, Dordrecht, Heidelberg, London.
- Casler MD, Stendal CA, Kapich L, Vogel KP. Genetic diversity, plant adaptation regions, and gene pools for switchgrass. *Crop Sci*, 2007; 47: 2261–2273.
- Dien BS, Miller DJ, Hector RE, Dixon RA, Chen F, McCaslin M, Reisen P, Sarath G, Cotta MA. Enhancing alfalfa conversion efficiencies for sugar recovery and ethanol production by altering lignin composition. *Bioresource Technol*, 2011; 102: 6479–6486. doi:10.1016/j.biortech.2011.03.022.
- The Energy Independence and Security Act [EISA], 2007. <http://www.gpo.gov/fdsys/pkg/PLAW-110publ140/pdf/PLAW-110publ140.pdf>. Accessed on 4 December 2012.
- Fu C, Xiao X, Xi Y, Ge Y, Chen F, Bouton J, Dixon RA, Wang Z-Y. Downregulation of cinnamyl alcohol dehydrogenase (CAD) leads to improved saccharification efficiency in switchgrass. *Bioenerg Res*, 2011; 4: 153–164. doi:10.1007/s12155-010-9109-z.
- Ghimire SR, Charlton ND, Bell JD, Krishnamurthy YL, Craven KD. Biodiversity of fungal endophyte communities inhabiting switchgrass (*Panicum virgatum* L.) growing in the native tallgrass prairie of northern Oklahoma. *Fungal Divers*, 2011; 47: 19–27. doi:10.1007/s13225-010-0085-6.

- Goldemberg J. Ethanol for a sustainable energy future. *Science*, 2007; 315: 808–810.
- Korbitz W. Biodiesel production in Europe and North America, and encouraging prospect. *Renew Energ*, 1999; 16: 1078–1083.
- McLaughlin SB. New switchgrass biofuels research program for the Southeast. In: *Proceedings Ann Auto Tech Dev Contract Coord Mtng*, 1992, pp. 111–115. Dearborn, MI.
- McLaughlin SB, Kszos LA. Development of switchgrass (*Panicum virgatum*) as a bioenergy feedstock in the United States. *Biomass Bioenerg*, 2005; 28: 515–535.
- Perlack RD, Wright LL, Turnhollow AF, Graham RL, Stokes BJ, Erbach DC. Biomass as a feedstock for a bioenergy and bioproducts industry: the technical feasibility of a billion ton supply, 2005. http://feedstockreview.ornl.gov/pdf/billion_ton_vision.pdf. Accessed on 3 September 2012.
- Sanderson MA, Adler PR, Boateng AA, Casler MD, Sarath G. Switchgrass as a biofuels feedstock in the USA. *Can J Plant Sci*, 2006; 86: 1315–1325.
- Schmer MR, Liebig MA, Vogel KP, Mitchell RB. Field-scale soil property changes under switchgrass managed for bioenergy. *Global Change Biol Bioenerg*, 2011; 3: 439–448.
- Schmer MR, Vogel KP, Mitchell RB, Perrin RK. Net energy of cellulosic ethanol from switchgrass. *Proc Natl Acad Sci U S A*, 2008; 105: 464–469.
- Strauss SH, Kershen DL, Bouton JH, Redick TP, Tan H, Sedjo RA. Far-reaching deleterious impacts of regulations on research and environmental studies of recombinant DNA-modified perennial biofuel crops in the United States. *BioScience*, 2010; 60: 729–741.
- USDOE. *US Billion-Ton Updates: Biomass Supply for a Bioenergy and Bioproducts Industry*. R.D. Perlack and B.J. Stokes (Leads), ORNL/TM-2011/224, 2011, p. 227. Oak Ridge National Laboratory, Oak Ridge, TN. http://www1.eere.energy.gov/biomass/pdfs/billion_ton_update.pdf. Accessed on 19 March 2013.
- USDA-NASS. Quick Stats: Agricultural Statistics Data Base, 2011. <http://quickstats.nass.usda.gov/results/D21A4E3A-A14A-3845-B2C7-7A62041B1648>. Accessed on 4 December 2012.

Chapter 2

Switchgrass Genetics and Breeding Challenges

Laura Bartley¹, Yanqi Wu², Aaron Saathoff³, and Gautam Sarath³

¹*Department of Microbiology and Plant Biology, University of Oklahoma, Norman, OK 73019, USA*

²*Department of Plant and Soil Science, Oklahoma State University, Stillwater, OK 74078, USA*

³*Grain, Forage, and Bioenergy Research Unit: USDA-ARS and Department of Agronomy and Horticulture, University of Nebraska, Lincoln, NE 68583, USA*

2.1 Introduction

Liquid biofuel production from biomass has the potential to reduce greenhouse gas emissions from transportation and dependence on fossil fuels extracted from politically volatile regions (Somerville, 2007; Bartley and Ronald, 2009; Vega-Sanchez and Ronald, 2010). Cultivated grasses are the most abundant sustainable class of biomass that can be produced in the United States (~57%, Perlack et al., 2005). Switchgrass (*Panicum virgatum* L.), in particular, is an attractive native species for development as a bioenergy crop given that largely unimproved varieties exhibit large biomass yield (up to 36.7 Mg ha⁻¹) and marked stress tolerance (Figure 2.1; Thomason et al., 2004; McLaughlin and Adams Kszos, 2005; Bouton, 2007). Even with typical, lower-yielding marginal land (5–11 Mg ha⁻¹), energy and emission measurements for switchgrass production give an approximately 5-fold net energy yield (output:input) and an approximately 10-fold reduction in greenhouse gas emissions compared with gasoline (Schmer et al., 2008). In order to realize greater benefits from the production of lignocellulosic fuels, there is an enormous need to apply various breeding methods and tools toward switchgrass improvement. Below, we outline switchgrass energy crop breeding goals and, in subsequent sections, provide an overview of the basic biology and genetic characteristics of switchgrass. We then discuss experiences and challenges related to switchgrass conventional and molecular breeding.

Biomass *yield* and *quality* are the two general classes of targets for genetic improvement of bioenergy crops. Selection of switchgrass for high biomass production is ongoing (Vogel et al., 2010). Recently released cultivars “BoMaster,” “Cimarron,” and “Colony” produce higher biomass yields than the current best commercial cultivar “Alamo” and are primarily targeted for cellulosic feedstock production (Burns et al., 2008a, 2008b, 2010; Wu and Taliaferro, 2009). Similarly, high-performing replacement proprietary cultivars for old standards such as Alamo, “Kanlow,” and “Cave-in-Rock” are currently sold in commercial



Figure 2.1. A seed production field of the lowland switchgrass cultivar “Cimarron” in September, toward the end of the growing season, near Perkins, OK.

bioenergy seed trade as EG1101, EG1102, and EG2101, respectively (<http://www.bladeenergy.com/SwitchProducts.aspx>).

Abiotic and biotic stress tolerance and improved agronomic characteristics, such as reduced seed dormancy (Burson et al., 2009), are important for establishing and obtaining consistent biomass production. In terms of biomass quality, the goals for the two current biomass to biofuel conversion platforms, biochemical and thermochemical, are roughly opposite (Figure 2.2). For biochemical conversion methods that mostly produce alcohol fuels, the quality goal is to optimize the quantity of sugar that can be obtained from the biomass. For




Figure 2.2. Potential applications and consequences of changing switchgrass biomass quality. CHO, carbohydrate.