


NEUROLOGY IN PRACTICE

Series editors **Robert A. Gross & Jonathan W. Mink**

Non-Parkinsonian Movement Disorders



Edited by

Deborah A. Hall and Brandon R. Barton

WILEY Blackwell

Non-Parkinsonian Movement Disorders

NEUROLOGY IN PRACTICE:

SERIES EDITORS: ROBERT A. GROSS, DEPARTMENT OF NEUROLOGY,
UNIVERSITY OF ROCHESTER MEDICAL CENTER, ROCHESTER, NY, USA

JONATHAN W. MINK, DEPARTMENT OF NEUROLOGY,
UNIVERSITY OF ROCHESTER MEDICAL CENTER, ROCHESTER, NY, USA

Non-Parkinsonian Movement Disorders

EDITED BY

Deborah A. Hall, MD, PhD

Department of Neurological Sciences

Section of Movement Disorders

Rush University Medical Center, Chicago

Illinois, USA

Brandon R. Barton, MD, MS

Department of Neurological Sciences

Section of Movement Disorders

Rush University Medical Center;

Neurology Section

Jesse Brown VA Medical Center

Chicago, Illinois, USA

WILEY Blackwell

This edition first published 2017, © 2017 by John Wiley & Sons, Ltd.

Registered Office

John Wiley & Sons, Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial Offices

9600 Garsington Road, Oxford, OX4 2DQ, UK

The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

111 River Street, Hoboken, NJ 07030-5774, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell

The right of the author to be identified as the author of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting a specific method, diagnosis, or treatment by health science practitioners for any particular patient. The publisher and the author make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of fitness for a particular purpose. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. Readers should consult with a specialist where appropriate. The fact that an organization or Website is referred to in this work as a citation and/or a potential source of further information does not mean that the author or the publisher endorses the information the organization or Website may provide or recommendations it may make. Further, readers should be aware that Internet Websites listed in this work may have changed or disappeared between when this work was written and when it is read. No warranty may be created or extended by any promotional statements for this work. Neither the publisher nor the author shall be liable for any damages arising herefrom.

Library of Congress Cataloging-in-Publication Data

Names: Hall, Deborah A., editor. | Barton, Brandon R., editor.

Title: Non-Parkinsonian movement disorders / edited by Deborah A. Hall and Brandon R. Barton.

Description: Chichester, West Sussex ; Hoboken, NJ : John Wiley & Sons Inc., 2016. |

Includes bibliographical references and index.

Identifiers: LCCN 2016023027 | ISBN 9781118473924 (Paperback) | ISBN 9781118474068 (Adobe PDF) |

ISBN 9781118474051 (epub)

Subjects: | MESH: Movement Disorders--diagnosis | Diagnosis, Differential | Diagnostic Techniques, Neurological

Classification: LCC RC376.5 | NLM WL 390 | DDC 616.8/3--dc23

LC record available at <https://lcn.loc.gov/2016023027>

A catalogue record for this book is available from the British Library.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Background cover image: iStockphoto.com

Set in 8.75/11.75pt Utopia by SPi Global, Pondicherry, India

Contents

List of Contributors	vii
Series Foreword	ix
Foreword	x
About the Companion Website	xi

1 Approach to Movement Disorders	1
<i>Deborah A. Hall and Brandon R. Barton</i>	

PART 1: HYPOKINETIC

2 Hypokinetic (Non-Parkinsonian) Movement Disorders	7
<i>Shyamal H. Mehta and Alberto J. Espay</i>	

PART 2: HYPERKINETIC

3 Common Types of Tremor	17
<i>Jeff Kraakevik and Bernadette Schoneburg</i>	

4 Myoclonus	26
<i>Daniel Burdick and Pinky Agarwal</i>	

5 Tics and Tourette Syndrome	35
<i>David Shprecher</i>	

6 Chorea, Athetosis, and Ballism	43
<i>Rohit Dhall</i>	

7 Dystonia	57
<i>Lauren Schrock, Tao Xie and Brandon R. Barton</i>	

8 Ataxia	68
<i>Samantha Holden and Deborah A. Hall</i>	

9 Restless Leg Syndrome	79
<i>Olga Klepitskaya</i>	

10 Hemifacial Spasm and Other Facial Movement Disorders	89
<i>Tao Xie, Ifeoma Nwaneri and Un Jung Kang</i>	

11 Periodic Limb Movements of Sleep and REM Sleep Behavior Disorder	99
<i>Aleksandar Videnovic</i>	

12 Stereotypies	106
<i>Michael Rotstein</i>	
13 Paroxysmal Movement Disorders	114
<i>Christina L. Vaughan</i>	
14 Uncommon Movement Disorders and Movement Disorder Mimics	121
<i>Nina Browner</i>	
PART 3: OTHER DISEASE SYNDROMES	
15 Tardive Syndromes	133
<i>Stephanie Lessig</i>	
16 Heavy Metal Accumulation Diseases	139
<i>Khashayar Dashtipour and Janice Fuentes</i>	
17 ICU Intensive Care Unit Movement Disorder Emergencies	150
<i>Florence C. F. Chang and Steven J. Frucht</i>	
18 Functional or Psychogenic Movement Disorders	161
<i>S. Elizabeth Zauber</i>	
PART 4: ADDITIONAL RESOURCES	
19 Genetics of Movement Disorders	171
<i>Deborah A. Hall</i>	
20 Neuroimaging Finding in Movement Disorders	175
<i>Kathleen L. Poston</i>	
21 Clinical Rating Scales in Movement Disorders	185
<i>Padmaja Vittal and Brandon R. Barton</i>	
22 Videotaping Suggestions for Movement Disorders	190
<i>Gian Pal and Deborah A. Hall</i>	
Index	195

List of Contributors

Pinky Agarwal MD

Movement Disorders Center
Evergreen Health Neuroscience Institute
Kirkland, WA

Brandon R. Barton MD, MS

Department of Neurological Sciences
Section of Movement Disorders
Rush University Medical Center;
Neurology Section
Jesse Brown VA Medical Center
Chicago, IL

Nina Browner MD

National Parkinson Foundation Center of
Excellence
University of North Carolina
Chapel Hill, NC

Daniel Burdick MD

Movement Disorders Center
Evergreen Health Neuroscience Institute
Kirkland, WA

Florence C. F. Chang MBBS, FRACP

Neurology Department
Westmead Hospital
Wentworthville, NSW

Khashayar Dashtipour MD, PhD

Department of Neurology, Movement Disorders
Loma Linda University School of Medicine
Loma Linda, CA

Rohit Dhall MBBS, MSPH

Parkinson's Institute and Clinical Center
Sunnyvale, CA

Alberto J. Espay MD, MSc

Gardner Center for Parkinson's Disease and
Movement Disorders
Department of Neurology
University of Cincinnati
Cincinnati, OH

Steven J. Frucht MD

Mount Sinai Medical Center
NY

Janice Fuentes MD

Department of Neurology, Movement Disorders
Loma Linda University School of Medicine
Loma Linda, CA

Deborah A. Hall MD, PhD

Department of Neurological Sciences
Section of Movement Disorders
Rush University Medical Center
Chicago, IL

Samantha Holden MD

University of Colorado School of Medicine,
Aurora, Colorado, USA

Un Jung Kang MD

Department of Neurology
Columbia University Medical Center
NY

Olga Klepitskaya MD

Department of Neurology
School of Medicine, University of Colorado
Aurora, CO

Jeff Kraakevik MD

Assistant Professor
Oregon Health and Science University
Portland, OR

Stephanie Lessig MD

Department of Neurosciences
University of California San Diego
La Jolla, CA

Shyamal H. Mehta MD, PhD

Department of Neurology
Movement Disorders Division
Mayo Clinic
Phoenix, AZ

Ifeoma Nwaneri MD

Reston Hospital Center
Springfield, VA

Gian Pal MD, MS

Department of Neurological Sciences
Section of Movement Disorders
Rush University Medical Center
Chicago, IL

Kathleen L. Poston MD, MS

Department of Neurology and Neurological
Sciences
Stanford University
Stanford, CA

Michael Rotstein MD

Tel Aviv Sourasky Medical Center
Tel Aviv, Israel

Bernadette Schoneburg MD

Northshore Medical Group
Glenview, IL

Lauren Schrock MD

University of Utah
Department of Neurology
Salt Lake City, UT

David Shprecher DO, MSc

Cleo Roberts Center, Banner Sun Health
Research Institute in Sun City, AZ

Christina L. Vaughan MD, MS

Hospice and Palliative Medicine Program
University of California San Diego/Scripps Health
La Jolla, CA

Aleksandar Videnovic MD

Department of Neurology
Massachusetts General Hospital
Harvard Medical School
Boston, MA

Padmaja Vittal MD, MS

Northwestern Medicine Regional Medical Group
Winfield, IL

Tao Xie MD, PhD

Department of Neurology
University of Chicago Medical Center
Chicago, IL

S. Elizabeth Zauber MD

Department of Neurology
Indiana University School of Medicine
Indianapolis, IN

Series Foreword

The genesis for this book series started with the proposition that, increasingly, physicians want direct, useful information to help them in clinical care. Textbooks, while comprehensive, are useful primarily as detailed reference works but pose challenges for uses at the point of care. By contrast, more outline-type references often leave out the “hows and whys”—pathophysiology, pharmacology—that form the basis of management decisions. Our goal for this series is to present books, covering most areas of neurology, that provide enough background information to allow the reader to feel comfortable, but not so much as to be overwhelming, and to associate that with practical advice from experts about care, combining the growing evidence base with best practices.

Our series will encompass various aspects of neurology, with topics and the specific content chosen to be accessible and useful.

Chapters cover critical information that will inform the reader of the disease processes and mechanisms as a prelude to treatment planning. Algorithms and guidelines are presented, when appropriate. “Tips and Tricks” boxes provide expert suggestions, while other boxes present cautions and warnings to avoid pitfalls. Finally, we provide “Science Revisited” sections that review the most important and relevant science background material, and references and further reading sections that guide the reader to additional material.

Our thanks, appreciation, and respect go out to our editors and their contributors, who conceived and refined the content for each volume, assuring a high-quality, practical approach to neurological conditions and their treatment.

Our thanks also go to our mentors and students (past, present, and future), who have challenged and delighted us; to our book editors and their contributors, who were willing to take on additional work for an educational goal; and to our original publisher, Martin Sugden, for his ideas and support, for wonderful discussions and commiseration over baseball and soccer teams that might not quite have lived up to expectations. And thanks, too, to Claire Bonnett, our current publisher, for her efforts to bring this volume forward.

This volume represents the end of our series. As readers will recognize, neurology encompasses far more than we have presented; still, we hope that the high points encompassed by these books will serve well.

We have dedicated the series to Marsha, Jake, and Dan, and to Janet, Laura, and David. And also to Steven R. Schwid, MD, our friend and colleague, whose ideas helped shape this project and whose humor brightened our lives; but he could not complete this goal with us. Our thanks to them are undiminished.

Robert A. Gross
Jonathan W. Mink
Rochester, NY, USA

Foreword

Non-Parkinsonian Movement Disorders, edited by my colleagues, Drs. Brandon Barton and Deborah Hall, is a new entry in the larger series, *Neurology in Practice* and an immediate compendium of the *Parkinsonian Movement Disorders*. The topics covered in this volume provide the practicing neurologist, psychiatrist, and primary care health professional with expert reviews that cover both hypokinetic and hyperkinetic disorders. Hypokinetic disorders discussed include stiff-person syndrome, catatonia, and catalepsy as well as a variety of stiff-muscle conditions. For the hyperkinetic disorders, the editors have assembled a group of expert authors to cover tremors, myoclonus, tics, chorea, dystonia, and involuntary movements due to toxins and drugs. Importantly, because neurological disorders can be both out-patient and in-patient consultations, a chapter on ICU movement disorders emergencies is included, and an important chapter on the knotty and complex problem of psychogenic movement disorders that focuses on a variety of functional movements, both consciously and unconsciously generated.

Each presentation is anchored in very practical descriptions of phenomenology, key clinical information from the history and neurological examination that guide the physician to the correct diagnosis and treatment options. The text is enriched with tables and figures and a number of unique learning tools not found in other books on this topic. These tools include special boxed “Tips and Tricks” and “Caution” warnings that can help prevent errors. Besides the focus on practical clinical medicine, the authors provide two special highlights in each chapter, “Science Revisited” to remind clinicians of the scientific anchors related to the disorders and “Evidence at a Glance” where clinical trial evidence-based review information is provided. All these

special additions allow a reader to study the full text, but also to retrieve rapidly needed key points.

With a long career devoted to the treatment of movement disorders, research, and education, I laud the editors and their recruited authors in providing the medical community with an accessible and accurate book with a unique format. In a busy environment, this text serves as a very solid neurological work with the essentials delivered in a succinct and highly readable format.

Because a movement disorders diagnosis always starts with accurate visual identification, a very strong advantage of this text is the video material that accompanies each chapter. The elegant examples assembled are well synchronized with the text materials and allow the reader to study very good examples of the disorders under consideration. I suggest that readers start a chapter with a short examination of the video materials, so as to be clear on the type of movement under discussion, then read the text, finally returning to the videos for a focused re-examination of the details of the carefully prepared examples. With this emphasis on the key role of expert visual recognition in movement disorders, the words of the celebrated nineteenth-century neurologist, Jean-Martin Charcot, resonate, and I offer this citation as the reader embarks on the very pleasant road of reading this volume:

Let someone say of a doctor that he really knows his physiology or anatomy, that he is dynamic—these are not real compliments; but if you say he is an observer, a man who knows how to see, this is perhaps the greatest compliment one can make (Charcot, 1888, *Leçons du mardi*).

Christopher G. Goetz, MD, Chicago, IL, USA,
2016

About the Companion Website

This book is accompanied by a companion website:



www.wiley.com/go/hall/non-parkinsonian_movement_disorders

The website includes:

Videos

Approach to Movement Disorders

Deborah A. Hall, MD, PhD¹ and Brandon R. Barton, MD, MS^{1,2}

¹Department of Neurological Sciences, Section of Movement Disorders, Rush University Medical Center, Chicago, Illinois, USA

²Department of Neurological Sciences, Rush University Medical Center; Neurology Section, Jesse Brown VA Medical Center, Chicago, Illinois, USA

Introduction

Patients with movement disorders typically present with a change in their overall pattern of movements: this may represent an increase of movement (hyperkinetic), decrease (hypo- or akinetic), uncoordinated movement (ataxia), or a combination of the aforementioned. The initial task is to properly categorize the appearance or “phenomenology” of the movement disorder, as this is the essential step to guide the clinician in developing a differential diagnosis and treatment plan. Given recent advances in neurology, the majority of movement disorder patients are candidates for treatment, such as medication, physical therapy, or surgical interventions.

The first part of this book provides a short chapter on non-parkinsonian hypokinetic movement disorders; parkinsonian disorders are covered in another volume in this series. The second part includes hyperkinetic disorders. Part three includes various syndromes that do not fit into the other categories or that overlap between categories. Broader chapters in part four, on genetics, neuroimaging, rating scales, and videotaping suggestions, are intended to serve as clinician resources.

This introductory chapter provides an approach that will facilitate the evaluation of a movement disorder patient. The phenomenological categorization of the most common movement disorders falls into seven major categories: parkinsonism, tremor, dystonia, myoclonus, chorea, ataxia, and

tics. Most of the commonly encountered disorders can be classified into one of these categories, but given the breadth of the diseases in the field, there are many unusual or rare types of movement that may not be easily categorized or may be consistent with more than one phenomenological category. A thorough history and examination are essential to defining the phenomenology. Home videotapes of the patient may also be useful if the movements are intermittent, variable, or not seen clearly in the office. Laboratory testing and imaging are necessary in some movement disorders, but are less helpful in many circumstances given that the disorders are diagnosed mainly on history and examination.

History

Start by asking *six questions* in the history.

1. Can you describe the movements?

Patients will usually be able to describe a decrease or increase (or both) in their overall movement from baseline, although often hyperkinetic aspects of abnormal movements can overshadow the hypokinetic movements from the patient's perspective. Hypokinetic movement disorders, also termed bradykinesia (slowed movement) or akinesia (loss of movement) are characterized by an overall decrease in the speed or amplitude of movement in any area of the body. Signs and symptoms could include decreased facial expression, slowed speech, reduced

Non-Parkinsonian Movement Disorders, First Edition. Edited by Deborah A. Hall and Brandon R. Barton.

© 2017 John Wiley & Sons, Ltd. Published 2017 by John Wiley & Sons, Ltd.

Companion website: www.wiley.com/go/hall/non-parkinsonian_movement_disorders

dexterity of the extremities, decreased arm swing, and slowed walking speed. Hyperkinetic movement disorders, also generally termed dyskinesia (abnormal movements), are characterized by an increase in baseline movements. Hyperkinetic movement disorders have highly variable manifestations, ranging from increased eye closure to arm flailing to jerking of the legs. Lastly, patients may complain of a change in the character of voluntary movements, such as becoming clumsy or unsteady with walking, which may be seen in ataxic disorders.

Certain features of abnormal movements are very important to elicit in the patient's description. Defining the conditions under which the movement occurs, such as with rest or with action, is necessary for accurate diagnosis and categorization of tremor. An ability to suppress the movement or an increase in the movement with suggestion are features common to tics. Specific triggers of the movements, especially with certain tasks, may be reported in dystonic disorders or paroxysmal movement disorders. Myoclonus can be triggered by startle. Asking about worsening of the disorder or improvement with certain foods or alcohol can narrow the differential diagnosis in forms of dystonia, myoclonus, or tremor disorders. A history of falls, especially the temporal course, is helpful in disorders that affect gait and balance, as falls are seen earlier or more frequently in some disorders as opposed to others.

2. When did the movements start and how have they changed over time?

Most movement disorders are subacute or chronic in nature. An acute onset is less common and may signify a secondary movement disorder related to an underlying inciting event, such as a stroke or medication change. Acute onset of movement disorders at maximal severity is also commonly seen in functional movement disorders, where patients will often present to emergency departments from the start. Most hypokinetic, hyperkinetic, and ataxic movement disorders will slowly worsen over time. Disorders that improve over time are less common; for example, tic disorders will typically improve from childhood into adolescence and adulthood. Static movement disorders may occur with birth injury or some dystonic disorders.

3. Are the movements continuous or intermittent?

Although many movement disorders start out as intermittent or suppressible, they tend to become

more continuous or constant when they progress over time. The rest tremor seen in parkinsonian disorders is a classic example, where the tremor starts intermittently in a limb before becoming more regular and spreading to other limbs. Early on, this type of tremor can be sometimes voluntarily suppressed or decreased with movement, but later the tremor is continuous. Episodic movement disorders are much less common. Paroxysmal disorders, which are typically choreic or dystonic in nature, can many times be diagnosed by history alone if specific triggers such as sudden movements cause the disorder to occur. Functional (psychogenic) movement disorders are also frequently episodic. The circumstances under which the movement occurs can be particularly helpful. For example, restless legs syndrome worsens at night when the patient is laying down.

4. Is there a family history?

All modes of inheritance patterns are seen in movement disorders and the genetic basis of these disorders is rapidly being discovered. It is not sufficient to inquire only about the particular movement disorder seen in the patient, since broadening the questioning to other biological family members may yield additional important clues. For example, patients with grandchildren with intellectual disabilities may be at risk for fragile X-associated disorders. Tic patients may have associated diagnoses in the family, such as attention deficit hyperactivity disorder.

5. Are there other medical illnesses?

The majority of movement disorders are restricted to the nervous system, but systemic organ involvement may provide diagnostic clues. For example, patients with underlying cancers may be at risk for paraneoplastic disorders and iron deficiency anemia or diabetes may predispose to restless legs syndrome. The presence of cardiomyopathy is associated with Friedreich ataxia or mitochondrial disorders. Enlargement of visceral organs (spleen, liver) may suggest a lysosomal storage disease.

6. Have the movements been treated in the past and what was the response to treatment?

A response to dopamine medications may facilitate diagnosis of dopa-response dystonia. Paroxysmal movement disorders may be exquisitely responsive to antiepileptic medications. Other substances may improve movements, such as the improvement

of essential tremor, essential myoclonus, and myoclonus-dystonia with alcohol.

Examination

Depending on the movement disorder, abnormal movements may be present in focal or contiguous areas of the body or may be generalized. By determining the location and phenomenology of the movement, most patients can be placed into one of *seven distinct patterns* of abnormal movement.

Parkinsonism

The main features of parkinsonism are tremor at rest, bradykinesia or akinesia, rigidity, loss of postural reflexes, flexed posture, and freezing. Parkinsonism, in particular, Parkinson disease, is the most common disorder seen in movement disorder clinics and is covered by another volume of this series.

Tremor

This pattern is typically rhythmical and oscillatory and may affect more than one body part. Tremor should be classified on examination by the conditions under which it is activated: at rest, with posture, or with action. Tremor may be present in multiple conditions, for example, essential tremor, which is frequently seen with posture and action or intention. Tremors may also be task specific, such as the dystonic tremor of writer's cramp.

Chorea

Choreic movement is random in nature and is purposeless, non-rhythmic, and unsustained. It may appear to flow from one body part to another. Huntington disease is a frequent cause of chorea and manifests with brief, irregular movements. Chorea can be suppressed or camouflaged. It can be accompanied by "negative chorea" or motor impersistence.

Dystonia

In dystonia, agonist and antagonist muscles contract simultaneously causing twisting movements that are frequently sustained. The speed of the movement is variable and when sustained, can lead to abnormal postures and contractures. Dystonia is typically worsened with action, sometimes only occurring with specific actions. It can be classified by location, age of onset, and etiology, and the classification system has recently been revised.

Myoclonus

This pattern consists of brief, sudden, typically irregular jerks from muscle contraction. Myoclonus may be synchronized and triggered by action or startle. Negative myoclonus is caused by inhibition of the muscles, with the classic example being asterixis. Myoclonus can be rhythmic or oscillatory and occur in various parts of the body, either focally or generally.

Tics

Tics are abnormal movements (motor) or sounds (phonic) that are abrupt, usually transient, and can be simple or complex. Tics can vary over time and can be accompanied by an uncomfortable urge or feeling. Tics may be suppressible, although severe tics may be continuous. Gilles de la Tourette syndrome is characterized by the presence of both motor and phonic tics, present for more than one year, with young onset.

Ataxia

Lack of coordination of movement distinguishes ataxia from other movement disorders. The pattern of ataxic movement varies, but may include clumsy limb movements (dysmetria), dysarthria, ataxic eye findings such as abnormal pursuit, and uncoordinated walking. Kinetic tremor can also accompany ataxic signs. Ataxia can be localized to the peripheral or central nervous system so a thorough sensory and vestibular examination is necessary in these patients.

Other patterns of movements

There are several other types of abnormal movements that, despite being distinctly recognizable, do not fit well into the preceding patterns. These include stiff-muscles, akathetic movements, myokymia, paroxysmal dyskinesias, restless legs, and stereotypy. In addition, some movement disorders have more than one pattern of movement, such as in the myoclonus-dystonia disorders. Functional movement disorders frequently do not fit well into the above-described patterns, but caution must be maintained, since many unusual movement disorders can be labeled functional.

Diagnostic testing

Accurate description of the phenomenology of the abnormal movements as a result of the history and examination is the first and most fundamental step

in diagnosis of movement disorders. Additional diagnostic testing is not warranted in many situations, for example, in the classic appearance of Tourette syndrome. However, there are some studies that may enhance or confirm clinical diagnosis. For example, laboratory studies can be useful particularly with tremor. Abnormalities of the thyroid, evidenced by elevated or low thyroid stimulating hormone (TSH), may cause or worsen tremor. Wilson disease, diagnosed by abnormal copper levels (in serum and/or urine), low ceruloplasmin, and the presence of Kaiser-Fleischer rings; should be considered in younger patients who present with bizarre tremors or other unusual movement patterns/combinations.

Genetic testing is available for many movement disorders and is driven by family history, age of the patient, and financial resources. For the more rare movement disorders, such as the inherited ataxias and Huntington disease, it may be the only testing that can give a definitive diagnosis. For individuals who are considering family planning, it may be necessary that genetic testing be accompanied by genetic counseling.

Neurophysiological assessment may be helpful in myoclonus, where myoclonic jerks show brief electromyography (EMG) bursts of 10–50 milliseconds. Rhythmicity in tremor can be demonstrated on EMG, but this is not frequently ordered by clinicians when evaluating a patient with tremor. Electromyography may also be helpful therapeutically in dystonic patients when used in conjunction with botulinum toxin treatment. Nerve conduction studies may be used to evaluate ataxic individuals for sensory abnormalities in peripheral nerves.

Imaging can be valuable in movement disorders that do not fit classic patterns or presentations. The most common movement disorders typically show normal basal ganglia structures on routine imaging, as in essential tremor, and dystonia. However, patients with movement disorders that are localized to one side of the body, that have abrupt

stroke-like onset, or that include ataxia should be imaged with computed tomography or preferably, magnetic resonance imaging. Atrophy of specific structures, such as the striatum in Huntington disease, or the cerebellum in degenerative ataxias, may support the clinical diagnosis. Functional or nuclear medicine imaging is playing an increasingly important role in diagnostics.

Treatment

The majority of treatment options in movement disorders are symptomatic, not curative. However, in a few circumstances, early intervention of treatable forms of movement disorders may be curative or halt the progression of the disease. While rare, such conditions should be considered in patients with particular disease profiles; examples include patients with young onset tremor, dystonia or parkinsonism (Wilson disease), or fluctuating dystonic and parkinsonian features (dopa-responsive dystonia).

The approach described in this chapter offers a straightforward approach to evaluating a movement disorder patient. Questions about the movements, course, family history, medical illnesses, and medication response will help the clinician with the evaluation. Correctly describing the phenomenology is key to narrowing the list of diagnostic possibilities and guides the need for additional testing. The subsequent chapters will fill in the details, and with this framework, the reader will gain an ease of diagnosis and treatment of movement disorders.

Further Readings

- Fahn S, Jankovic J, Hallett M. *Principles and Practice of Movement Disorders*, 2nd ed. Philadelphia: Saunders, 2007.
- Fernandez HH, Rodriguez RL, Skidmore FM, Okun MS. *A Practical Approach to Movement Disorders: Diagnosis, Medical and Surgical Management*. New York: Demos Medical Publishing, 2007.

Part 1

Hypokinetic

Hypokinetic (Non-Parkinsonian) Movement Disorders

Shyamal H. Mehta, MD, PhD¹ and Alberto J. Espay, MD, MSc²

¹Department of Neurology, Movement Disorders Division, Mayo Clinic, Phoenix, Arizona, USA

²Gardner Center for Parkinson's Disease and Movement Disorders, Department of Neurology, University of Cincinnati, Cincinnati, Ohio, USA

Introduction

Movement disorders can be broadly classified into two categories, based on the presence of excess movement or a deficiency of movement. Hyperkinetic movements involve the presence of excessive involuntary movements that may manifest as tremor, chorea-ballism, myoclonus, and tics, among other disorders. Hypokinetic movements show paucity of movement and are described with terms such as bradykinesia (slowness of movement) or akinesia (absence or extreme poverty of movement). The most common form of hypokinetic disorders are the parkinsonian syndromes, including idiopathic Parkinson Disease (PD), atypical parkinsonism, (multiple system atrophy, corticobasal syndrome, progressive supranuclear palsy, etc.), and secondary causes of parkinsonism (midbrain tumors, paraneoplastic disorders, etc.). These topics are covered in another book in this series. A different category of slowness comes from disorders that affect motor function to the extent of rendering it “parkinsonian” but that cannot be explained by traditional impairments in the basal ganglia circuitry. This chapter focuses on the *non-parkinsonian* causes of hypokinetic movement disorders, which may not be traditionally included in the differential diagnosis of parkinsonism.

This chapter aims to highlight some of the important and treatable causes of non-parkinsonian hypokinetic syndromes such as stiff person syndrome, primary lateral sclerosis, catatonia and psychomotor depression, hypothyroidism, and normal pressure hydrocephalus. These, along with other general causes, which can result in paucity or absence of movement, are listed in Table 2.1.

Stiff-person syndrome (SPS)

Moersch and Woltman described “stiff man syndrome” in 1956 in 14 patients who had progressively fluctuating rigidity and painful spasms affecting the muscles of the back and abdomen. An association between SPS and DM was established in the late 1980s; however, only a few of the originally reported patients had concomitant diabetes mellitus (DM). Solimena and colleagues later reported the presence of anti-glutamic acid decarboxylase (anti-GAD) antibodies in patients with SPS and DM. Since GAD is the rate-limiting enzyme for the synthesis of GABA, the major inhibitory neurotransmitter, in the central nervous system, GABAergic depletion at the cortical and spinal interneuronal level is central to the pathogenesis of SPS.

About 80% of SPS patients have a high titer of anti-GAD antibodies detectable in the serum or CSF.

Non-Parkinsonian Movement Disorders, First Edition. Edited by Deborah A. Hall and Brandon R. Barton.

© 2017 John Wiley & Sons, Ltd. Published 2017 by John Wiley & Sons, Ltd.

Companion website: www.wiley.com/go/hall/non-parkinsonian_movement_disorders

Table 2.1 Non-parkinsonian causes of hypokinesia

Stiff-person syndrome and related disorders
• Primary lateral sclerosis (PLM) • Catatonia • Neuromuscular causes: hypothyroidism, Brody syndrome, myotonia • Akinetic mutism
Sequelae of vascular events (affecting anterior cerebral artery distribution)
Structural lesions: tumors, hydrocephalus, traumatic brain injury
Post-infectious: Creutzfeldt–Jakob disease, post-encephalitic parkinsonism
• Functional or psychogenic slowness

GAD is synthesized in the presynaptic GABAergic neurons in both the central nervous system and in the islet of Langerhans β -cells of the pancreas, hence the association with DM. There are two GAD isoforms—GAD65 and GAD67—but it is GAD65 that has been implicated in both SPS and DM. The GAD antibodies recognize different regions (epitopes) of the GAD molecule. The GAD antibodies in type 1 DM recognize the carboxy-terminal end or the center of the GAD molecule, while in SPS they recognize the amino-terminal fragment of GAD. Even though GAD antibodies are the most common antibodies associated with SPS, other proteins, both pre- and post-synaptic, in the GABAergic neuron have been implicated in the etiology of SPS or its variants (Box 2.1).

Clinical features

SPS disease is sporadic in nature, affecting women more often than men (in a recent series ~70% patients were women). Although the age of onset is variable, most of the afflicted adults are between 29 to 59 years of age. Symptoms start slowly and insidiously with episodic aching and stiffness of the axial musculature (paraspinal and abdominal muscles). Symptoms are usually symmetric and progress to involve the proximal muscles in all four extremities (Video 2.1). Typically the distal limb and facial

muscles are spared. Patients have a characteristic hyperlordosis of the spine, which makes it very difficult for them to bend over to touch their toes. The hyperlordosis persists even when they are laying down. The rigidity and stiffness may fluctuate on an hour-to-hour or daily basis. If it affects the neck, patients should be counseled not to drive until adequately treated, as it may significantly limit their ability to turn their heads.

Superimposed on the stiffness, patients also have intermittent severe spasms. These can be precipitated by various triggers, such as loud noise, sudden movement, touch, stress, and fatigue. Spasms usually last for minutes and abate once the offending stimuli are removed. However, during the spasms, patients can experience significant pain. The spasms can be variable in magnitude and severity, may occur in rapid succession, leading to a “spasmodic storm,” and can be severe enough to cause fracture of the long bones.

The fear of precipitating the spasms causes patients to have anxiety and task-related phobias. More than 50% of patients fear open spaces. The presence of phobias, excessive startle, and exacerbation of symptoms when emotionally upset many times leads to an erroneous diagnosis of a functional or psychogenic disorder and, unfortunately, delays proper treatment.

Electromyography (EMG) studies show the presence of continuous motor unit activity at rest without any abnormality in the motor unit morphology. Reflex-induced spasms are short-latency (<80ms), stereotyped motor responses to nerve stimulation. These are expressed as one or more hypersynchronous bursts of EMG activity followed by short pauses and then slow cessation.

Diagnosis of SPS variants and related conditions

SPS is a clinical diagnosis. Dalakas and colleagues have outlined criteria, which can assist the clinician in making the diagnosis (Box 2.2). However, there are patients who do not meet all of these criteria or have other additional features: they are often categorized as SPS variants, which we describe below (Box 2.3).

Box 2.1 Autoimmune antibodies in SPS and variants	
Presynaptic:	GAD and amphiphysin
Postsynaptic:	GABA-A receptor associated protein, glycine and gephyrin

Progressive encephalomyelitis with rigidity and myoclonus

Progressive encephalomyelitis with rigidity and myoclonus (PERM) variant of SPS (formerly, “stiff-person plus syndrome”) is a rare paraneoplastic